


UC-NRLF



B 4 091 732



UNIVERSITY OF CALIFORNIA  
MEDICAL CENTER LIBRARY  
SAN FRANCISCO



*Ex Libris*  
C. K. OGDEN



Digitized by the Internet Archive  
in 2007 with funding from  
Microsoft Corporation



HYMENOMYCETES BRITANNICI

“Pleniorē dabit lucē futura ætas.”

*Hymenomycetes Britannici*

---

BRITISH FUNGI

(*HYMENOMYCETES*)

BY

REV. JOHN STEVENSON

AUTHOR OF 'MYCOLOGIA SCOTICA'

HON. SEC. CRYPTOGAMIC SOCIETY OF SCOTLAND

WITH ILLUSTRATIONS

IN TWO VOLUMES

VOL. II. CORTINARIUS—DACRYMYCES

QK607

S84b

v.2

1886

WILLIAM BLACKWOOD AND SONS

EDINBURGH AND LONDON

MDCCCLXXXVI

172552



# HYMENOMYCETES.

---

## ORDER I.—AGARICINI.

GENUS V.—**Cortinarius** (*cortina*, a veil). Fr. Epicr. p. 255.

*Veil arachnoid, distinct from the cuticle of the pileus, superficial. Hymenophore continuous with the stem. Gills persistent, arid, changing colour, pulverulent with the spores which fall off slowly. Trama fibrillose. Spores globose or oblong, here and there apiculate, somewhat ochraceous on white paper. Growing on the ground in woods, putrefying. Fr. Hym. Eur. p. 335.*

The most natural of the genera of Agaricini, but not easily defined by artificial characters. The veil and the gills present the chief marks of distinction. The gills generally become cinnamon-coloured. Very rare or altogether wanting in hot countries, but generally abundant in northern woods. All autumnal. They present a very different appearance in different stages of growth, and in the various stages according as they are moist or dry. It is necessary, therefore, to examine species both in their earlier and later stages. Some *Flammulæ* resemble them, but can scarcely be confounded with them.

---

**Tribe I. PHLEGMACIUM** (φλέγμα, clammy moisture). Par- Phlegma-  
tial veil arachnoid. Pileus equally fleshy, *viscous*. Stem firm, cium.  
*dry. Fr. Hym. Eur. p. 336.*

† CLIDUCHII (*C. cliduchus*. κλειδοῦχος, κλείς, a club; ἔχω, to have). Partial veil superior, pendulous in the form of a ring from the apex of the nearly equal or clavate stem.

\* *Gills pallid then clay-colour.*

Phlegma-  
cium.

\*\* Gills violaceous or purplish then cinnamon.

\*\*\* Gills yellow, cinnamon, ferruginous (not at first whitish or violaceous).

\*\*\*\* Gills olivaceous.

†† SCAURI (*C. scaurus*, club-footed). Bulb depressed or turbinate, *marginate*. Stem fleshy, fibrous, *cortina* commonly *inferior*, arising from the margin of the bulb. Pileus equally fleshy. Gills somewhat sinuate.



XL. *Cortinarius (Phlegmacium) fulgens*. One-third natural size.

The margin of the bulb is a kind of circularly split volva. In many instances it has been observed conjoined with the covering of the pileus, in others free and volvaecous. Hence the *stem* is never sheathed with the universal veil, nor *at the first*, as in the next group, *extended*, and scarcely ever farinaceous or clothed at the apex, but like group \*\* of *Amanitæ* at first enclosed within the pileus and bulb, which are applied to each other like two small balls. The impression on the bulb around the stem arises from the margin of the pileus pressing upon it at first. There are a few species which are intermediate between *Scauri* and *Cliduchii*, the pileus at first embracing the stem to the very base, whence the bulb is nearly wanting, but the base itself is *marginate*.

\* Gills whitish, then clay-coloured or pale cinnamon.

\*\* Gills violaceous, dark blue, purplish, at length cinnamon.

\*\*\* Gills ferruginous, tawny or yellow.

\*\*\*\* Gills olivaceous.

††† ELASTICI. *Cortina* simple, thin, fugacious, medial or inferior. *Stem* at the first extended, somewhat thin, never marginato-bulbous or sheathed, but rigid-elastic, externally somewhat cartilaginous, polished, shining, naked at the apex, at length most frequently hollow. Pileus thin, often hygroph-anous.

\* Gills whitish, then clay-coloured or dirty-cinnamon.

\*\* Gills violaceous, purplish, flesh-coloured.

\*\*\* Gills pure ochre, tawny, or ferruginous.

\*\*\*\* Gills olivaceous, fuliginous.

## † CLIDUCHII.

\* Gills pallid then clay-coloured.

1. *C. triumphans* Fr.—Pileus 7.5–12.5 cent. (3–5 in.) broad, yellow with a tinge of brown or ochraceous when moist, yellow when dry, variegated at the disc with minute adpressed spot-like scales or wholly naked, fleshy, not very thick, convexo-plane, obtuse, regular, margin even, not bent inwards; flesh alike of pileus and stem compact, white. Stem 7.5–12.5 cent. (3–5 in.) long, 12 mm. ( $\frac{1}{2}$  in.) and more thick, solid, firm, attenuated from the ovato-bulbous base, striate, yellowish-white, adorned with tawny scales arranged in many circles (or rings); the partial *cortina* superior, woven, somewhat ringed. Gills emarginate, crowded,

6 mm. (3 lin.) broad, with a small decurrent tooth which ends determinately in a continuous mealy white zone, densely veined at the sides, plane, whitish or inclining to very pale bluish grey, at length clay-colour and somewhat cinnamon. Phlegma-  
cium.

Large, splendid, from its zones of scales recalling *A. Vittadinii*. The scales readily fall off. A smaller form occurs in dry birch woods.

In grassy ground. Rare. Sept.

Spores ellipsoid or oblong-ellipsoid,  $12-16 \times 5-6$  mk. *K.*; pruniform, punctate,  $12-13$  mk. *Q.* Name—from the stem resembling a triumphal column. *Fr. Monogr.* ii. p. 4. *Hym. Eur.* p. 336. *Icon. t.* 141. f. 1. *B. & Br. n.* 1263. *S. Mycol. Scot. n.* 445. *A. sublanatus* *Huss.* ii. t. 22.

**2. C. claricolor** Fr.—**Pileus** 7.5–10 cent. (3–4 in.) and more broad, yellow, unchangeable, wholly fleshy, convexo-flattened, and at length depressed, at the first everywhere, soon only round the margin veiled with superficial, silky-pruinose, villous down, then wholly smooth, even, *for the most part however broken into scales*. **Stem** solid, hard, white, wholly clothed as far as the superior cortina (above which it is white-mealy) *with white scaly flocci* or lax down, but when full grown the cortina and scales alike disappear, otherwise sometimes curt, bulbous, sometimes elongated and conico-attenuated or cylindrical; flesh of the pileus and stem white. **Gills** sometimes emarginate, almost free, sometimes adnate, crowded, at the first whitish then clay-colour, the edge unequal.

Very changeable in form, but always very robust and compact. Intermediate between *C. triumphans* and *C. turmalis*. In a var. the gills are at first slightly bluish-grey.

In mixed woods. Glamis, &c. Sept.–Nov.

Spores pruniform, nucleate,  $11-12 \times 6-8$  mk. *C.B.P.*;  $12 \times 6$  mk. *W.P.* Name—*clarus*, clear; *color*, colour. Of pure colour. *Fr. Monogr.* ii. p. 5. *Hym. Eur.* p. 336. *Icon. t.* 141. f. 2. *B. & Br. n.* 1541. *S. Mycol. Scot. n.* 446. *Quel. Grev. t.* 102. f. 1.

**3. C. turmalis** Fr.—**Pileus** yellow-tan, most frequently darker at the disc, not changeable, compact, convex then plane, very obtuse, even, smooth (sometimes obsoletely piloso-virgate), when young veiled with pruinose but very fugacious villous down, soon naked, viscid; flesh white. **Stem** sometimes 7.5 cent. (3 in.), sometimes 15 cent. (6 in.) long, 2.5 cent. (1 in.), thick, solid, very hard, rigid, *cylindrical*, here and there attenuated at the base, shining white when dry, *when young sheathed with a white woolly veil*, naked when full grown. Cortina entirely fibrillose, superior and persistent in the form of a ring, at length ferruginous with

Phlegma-  
cium.

the spores. **Gills** variously adnexed, rounded or emarginate, even decurrent with a tooth, crowded, *serrated*, white then clay-colour.

Densely crowded in large heaps. Dependent on the weather. The flesh is not so hard and compact as that of *C. claricolor*, wherefore *the pileus is never broken into scales*. The gills never turn bluish-grey. *The mealy and floccose covering is a universal veil*; in *C. triumphans* it is concentrically tawny-scaly, in *C. claricolor*, white-woolly-scaly, in *C. turmalis*, white-woolly, soon naked.

In mixed woods. Glamis, &c. Sept.-Oct.

Spores 8-9 × 5 mk. *K.*; 8 × 5 mk. *W.P.* Name—*turma*, a troop. From its habit of growth. *Fr. Monogr.* ii. p. 5. *Hym. Eur.* p. 336. *B. & Br. n.* 1774. *S. Mycol. Scot. n.* 447.

**4. C. sebaceus** Fr.—**Pileus** 6-12.5 cent. (2½-5 in.) broad, unicolorous, *pale*, of the colour of tallow, equally fleshy, convex then rather plane, commonly very repand, viscid, smooth, but at the first *covered over with a whitish pruinose lustre*; flesh white. **Stem** 7.5-10 cent. (3-4 in.) long, 12 mm.-2.5 cent. (½-1 in.) thick, solid, stout, compact, never bulbous, often twisted and compressed, slightly fibrillose, pale white. Cortina delicate, fugacious, adhering only to the margin of the pileus. **Gills** emarginate, *not crowded*, connected by veins, 8 mm. (4 lin.) broad, clay-colour or pallid cinnamon, paler at the sides.

The flesh of the pileus is not compact at the disc and abruptly thin at the circumference, but equally attenuated towards the margin. The flesh of the stem is white. The gills never turn bluish-grey. Taste mild.

In mixed woods. Glamis, &c. Sept.-Nov.

Spores pale cinnamon, *Fr.* Name—*sebum*, tallow. From the colour. *Fr. Monogr.* ii. p. 7. *Hym. Eur.* p. 337. *Icon. t.* 143. f. 1. *B. & Br. n.* 1542. *S. Mycol. Scot. n.* 448. *Grevillea, t.* 83.

**\*\* Gills violaceous, &c.**

**5. C. varius** Fr.—**Pileus** 5 cent. (2 in.) and more broad, bright *ferruginous-tawny*, compact, hemispherico-flattened, very obtuse, regular, slightly viscid, even, smooth, the thin margin at first incurved, appendiculate with the cortina; flesh firm, white. **Stem** curt, 4-6 cent. (1½-2½ in.) long, 2.5 cent. (1 in.) and more thick, *bulbous*, absolutely immarginate, compact, *shining white*, adpressedly flocculose, the superior veil pendulous. **Gills** emarginate, thin, somewhat crowded, *at first narrow, violaceous-purplish*, then broader and ochraceous-cinnamon, always quite entire.

Variable in stature, but the habit and colours are always unchangeable. It varies with the stem taller and somewhat equal, the pileus yellow-tawny, and the gills dark blue.



In woods. Uncommon. Sept.—Nov.

Phlegma-  
cium.

Name—*varius*, changeable. *Fr. Monogr.* ii. p. 8. *Hym. Eur.* p. 338. *Berk. Out.* p. 183. *C. Hbk. n.* 487. *Ag. Schæff. t.* 42.

**6. C. cyanopus** Fr.—**Pileus** 6 cent. (2½ in.) broad, at first date-brown-livid, then tan and opaque, fleshy when unfolded, flattened, obtuse, regular, viscid in rainy weather, in fine weather dry opaque and even, *smooth*. **Stem** 5–7.5 cent. (2–3 in.) long, 12 mm. (½ in.) thick, solid, firm, ventricoso-bulbous, but always immarginate, above the bulb attenuated upwards, *naked* (neither fibrillose nor mealy) and white, *violaceous above*. **Gills** at first adnate, then emarginate, broad (6–8 mm., 3–4 lin.), *not much crowded*, in the same cluster *intensely violaceous* and pallid bluish-grey, soon changing colour to cinnamon, edge unequal.

The pileus when young is hemispherical, not embracing the bulb but the apex of the stem, the cortina extending from the closely involute margin to the apex of the stem. The stem when young is ovato-bulbous, when full grown depressed and oblique (somewhat rhomboid) at the base, more than 2.5 cent. (1 in.) thick, soft and spongy. The gills are less crowded than those of its allies. The flesh of the stem in its earliest stage is dingy white, variegated purple-violaceous above, that of the pileus dingy *white*, when full grown wholly white.

In woods. Frequent. Sept.—Nov.

Spores 10 × 5 mk. *W.P.* Name—*κύανος*, dark blue; *πούς*, a foot. From the colour of the stem. *Fr. Monogr.* ii. p. 9. *Hym. Eur.* p. 338. *Berk. Out.* p. 184. *C. Hbk. n.* 488. *S. Mycol. Scot. n.* 449. *Quel. Grev. t.* 102. f. 2. *A. glaucopus* *Sow. t.* 223.

**7. C. largus** Fr.—**Pileus** 10–15 cent. (4–6 in.) broad, commonly date-brown-tawny, fleshy, compact at the broad disc, thin at the circumference, convexo-flattened, very obtuse, slightly viscid on the surface, adpressedly silky-fibrillose when dry, commonly rivuloso-squamulose, sometimes fibrillose towards the margin; flesh wholly fibrous, firm, *whitish-bluish-grey*, becoming white when exposed to the air. **Stem** *solid*, sometimes shorter, slightly bulbous, 6 cent. (2½ in.) long, commonly elongated, 10–12.5 cent. (4–5 in.) long, almost 2.5 cent. (1 in.) thick and equal, often *curved* and ascending, *wholly fibrillose*, the superior cortina *pendulous*, above which the stem is *pruinose*, white *tinted violaceous*; flesh same as that of pileus. **Gills** sometimes adnate, sometimes emarginate, crowded, 12 mm. (½ in.) broad, at first *bluish-grey-clay-colour*, then cinnamon.

Somewhat *cæspitose*. The pileus when young is sometimes violet such as is here and there observed round the margin of full-grown specimens. When bruised the stem is sometimes bloody.

Phlegma-  
cium.

In pine woods. Herefordshire. Sept.—Oct.

Name—*largus*, large. *Fr. Monogr.* ii. p. 10. *Hym. Eur.* p. 339. *Grevillea*, vol. viii. p. 76. *Quel. Grev. t.* 103. f. 1.

**8. C. Riederi** Fr.—**Pileus** 7.5 cent. (3 in.) broad, ochraceous, compact, campanulato-expanded, obtusely umbonate, even, *glutinous*, shining when dry; flesh watery. **Stem** 10–12.5 cent. (4–5 in.) long, solid, clavate, *lilac-silky* and tawny-fibrillose. **Gills** adnate, rather thick, eroded, lilac then cinnamon.

Pileus slimy when moist, obsoletely streaked. Allied to the *Myxacia*.

In pine woods. Herefordshire. Sept.—Oct.

Name—After Rieder. *Fr. Hym. Eur.* p. 339. *Grevillea*, vol. viii. p. 76. *Quel. Grev. t.* 104. f. 1.

\*\*\* *Gills yellow, cinnamon, &c.*

**9. C. saginus** Fr.—**Pileus** 10–12.5 cent. (4–5 in.) broad, yellow, fleshy, plano-convex, irregular, *repand*, viscous; flesh soft, white. **Stem** 7.5 cent. (3 in.) long, 2.5 cent. (1 in.) and more thick, solid, somewhat bulbous, fibrillose, *light yellowish*, *naked at the apex*. Cortina fibrillose, *fugacious*, not very conspicuous. **Gills** truly *decurrent*, 8–10 mm. (4–5 lin.) broad, attenuated at both ends, dingy pallid then cinnamon, *eroded* at the edge.

Very gregarious, somewhat cæspitose.

In woods. Hereford, &c.

Name—*sagina*, corpulence. From its size. *Fr. Monogr.* ii. p. 12. *Hym. Eur.* p. 340. *Grevillea*, vol. v. p. 56, vi. t. 92. *S. Mycol. Scot. Supp. Scot. Nat.* 1882, p. 216.

**10. C. russus** Fr.—**Pileus** 10 cent. (4 in.) broad, unicolorous, rufous, fleshy, convex then flattened, obtuse, viscid, smooth at the disc, *innately fibrillose round the margin*. **Stem** 7.5 cent. (3 in.) long, scarcely 2.5 cent. (1 in.) thick, stuffed then hollow, attenuated upwards, not bulbous, often curved-ascending, soft, adpressedly fibrillose, pale-white, delicately pruinose at the apex. Cortina very tender, fugacious. **Gills** obtusely adnate (scarcely perceptibly rounded), 8–10 mm. (4–5 lin.) broad, crowded, connected by veins, *rufous-ferruginous*.

Well distinguished by its *coppery-rufous pileus*, bitter nauseous taste, and by the soft *flesh being whitish-flesh-colour*.

In woods. Uncommon. Sept.—Oct.

Spores 8×6 mk. *W.G.S.* Name—*russus*, red. *Fr. Monogr.* ii. p. 13. *Hym. Eur.* p. 341. *C. Hbk. n.* 489. *S. Mycol. Scot. n.* 450. *Worth. Smith, Trans. Woolh. Cl.* 1870, t. 1.

\*\*\*\* *Gills olivaceous.*Phlegma-  
cium.

**11. C. anfractus** Fr.—**Pileus** 5-7.5 cent. (2-3 in.) broad, *dark olivaceous*, becoming fuliginous-tawny, equally fleshy, but at length very *irregular* at the thin bent in margin, even, smooth. **Stem** 5 cent. (2 in.) or a little more long, 12 mm. ( $\frac{1}{2}$  in.) and more thick, clavate, immarginate, ascending, fibrillose, very dingy, *violaceous at the apex*, here and there hollow within. Cortina superior, but not in the form of a ring. Flesh whitish, obsoletely violaceous at the apex of the stem. **Gills** emarginate, *distant*, 6-8 mm. (3-4 lin.) broad, *crisped, olivaceous-fuliginous*.

The pileus is always zoneless, but it varies fuliginous when moist, when dry yellowish-olivaceous, with tiger-spots. It is often pitted and depressed round the margin, which is at length lobed. It varies regular.

In woods. Rare. King's Cliffe, &c.

The whole plant when young is covered with a white volva. *M. J. B.* Spores  $6 \times 4$  mk. *W. P.* Name—*anfractus*, bent, waved, repand. Of the margin of the pileus. *Fr. Monogr.* ii. p. 14. *Hym. Eur.* p. 341. *Berk. Out.* p. 184. *C. Hbk. n.* 496. *Quel. Grev. t.* 104. f. 3.

## †† SCAURI.

\* *Gills whitish, then clay-coloured, &c.*

**12. C. multiformis** Fr.—**Pileus** unicolorous, light-yellow, clay-yellow, tawny, &c., fleshy, regular, convex then flattened, very obtuse and at length depressed, for the most part very viscous, even, smooth. **Stem** 5-10 cent. (2-4 in.) long, about 12 mm. ( $\frac{1}{2}$  in.) thick, solid, equal or attenuated, somewhat smooth or adpressedly fibrillose, naked (not pruinose), *white then yellowish*, with a somewhat marginate bulb. Cortina fibrillose, fugacious, *white*. **Flesh** *white*, at first compact then soft, that of the pileus at length slightly yellow. **Gills** emarginate, free or with a small decurrent tooth, very thin, crowded, tense and straight, not broad, serrulated, *at first white*, then clay-colour.

*Very changeable.* Odour and taste mild. It is distinguished by the *colour being by no means violet and not hygrophanous, by the white flesh, and by the crowded serrulated gills being at the first whitish.* According to the state of the atmosphere it is very viscid or somewhat dry, then often sprinkled with the universal white veil, the thin margin deflexed. The pileus often varies spotted, and when taller streaked with fibrils; moreover lacunoso-sulcate, &c. The bulb is about 2.5 cent. (1 in.) thick.

In woods. Uncommon. Aug.-Oct.

I have seen the bulb so margined as to give the impression of a volva like that of *A. pantherinus*. *M. J. B.* Spores ochraceous, *Fr.*; pruiniform, nucleate, punctate,  $5-6 \times 2-3$  mk. *C. B. P.*;  $7 \times 4$  mk. *W. P.* Name—*multus*,

Phlegma-  
cium.

*forma.* Very various in shape. *Fr. Monogr.* ii. p. 15. *Hym. Eur.* p. 343. *Berk. Out.* p. 184. *C. Hbk.* n. 491. *S. Mycol. Scot.* n. 451. *Quel. Grev. t.* 104. f. 4. *A. turbinatus* Sow. t. 102.

**13. C. napus** Fr.—**Pileus** 5–7.5 cent. (2–3 in.) broad, fuliginous, then date-brown-tawny, fleshy, convexo-plane, obtuse, even, smooth, glutinous, the regular *margin bent inwards*; flesh compact, white, with a horny line (hymenophore) next the gills. **Stem** 5 cent. (2 in.) long, 12 mm. ( $\frac{1}{2}$  in.) thick, solid, equal, ascending, smooth, firm, white, at length becoming yellow at the base, inserted in an obconic, acutely and obliquely marginate bulb. **Gills** emarginate, *somewhat distant*, broad, *crisped*, whitish-fuliginous, hyaline at the sides.

It differs conspicuously from species near to it in the *broad, crisped, somewhat distant, whitish-fuliginous gills*.

In pine woods. Hereford. Oct.

Name—*napus*, a turnip. From the bulb. *Fr. Monogr.* ii. p. 16. *Hym. Eur.* p. 343. *Grevillea*, vol. xii. p. 41.

**14. C. talus** Fr.—**Pileus** 5–7.5 cent. (2–3 in.) broad, of a yellowish dirty colour, becoming pale, margin somewhat olivaceous, yellowish, fleshy, thin, equal, convexo-plane, even, smooth, viscous; *flesh watery*, dingy pallid-whitish, *with hyaline spots, and variegated with a horny line next the gills*. **Stem** 7.5 cent. (3 in.) long, 12 mm. ( $\frac{1}{2}$  in.) thick, solid, equal, cylindrical (not oblique), becoming smooth, marginato-bulbous at the base, pale. **Gills** emarginate, somewhat crowded, *beautiful straw-colour* or ochrey-pallid, scarcely changing colour.

Very pretty, very much allied to *C. turbinatus*, but quite distinct in its colours. The bulb is small, somewhat round, marginate.

In woods. Epping Forest. Oct.

Name—*talus*, an ankle bone. From the swelling at the base of the stem. *Fr. Monogr.* ii. p. 17. *Hym. Eur.* p. 344. *Icon. t.* 145. f. 2. *Grevillea*, vol. xiv. p. 38.

**\*\* Gills violaceous, &c.**

**15. C. glaucopus** Fr.—**Pileus** dingy yellow, tan-tawny or clay-colour, very fleshy, *at first bent inwards*, then flattened and unequal, somewhat repand, slightly viscid, rarely even, commonly floccoso-scaly, often marked with a raised fuscous-blackish zone round the margin; flesh *compact*, white. **Stem** solid, very stout, at first in the form of a bulb (2.5 cent., 1 in.), and marginato-bulbous, fibrillose, woolly, chiefly at the apex, at length elongated (7.5 cent., 3 in.), the bulb vanishing, but the base remaining

marginate, more than 2.5 cent. (1 in.) thick, azure-blue-pallid; <sup>Phlegma-</sup>internally pale azure-blue, becoming yellow at the base. **Gills** rounded and emarginate, crowded, sometimes crisped, azure-blue, at length cinnamon.

Very changeable and variable. The above is the earlier form. Later it appears in great troops and is very robust. Pileus compact, very large, undulated and tiger-spotted, ochraceous or becoming tawny, when young regular and convex, 7.5-10 cent. (3-4 in.) broad, at length flattened and depressed, 20 cent. (8 in.) broad. Stem somewhat equal, marginate at the base, white or yellowish, fibrillose and striate, naked at the apex. *Flesh* alike of the pileus and stem *yellowish*, that of the stem dark yellow below. Gills sinuate, with a small decurrent tooth, 12 mm. ( $\frac{1}{2}$  in.) broad, somewhat crowded, bluish-grey then clay-colour and somewhat cinnamon.

In pine woods. Frequent. Aug.-Oct.

Spores sphaeroid-ellipsoid,  $8-9 \times 5$  mk. *K.* Name—*γλαυκός*, pale blue; *πούς*, a foot. With pale blue stem. *Fr. Monogr.* ii. p. 18. *Hym. Eur.* p. 344. *Berk. Out.* p. 184. *C. Hbk. n.* 492. *S. Mycol. Scot. n.* 452. *Quel. Grev. t.* 104. f. 5. *Ag. Schaff. t.* 53.

**16. C. calochrous** Fr.—**Pileus** tawny, yellow round the margin, *compact*, convex, then plane, obtuse, viscid, guttate, commonly stained with soil; margin short, regularly involute, when unfolded often flexuous; *flesh very white*. **Stem** *curt*, 4-5 cent. ( $1\frac{1}{2}$ -2 in.) long, solid, firm, *marginato-bulbous*, bulb very depressed, otherwise equal, fibrillose, yellowish, *never becoming dark blue*. Cortina marginal, fugacious, yellow. **Gills** emarginate, crowded, serrated, of a beautiful dark blue-purple.

In woods and pastures. Frequent.

Name—*καλός*, beautiful; *χρῶς*, colour. *Fr. Monogr.* ii. p. 19. *Hym. Eur.* p. 345. *Berk. Out.* p. 185. t. 12. f. 3. *C. Hbk. n.* 493. *Quel. Grev. t.* 105. f. 1. *S. Mycol. Scot. n.* 453.

**17. C. cærulescens** Fr.—**Pileus** 5 cent. (2 in.), and more broad, dingy yellow, somewhat clay-colour, but varying fuscous-yellowish, &c., equally fleshy, convex, at length plane, obtuse, regular, even, rather smooth, but often streaked with fibrils, viscid, when dry shining or opaque; *flesh violaceous, growing white*. **Stem** 5 cent. (2 in.) long, 12 mm. ( $\frac{1}{2}$  in.) thick (the bulb more than 2.5 cent., 1 in. thick), solid, equally attenuated upwards from the marginate bulb, firm, at first fibrillose, *of a charming violet colour*, then naked, becoming pale, whitish. Cortina fibrillose, fugacious. **Gills** somewhat adnate, slightly rounded, thin, very crowded, 4 mm. (2 lin.) broad, at first *beautifully* and intensely *azure-blue*, then purplish, at length dingy-cinnamon.

When young the plant is here and there wholly azure-blue. The bulb is here

Phlegma-  
cium.

and there obliterated with age, always white beneath. Neither the gills when bruised nor the flesh when broken ever change colour (intensely azure-blue), but the gills become purple and at length cinnamon. Odour scarcely any.

In mixed woods. Frequent. Aug.—Nov.

Spores subellipsoid,  $9-10 \times 5-6$  mk. *K.* Name—*cæruleus*, azure. *Fr. Monogr.* ii. p. 19. *Hym. Eur.* p. 345. *Berk. Out.* p. 185. *C. Hbk. n.* 494. *S. Mycol. Scot. n.* 454. *Quel. Grev. t.* 105. f. 3. *A. calochrous b. Letell. t.* 651.—*Schæff. t.* 34 (not good). *Vent. t.* 32. f. 1-3.

**18. *C. purpurascens* Fr.**—**Pileus** 10–12.5 cent. (4–5 in.) broad, bay-brown or date-brown-olivaceous, then tawny-olivaceous, tiger-spotted, and often depressed round the margin, which is at first inflexed then repand, marked with a raised fuscous zone, fleshy, compact at the disc, obtuse, repand, variable, smeared with dense gluten, but opaque when dry; *flesh wholly azure-blue*. **Stem** solid, thick, bulbous, wholly fibrillose, intensely pallid azure-blue, darker when touched, bulb somewhat marginate. **Gills** broadly emarginate, 6 mm. (3 lin.) and more broad, crowded, azure-blue-clay, then cinnamon, *violaceous-purple when bruised*.

Readily distinguished from species near to it (especially *C. glaucopus*) by its *gills* soon changing colour and becoming *spotted with purple* when touched. The primary form is curt, robust, very compact, juicy, azure-blue-purplish when touched. There is another form, in dense beech woods, somewhat caespitose, the longer stem somewhat twisted, pileus undulated, as much as 20 cent. (8 in.) broad, toast-brown then clay-colour and at length cinnamon with the spores. *Gills* at first bluish-grey-purplish; otherwise as above.

In mixed woods. Common. Sept.—Nov.

The marginal zone is not constant. *M. J. B.* Name—from the gills becoming purple when bruised. *Fr. Monogr.* ii. p. 20. *Hym. Eur.* p. 345. *Berk. Out.* p. 185. *C. Hbk. n.* 495. *S. Mycol. Scot. n.* 455. *Quel. Grev. t.* 105. f. 2.

\* ***C. subpurpurascens* Fr.**—**Pileus** tawny-fuliginous, becoming pale, not compact, obtuse, at length very much dilated and undulato-flexuous, viscous, obsoletely streaked with fibrils, somewhat spotted; flesh soft, whitish (very faint azure-blue), *un-changeable*. **Stem** 7.5–10 cent. (3–4 in.) long, 12 mm. (½ in.) thick, at length hollow, somewhat cylindrical, marginato-bulbous at the base, fibrillose below, naked above, smooth, slightly striate, pale azure-blue or whitish. **Gills** sometimes emarginate, almost free, sometimes decurrent, *pallid* then cinnamon, becoming purple when bruised.

In woods. Epping Forest. Oct. *Fr. Monogr.* ii. p. 21. *Hym. Eur.* p. 346. *Grevillea, vol. xii.* p. 42. *Batsch f.* 74.

\*\*\* *Gills ferruginous, &c.*Phlegma-  
cium.

**19. C. dibaphus** Fr.—**Pileus** 7.5–10 cent. (3–4 in.) broad, purplish, disc yellowish, and at length *variegated* with lilac, fleshy, convex then plane, at length depressed, somewhat repand, viscous, smooth; *flesh yellow, variegated under the pellicle with a violet line.* **Stem** 7.5 cent. (3 in.) long, 12 mm. ( $\frac{1}{2}$  in.) thick, marginato-bulbous, stuffed, fibrillose, yellow, shining, purplish at the apex, light yellowish within. **Gills** adnate, slightly rounded, somewhat crowded, 6 mm. (3 lin.) broad, quite entire, *purplish-ferruginous.*

Odour and taste mild.

In woods, beech and oak. Fordingbridge, Hants.

Spores  $8 \times 11$  mk. *W.G.S.* Name—*δι-βαφος*, twice dipped or dyed. Of two colours. *Fr. Monogr.* ii. p. 21. *Hym. Eur.* p. 346. *B. & Br. n.* 1265. *Saund. & Sm. t.* 10. *Quel. Grev. t.* 105. f. 4.

**20. C. turbinatus** Fr.—**Pileus** *unicolorous*, dingy yellow or green, *becoming pale*, hygrophanous, opaque when dry, fleshy, convex then flattened, obtuse, at length depressed, orbicular, even, *smooth*, viscid; *flesh soft, white.* **Stem** commonly curt (5 cent., 2 in.), but varying elongated, yellowish, springing from a globoso-depressed distinctly marginate bulb, otherwise equal, cylindrical, *stuffed then hollow.* **Gills** attenuato-adnate, thin, crowded, broad, *quite entire*, at first pallid light-yellowish, at length somewhat ferruginous.

The typical form is *regular*, distinct from its allies in the *hygrophanous pileus*, in the gills being *isabelline-ferruginous* and *quite entire*, and in being *without any dark blue or purple colour.* Easily distinguished by its turbinate bulb.

In woods. Uncommon.

Pileus variable in width. Margin of the bulb so strongly developed as to appear volvate like *A. pantherinus*. *M.F.B.* Name—*turbo*, a spinning-top; conical. Of the bulb. *Fr. Monogr.* ii. p. 22. *Hym. Eur.* p. 346. *Berk. Out.* p. 185. *C. Hbk. n.* 497. *Quel. Grev. t.* 107. f. 1. *Ag. Vent. t.* 38. f. 5. 6. *Bull. t.* 110.

**21. C. fulgens** Fr.—**Pileus** 5–7.5 cent. (2–3 in.) broad, orange-tawny, very fleshy, convexo-plane, obtuse, viscid, occasionally punctate as with drops, at length *silky-fibrillose* or squamulose; *flesh very compact, white-yellow, at length spongy and tan-colour.* **Stem** solid, curt, equal above the depressed and obliquely, at the first acutely, marginate bulb, woolly with the yellow, densely fibrillose (in wet weather viscid) cortina, when full grown elon-

Phlegma-  
cium.

gated to 7.5 cent. (3 in.). **Gills** emarginate, 6 mm. (3 lin.) broad, somewhat crowded, quite entire, at first bright yellow, at length somewhat distant, tawny or ferruginous with the spores.

Very handsome, robust. The colour of the pileus is deeply and persistently orange-tawny. The spores tinge even the wool on the stem.

In pine woods. Rare. Sept.

Spores  $9 \times 6$  mk. *W.G.S.* Name—*fulgeo*, to shine. Of its bright colour. *Fr. Monogr.* ii. p. 23. *Hym. Eur.* p. 347. *C. Hbk. n.* 498. *Saund. & Sm. t.* 12.

**22. C. fulmineus** Fr.—**Pileus** about 7.5 cent. (3 in.) broad, tawny, variegated with dense irregular agglutinated scales, very fleshy, at first hemispherical and attached to the bulb, then convex, very viscous, margin regular, at first involute; flesh thick, white, often yellow at the circumference or wholly yellowish. **Stem** when young enclosed in the bulb, apparently awanting, then extended but scarcely more than 2.5 cent. (1 in.), solid, obese, yellow, naked, white-cortinate at the apex. Bulb very depressed, marginate, rooting, wider than the young pileus. **Gills** rounded, thin, very crowded, golden-yellow at length tawny.

The bulb and the scales on the pileus resemble the volva of the *Amanitæ* of the group represented by *A. rubescens*. The whole plant is robust, very hard. Mild. The pileus is tawny, almost brown, margin orange. Stem not filamentous, at first slightly viscid.

In shady woods. Ledbury.

Name—*fulmen*, lightning. Shining. *Fr. Monogr.* ii. p. 23. *Hym. Eur.* p. 347. *Grev.* xii. p. 42. *Ag. Schæff. t.* 24.

**23. C. orichalceus** Batsch.—**Pileus** 7.5 cent. (3 in.) broad, normally blood-red or liver-rufescent, with the margin livid, or bay-brown-ferruginous, and cinereous-olive round the margin, fleshy, convex, soon flattened, at length depressed, with a viscous pellicle or glutinous, the disc spotted with scales. **Stem** 7.5 cent. (3 in.) long, 12 mm. ( $\frac{1}{2}$  in.) thick, solid, equal, but springing from a marginate somewhat volvaceous bulb, very fibrillose (the fibrils viscid in wet weather from the gluten of the pileus flowing down), yellow-green or dingy yellow-pale. **Gills** adnate, broad, crowded, sulphur-yellow becoming green.

The flesh alike of pileus and stem white, but lemon-yellow at the circumference, at length wholly greenish-yellow. Easily distinguished by its dark dirty colours, which are nevertheless variable (the pileus sometimes olivaceous). The gills are very broad behind.

Under trees. Lyndhurst, New Forest.

Name—*orichalchum*, yellow copper ore, or brass. Of the colour. *Fr.*



*Monogr.* ii. p. 24. *Hym. Eur.* p. 348. *Grevillea*, vol. xii. p. 42. *Batsch f.* Phlegma-  
184. *Quel. Grev. t.* 106. f. 1. cium.

\*\*\*\* *Gills olivaceous.*

**24. C. prasinus** Fr.—**Pileus** sometimes æruginous, sometimes tawny-fuliginous, *tiger-spotted*, as if scaly, compact, convexo-plane, very obtuse, regular, viscid, the margin regularly involute; flesh dingy white. **Stem** curt, very thick, solid, greenish-pallid, not becoming violet; *flesh greenish-white*. Cortina scanty, whitish or pallid-green. **Gills** emarginate, not crowded, when full grown rather *distant*, somewhat crisped, *yellow-olivaceous* or somewhat olivaceous, darker and cinereous-olivaceous at the base.

Commonly sporadic. Taste mild.

In beech woods. King's Cliffe.

Name—*πράσινον*, leek. Leek-green. *Fr. Monogr.* ii. p. 25. *Hym. Eur.* p. 348. *Berk. Out. p.* 186. *C. Hbk. n.* 500. *Quel. Grev. t.* 106. f. 2. *Ag. Schæff. t.* 218.

**25. C. scaurus** Fr.—**Pileus** commonly 5–7.5 cent. (2–3 in.), very rarely 10 cent. (4 in.) broad, tiger-spotted, of a peculiar tawny-fuliginous colour, when dry more tawny, fleshy, convex then plane, depressed when old, viscous, smooth, *the thin margin at length slightly striate*; flesh thin, soft. **Stem** solid, soft and spongy (especially the bulb) with age, equally attenuated upwards from the marginato-bulbous base (more than 2.5 cent., 1 in. thick at the base, 8 mm., 4 lin. at the apex), fibrillose or striate, azure-blue or olivaceous then becoming white, and also yellowish when old. Cortina fibrillose, somewhat green. **Gills** attenuato-adnexed, rounded, thin, *very crowded*, 2–4 mm. (1–2 lin.) broad, broader (6 mm., 3 lin.) in front, normally *purplish-olivaceous*, but varying pure olivaceous and even fuliginous.

Odour and taste mild. It differs from all neighbouring species in the very crowded and very narrow gills. The bulb here and there vanishes.

In woods. Rare. Oct.–Nov.

Stem purplish in my specimens with a red tinge on the edge of the bulb. *M. J. B.* Name—*scaurus*, with projecting ankles. Of the bulb. *Fr. Monogr.* ii. p. 25. *Hym. Eur. p.* 349. *Icon. t.* 146. f. 1. *Berk. Out. p.* 186. *C. Hbk. n.* 499. *S. Mycol. Scot. n.* 456. *Quel. Grev. t.* 107. f. 2.

††† ELASTICI.

\* *Gills whitish, then clay-coloured, &c.*

**26. C. cumatilis** Fr.—**Pileus** 7.5 cent. (3 in.) broad, *of a very charming violet* or purple-violet colour, convex, obtuse, not de-

Phlegma-  
cium.

pressed, often irregular, with a viscous pellicle, even, smooth; flesh both of pileus and stem hard, shining white, unchangeable. **Stem** 5-7.5 cent. (2-3 in.) long, 12 mm. ( $\frac{1}{2}$  in.) thick, but variable and often curved, *solid*, firm, wholly shining white, furnished with a cortina only at the apex, but not pulverulent; *the universal veil*, which serves as a pellicle of the pileus, *ruptured at the base and adnate to it, as a separable agglutinated membrane of the same colour as the pileus*. **Gills** attenuato-adnexed, almost free, crowded, narrow, scarcely 4 mm. (2 lin.) broad, with a small decurrent tooth, acute at the apex, *white* then clay-colour.

Handsome, distinguished, not allied to any. The colours of the pileus and base of the stem depend upon the universal veil, which on being ruptured forms the pellicle of the former, and the sheathing membrane of the latter. Yellow spots are left where these are rubbed off. It varies sometimes solitary and larger, compact, sometimes somewhat caespitose with the stems equal and ascending.

In woods. Honningham, Norfolk. Oct.

The colour agrees better with the name and with Fries's fig. than with his description. Spores  $7 \times 4$  mk. *W.P.* Name—*cumatilis*, sea-coloured, blue. *Fr. Monogr.* ii. p. 27. *Hym. Eur.* p. 349. *Icon. t.* 146. f. 2. *C. cærulescens* Saund. & Sm. t. 22.

**27. C. decoloratus** Fr.—**Pileus** 5-10 cent. (2-4 in.) broad, *clay-colour, disc darker*, thin, but equally fleshy, campanulate then convex, obtuse, soft, viscous, *soon dry*, smooth, corrugated when old; flesh soft, *white*, watery. **Stem** 7.5 cent. (3 in.) long, 12 mm. ( $\frac{1}{2}$  in.) or less thick, stuffed, thin, somewhat equal, slightly thickened at the base, ascending, fibrillose, *silvery*, often curved, smooth and *naked at the apex*. Cortina inferior, fibrillose. **Gills** emarginate, adnate or decurrent according to situation, not much crowded, 6 mm. (3 lin.) broad (broader than the flesh of the pileus), *clay-colour then cinnamon*.

Taste slightly acrid. Very protean. The above is the typical form found in beech woods. The habit and stature are quite those of *C. tabularis*, but the pileus is viscous, never silky-veiled, stem never scaly, &c. In birch woods: pileus a little darker, floccoso-scaly or rivuloso-granulose when dry; bulb at the first present, small and somewhat round, but soon vanishing, soft, stem attenuated from it, sometimes yellowish; gills at the first bluish-grey-whitish. In pine woods: smaller and thinner, pileus even; gills pallid clay-colour.

In woods. Epping. Edinburgh Fungus Show. Oct.

Name—*de*, and *coloro*, to colour. Changing colour. *Fr. Monogr.* ii. p. 30. *Hym. Eur.* p. 351. *B. & Br. n.* 1541. *S. Mycol. Scot. n.* 457. *Quel. Grev. t.* 107. f. 4.

\*\* *Gills violaceous, &c.*Phlegma-  
cium

**28. C. decolorans** Fr.—**Pileus** 4-6 cent. ( $1\frac{1}{2}$ - $2\frac{1}{2}$  in.) broad, *wholly persistently yellow*, fleshy, convex then flattened, somewhat gibbous, equal, even, smooth, viscous; *flesh thin, firm, white*. **Stem** 5-7.5 cent. (2-3 in.) long, 6-8 mm. (3-4 lin.) thick, stuffed, equally attenuated, *shining white*, smooth above the somewhat persistent medial cortina. **Gills** sinuato-adnexed, crowded, thin, 6 mm. (3 lin.) broad, *juiceless, purplish then soon cinnamon*.

The gills are unchangeable when bruised. Easily mistaken for a thin form of *C. varius*, differing, however, in the thinner unicolorous pileus, in the stem being more slender, externally rigid, polished, without a bulb, and smooth with exception of the cortina.

In fir woods. Epping Forest. Oct.

Name—*de*, and *coloro*, to colour. From the gills *changing colour*. *Fr. Monogr.* ii. p. 31. *Hym. Eur.* p. 351. *Grevillea*, vol. xiv. p. 38.

**29. C. porphyropus** Fr.—**Pileus** 4-7.5 cent. ( $1\frac{1}{2}$ -3 in.) broad, livid-light yellowish or clay-colour, unchangeable when touched, fleshy, very thin at the margin, convexo-plane, obtuse, even, innately streaked, viscid; flesh thin, soft, whitish, *soon becoming purple-lilac when broken*. **Stem** 5-10 cent. (2-4 in.) long, about 6 mm. (3 lin.) thick, stuffed, at length hollow, somewhat bulbous or rather equally attenuated from the thickened base (sometimes equal), fragile, *externally and internally violaceous-lilac*, soon becoming pale, even whitish, but *soon becoming violaceous-lilac again when touched*. Cortina inferior, fibrillose. **Gills** rounded or emarginate, somewhat crowded, rather broad (4-8 mm., 2-4 lin.) *purplish then watery cinnamon, becoming purple again when touched*.

It approaches *C. purpurascens*, but the gills are crowded in that species.

In woods. Coed Coch, 1871. Oct.

Name—*πορφύρεος*, purple; *πούς*, a foot. From the stem becoming purple when touched. *Fr. Monogr.* ii. p. 31. *Hym. Eur.* p. 351. *B. & Br. n.* 1351.

**30. C. croceo-cæruleus** Fr.—**Pileus** 2.5 cent. (1 in.), or a little more broad, *lilac* or faintly violaceous, fleshy but thin, convex then plane, obtuse or gibbous, even, smooth, viscous; *flesh watery, pallid*. **Stem** 5 cent. (2 in.) long, *hollow*, somewhat equal (sometimes attenuated downwards), 4-6 mm. (2-3 lin.) thick, even, smooth, naked, *fragile, whitish*. Cortina white, fibrillose, fugacious. **Gills** attenuated or broadly emarginate, with a small very thin decurrent tooth, plane, somewhat distant, lilac then clay-saffron.

Phlegma-  
cium.

Spores saffron, cinnamon. The flesh is unchangeable in colour, but otherwise the plant becomes very slightly purple when touched or broken.

In woods. Laxton Park, Norths. Oct.

Name—*croceus*, saffron; *cæruleus*, azure. *Fr. Monogr.* ii. p. 32. *Hym. Eur.* p. 352. *B. & Br. n.* 1543. *Ag. Pers. Ic. and descr. t.* 1. f. 2. *Bucknall, Brist. Nat. Soc. Proc.* vol. iv. t. 3. f. 3.

\*\*\* *Gills pure ochre, tawny, &c.*

**31. C. papulosus** Fr.—**Pileus** 6–9 cent. ( $2\frac{1}{2}$ – $3\frac{1}{2}$  in.) broad, honey-tan-colour at the circumference, *always darker* (ferruginous or fuscous) *at the disc*, fleshy, at first convex, obtuse, then plane and at length depressed (disc here and there gibbous), smooth, viscid; *the cuticle breaking up into minute, granular, fuscous patches when dry*; flesh white, thick at the disc, thin at the margin. **Stem** 7.5 cent. (3 in.) long, 1–2.5 cent. ( $\frac{1}{2}$ –1 in.) or a little more thick, solid, firm, wholly fleshy within, equal or thickened at the base, densely fibrillose, but quite *naked at the apex, white*. Cortina inferior, very fugacious, white. **Gills** *adnato-decurrent*, crowded, slightly joined behind, but at length separating from the stem in older specimens and forming a spurious collar, 4–6 mm. (2–3 lin.) broad, at the first pallid, soon *ochraceous*, at length very pale yellow-cinnamon.

There are forms with the stem longer, here and there flexuous and hollow.

In woods. Durdham Down, Bristol. Glamis. Oct.–Nov.

Name—*papula*, a pimple. From the minute granular patches on the pileus. *Fr. Monogr.* ii. p. 33. *Hym. Eur.* p. 352. *Bucknall, Brist. Nat. Soc. Proc.* vol. iv. t. 3. f. 1.

\*\*\*\* *Gills olivaceous, fuliginous.*

No British species.

Myxacium.

**Tribe II. MYXACIUM** ( $\mu\acute{\upsilon}\xi\alpha$ , mucus). Universal veil glutinous, hence the scarcely bulbous *stem is also viscosus*. Pileus fleshy, somewhat thin. Gills adnate, decurrent. (Certain *Phlegmacia* having the fibres of the cortina slightly viscid by gluten from the pileus must not be confounded with the *Myxacia*.) *Fr. Hym. Eur.* p. 354.

† COLLINITI (*C. collinitus*, besmeared). Stem floccoso-peronate, the flocci at first covered with gluten.

†† DELIBUTI (*C. delibutus*, besmeared). Veil entirely viscid, hence the *Myxaci*um. stem is not floccoso-peronate but only viscid, acquiring a varnished appearance when dry.

\* Gills whitish, then clay-coloured.

\*\* Gills at first violaceous, dark blue, or reddish.

\*\*\* Gills at first ochraceous or cinnamon.

### † COLLINITI.

**32. *C. arvinaceus* Fr.—Pileus** 7.5–10 cent. (3–4 in.) broad, orange-tawny, fleshy, soft, convex then soon flattened, at length reflexed and undulated, even, smooth, viscous (not glutinous), glistening when dry, margin slightly striate when in full vigour. **Stem** tall, as much as 20 cent. (8 in.) long, 12 mm. ( $\frac{1}{2}$  in.) thick, solid, equal, *silky-viscous*, never broken up into scales, *white*. Cortina soon fibrillose and fugacious. **Gills** *adnato-decurrent*, very broad (12–18 mm.,  $\frac{1}{2}$ – $\frac{3}{4}$  in.), even at the sides, *somewhat distant, crenulated at the edge, at first straw-colour then bright ochraceous*.

Spores plentiful, of the same colour as the gills. Cortina not in the form of a ring. Perhaps too nearly allied to *C. collinitus*.

In beech woods. Blaize Castle, Bristol. Oct.

Name—*arvina*, grease. Greasy, viscid. *Fr. Monogr.* ii. p. 36. *Hym. Eur.* p. 354. *Grevillea*, vol. xii. p. 42. *Bucknall, Brist. Nat. Soc. Proc.* vol. iv. t. 5. f. 1. *Ag. Krombh.* t. 73. f. 16–18.

**33. *C. collinitus* Fr.—Pileus** orange-tawny, *fleshy*, not compact, convex with the margin bent inwards then expanded, obtuse, clothed with a thick cuticle, which in wet weather swells into persistent *orange-tawny gluten*, shining when dry, margin even. **Stem** solid, firm, elongated, exactly *cylindrical*, at length soft, for the most part covered over with a *floccose* and glutinous veil, *which is commonly broken up into concentric scales*; near the apex the gluten is continuous with the glutinous cuticle of the pileus and forms an entirely viscous fugacious ring, colour variable, violaceous, white, even yellowish. **Gills** *adnate*, somewhat crowded, *whitish-bluish-grey or clay-colour then cinnamon*.

There are numerous forms departing from the type: b) pileus repand, stem inclining to fuscous: *Batsch* f. 197. c) small, pileus pallid yellow or straw-



XLI. *Cortinarius (Myxaci)um collinitus*. One-fifth natural size.

*Myxarium*. colour, stem concentrically light yellow-scaly, gills white. d) very small, pileus date-brown-fuscous, stem scarcely 2.5 cent. (1 in.)

In woods. Common. July–Nov.

Spores ferruginous, *Fr.*;  $9 \times 6$  mk. *W.G.S.*;  $12-14 \times 6-7$  mk. *K.* Name—*collino*, to besmear. From the gluten. *Fr. Monogr.* ii. p. 36. *Hym. Eur.* p. 354. *Berk. Out.* p. 186. *C. Hbk. n.* 501. *S. Mycol. Scot. n.* 458. *Ag. Bull. t.* 549, 596. *Ventur. t.* 32. f. 4–6 (abnormal). *Buxb. C. iv. t.* 9.

**34. *C. mucifluus* Fr.—Pileus** livid-clay, when dry tan, opaque, somewhat fleshy, campanulato-expanded, *smeared with separating hyaline gluten, margin striate*. **Stem** attenuated downwards, soft, viscid with the floccoso-scaly fugacious veil, white or inclining to azure-blue. **Gills** adnate, distinct, clay-colour then watery cinnamon.

*Odour sweet*. Intermediate between *C. collinitus* and *C. elatior*, so allied to the former that it was long considered a variety. It differs especially as follows: 1) stem spongy, attenuated downwards, white: 2) pileus thinner, campanulate then expanded, at length reflexed and repand, the membranaceous margin striate: 3) colour of pileus livid-clay, when dry tan, opaque: 4) the gluten of the pileus thin, hyaline, fluid, not forming a thick persistent brightly coloured pellicle: 5) odour sweet. Gills clay then cinnamon. There is nothing of a violaceous tint in the whole plant.

On the ground. Hereford, &c. Oct.

Name—*mucus, fluo*, to flow. From its being covered with mucus which flows off and disappears. *Fr. Monogr.* ii. p. 37. *Hym. Eur.* p. 355. *Icon. t.* 148. f. 1. *Grevillea, vol. xi. p.* 70. *Quel. Grev. t.* 108. f. 4. *S. Mycol. Scot. Supp. Scot. Nat.* 1881, p. 35.

**35. *C. elatior* Fr.—Pileus** about 7.5–10 cent. (3–4 in.) broad, commonly livid-light yellow when damp, dingy ochraceous when dry, slightly fleshy only at the disc, cylindrical or bullate then *campanulate*, afterwards flattened and somewhat reflexed, the disc above the stem obtuse, even, otherwise *membranaceous and longitudinally plicato-wrinkled at the sides*, fragile, slightly viscous. **Stem** 12.5–17.5 cent. (5–7 in.) long, 12 mm. ( $\frac{1}{2}$  in.) and more (5 cent., 2 in.) thick, commonly attenuated at both ends (especially at the base), longitudinally fibrous, clothed with somewhat viscous, whitish, torn up scales, rarely becoming violet. Cortina viscous, fugacious. **Gills** adnate, linear, at first 6 mm. (3 lin.) broad, at length broader (as much as 2.5 cent., 1 in.), somewhat distant, connected by veins or wrinkled at the sides, always dark, *brown-cinnamon*.

The colour of the pileus varies whitish, tan-fuscous, date brown, violaceous-brown, black, whitish round the margin, grey with the margin violaceous. Gills *violet-brown* in Sowerby's fig.

In mixed woods. Common. Sept.–Nov.

Name—*elatus*, tall. *Fr. Monogr.* ii. p. 37. *Hym. Eur.* p. 355. *Icon. t.* Myxaciium. 149. f. 1. *Berk. Out.* p. 186. *C. Hbk.* n. 502. *S. Mycol. Scot.* n. 459. *Sow. t.* 9.

**36. C. grallipes** Fr.—**Pileus** 7.5 cent. (3 in.) broad, ferruginous when moist, when dry ochraceous-tan, opaque, *almost membranaceous* with exception of the disc, *with a prominent* often acute *umbo*, campanulate then flattened, *even, hygrophanous*, slightly viscid (not glutinous). **Stem** 10–15 cent. (4–6 in.) long, 4–6 mm. (2–3 lin.) thick, stuffed then *hollow*, wholly equal, but flexuous, tough, *fibroso-striate*, viscous, *yellowish-tawny*, ochraceous when dry, naked at the apex. **Gills** wholly adnate, with a decurrent tooth, 12 mm. ( $\frac{1}{2}$  in.) broad, plane, attenuated in front, crowded, even at the sides, clay-colour then ferruginous.

The habit is that of *C. hinnuleus*, but it is viscous not torn.

Among grass near trees. Ashton Park, Bristol. Oct.

Pileus very slimy when wet; veil fugacious, cobweb-like, pale, whitish brown; spores dark ferruginous. *W.G.S.* Name—*grallæ*, stilts; *pes*, a foot. Long-stemmed as if on stilts. *Fr. Monogr.* ii. p. 38. *Hym. Eur.* p. 355. *Saund. & Sm. t.* 27.

**37. C. livido-ochraceus** Berk.—**Pileus** 2.5 cent. (1 in.) broad, livid-ochre, *somewhat membranaceous*, plane, viscid, *margin not striate*. **Stem** 2.5 cent. (1 in.) high,  $\frac{3}{8}$  thick in the middle, beautiful violet, ochraceous at the base, stuffed with cottony fibres, attenuated at either end, somewhat scaly, striate above the fugitive veil. **Gills** somewhat adnexed, broad in front, moderately distant, cinnamon, the extreme margin pale.

Pileus quite smooth, shining, covered with a thick subcartilaginous skin, the margin very thin; edge with a few indistinct fragments of the veil. Inodorous. Nearest to the smooth-stemmed form of *C. collinitus*.

In woods. Uncommon. Sept.–Oct.

An undoubted specimen has been gathered with the margin of the pileus striate. Spores lemon-shaped, 13–15 × 6–8 mk. *C.B.P.* Name—*livid-ochre* in colour. *Berk. Out.* p. 186. *Eng. Fl. v.* p. 89. *Fr. Hym. Eur.* p. 356. *C. Hbk.* n. 504. *S. Mycol. Scot.* n. 460.

†† DELIBUTI.

\* No British species. -

\*\* *Gills at first violaceous, &c.*

**38. C. salor** Fr.—**Pileus** grey, bright *violaceous* round the inflexed *margin*, at length of the same colour, when young obtusely conical or parabolic, soon campanulate and at length flattened with a broad umbo on account of the compactly fleshy disc, thin

**Myxadium.** towards the circumference, small in comparison with the stem, even, smooth, thinly viscous, fibrillose towards the margin when dried; flesh continuous, white, becoming yellow (or faintly azure-blue). **Stem** 4-7.5 cent. (1½-3 in.) long, 2.5 cent. (1 in.) thick at the base, 12 mm. (½ in.) at the apex, solid, *conico-attenuated from the bulbous base*, gradually elongated, white, but when young *covered to the apex with the azure-blue glutinous veil*, when old becoming pale, softer. **Gills** *adnate, distant*, distinct, 4-6 mm. (2-3 lin.) broad, quite entire, at first pale grey with the *edge violaceous* or bluish-grey, when full grown grey-clay-colour or cinnamon.

In proportion as the stem becomes more elongated, the bulb becomes less in size. The stem is white externally and internally, but when young it is blue with the glutinous veil. *Very distinguished, firm.*

In woods. Coed Coch.

Name—*salum*, sea. Sea-colour. *Fr. Monogr.* ii. p. 40. *Hym. Eur.* p. 357. *Icon.* t. 150. f. 1. *B. & Br. n.* 1878. *Quel. Grev.* t. 108. f. 1.

**39. C. delibutus** Fr.—**Pileus** 5-7.5 cent. (2-3 in.) broad, *light-yellowish*, fleshy, thin, especially towards the margin, convex then flattened, obtuse, at length somewhat depressed, viscid with hyaline gluten, and slightly silky-fibrillose when the gluten disappears. **Stem** 5-10 cent. (2-4 in.) long, 6-8 mm. (3-4 lin.) thick, *stuffed* or hollow upwards, *equally attenuated* from the slightly bulbous base or somewhat equal, elastic, viscous as far as the scanty, fibrillose, fugacious cortina, when dry even, smooth, varnished, yellowish white, snow-white at the apex. **Gills** *adnate*, at length rounded or slightly emarginate, more or less *distant*, 4-6 mm. (2-3 lin.) broad, *serrulated* pallid or often crisped at the edge, at the first dark blue, or *violaceous-dark blue*, then clay, cinnamon.

The pileus and especially the gills vary in the colour being more or less intense. Taste at first watery, then slightly pungent. It loves grassy and rather damp places.

On the ground. King's Lynn.

Spores 7×5 mk. *W.P.* Name—*delibuo*, to besmear. From the gluten. *Fr. Monogr.* ii. p. 41. *Hym. Eur.* p. 357. *Grevillea*, vol. viii. p. 97. *Quel. Grev.* t. 108. f. 2. *S. Mycol. Scot. Supp. Scot. Nat.* 1882, p. 216.

**40. C. illibatus** Fr.—**Pileus** 2.5-5 cent. (1-2 in.) broad, yellow, disc darker, *slightly fleshy*, campanulate then convex, at length plane and *somewhat umbonate*, with a viscous pellicle, even, smooth; flesh white, very thin at the circumference. **Stem** 7.5 cent. (3 in.) long, only 4 mm. (2 lin.) thick, *stuffed* then soon hol-



low, soft, slightly attenuated upwards, smooth, viscid, white, commonly with reddish dots upwards. Cortina superior, fibrillose, very fugacious. **Gills** adnato-decurrent, arcuate, 4 mm. (2 lin.) broad, thin, *crowded, quite entire, flesh-colour then clay, cinnamon* with the spores. Myxadium.

Much smaller and thinner than *C. delibutus*, &c., and without any violet colour.

In woods. Coed Coch.

Name—*illibatus*, unimpaired, entire. Perhaps from the gills. *Fr. Monogr.* ii. p. 42. *Hym. Eur.* p. 358. *B. & Br. n.* 1879.

\*\*\* *Gills at the first ochraceous or cinnamon.*

**41. C. stillatitius** Fr.—**Pileus** scarcely 5 cent. (2 in.) broad, slightly fleshy, convex then plane, somewhat umbonate, even, smooth, *clothed with azure-blue gluten*, which is at first continuous with that of the stem, *fuscous-livid when the gluten separates* in the form of drops, and at length grey-white; flesh soft, watery, hygrophanous. **Stem** 5 scarcely 7.5 cent. (2, 3 in.) long, 6–8 mm. (3–4 lin.) thick, *hollow, very soft*, wholly equally attenuated, at first sheathed with *thick azure-blue gluten* which is extended into the cortina, naked at the apex. **Gills** *emarginate*, scarcely crowded, distant rather, 6 mm. (3 lin.) broad, *dark cinnamon*.

Odour none. The gluten disappears with age. Rather allied to the *Col-liniti*, but the veil is by no means floccose. The gills are almost those of *C. elatior*.

Among dead leaves. Uncommon.

**Pileus** striate near the margin. Gills with a whitish ragged margin. *W.G.S.* Spores sphaeroid-ellipsoid, uniguttate, yellowish, 8 × 6 mk. *K.* Name—*stillo*, to drip. Dripping. *Fr. Monogr.* ii. p. 42. *Hym. Eur.* p. 358. *C. Hbk. n.* 503. *S. Mycol. Scot. Supp. Scot. Nat.* 1882, p. 216. *Saund. & Sm. t.* 3.

**42. C. pluvius** Fr.—**Pileus** 1–2.5 cent. (½–1 in.) broad, pale yellow-tawny when moist, ochrey-tan and opaque when dry, *slightly fleshy*, at first somewhat globose, then convex, commonly gibbous, when more fully grown and moist slightly pellucid-striate, hygrophanous, viscid and shining in rainy weather; flesh thin, of the same colour. **Stem** more or less elongated to as much as 7.5 cent. (3 in.), 4–6 mm. (2–3 lin.) thick, at first stuffed, then hollow, *soft, equal* or slightly attenuated upwards, even, naked or *obsoletely viscid* with whiter silky spots. Cortina white and *entirely fibrillose*, slightly covered over with slime, soon fugacious. **Gills** adnexed, separating, ventricose, crowded, light yellowish or at the first whitish, then ochraceous.

**Myxadium.** The pileus in the earliest stage is covered over, especially round the margin, with a very thin, silky, white film. Taste at first watery, then acrid and pungent.


In woods. Lea, Gainsborough. Glamis. Sept.-Oct.

Name—*pluo*, to rain. From being viscid in rainy weather. *Fr. Monogr.* ii. p. 43. *Hym. Eur.* p. 359. *B. & Br. n.* 1267. *S. Mycol. Scot. n.* 461. *Ag. Batsch f.* 190.

**Inoloma.**

**Tribe III. INOLOMA** (ἰσ, a fibre; λῶμα, a fringe). *Fr. Syst.*

*Myc. i.* p. 216. Pileus equally fleshy, dry, at first silky with innate scales or fibrils, flesh continuous, not hygrophanous. Veil single. Stem fleshy, somewhat bulbous. *Species very distinguished.* There are *Telamonias* which resemble *Inolomata*, but they differ from them by having the pileus smooth, moist, then torn (whereas the *Inolomata* become smooth), or by being sprinkled over with white superficial fibrils, and by possessing a double veil. *Fr. Hym. Eur. p.* 359.



**XLII.** *Cortinarius (Inoloma) calisteus.* One-fourth natural size.

\* *Gills at first white or pallid.*

\*\* *Gills, as well as the veil and stem, violaceous.*

\*\*\* *Gills or veil cinnamon, red, or ochraceous.*

(*C. subferrugineus* and *C. armillatus* are apt to be sought for here.)

\*\*\*\* *Gills or veil dark, fuscous, olivaceous.*

\* *Gills at first white or pallid.*

**43. C. argentatus** *Fr.*—**Pileus** 10 cent. (4 in.) broad, *silvery-shining*, disc becoming pale, at the first *silky-lilac round the margin* then dun-coloured, fleshy, convexo-plane, at length broadly gibbous, *silky-even* becoming smooth; flesh whitish. **Stem** 10 cent. (4 in.) long, 12 mm. ( $\frac{1}{2}$  in.) and more thick, solid, attenuated from the (scarcely bulbous) base, smooth, silvery-white, at length becoming yellow at the base, internally white. *Cortina* fibrillose, fugacious, adhering only to the margin of the pileus, pallid. **Gills** emarginate, crowded, slightly serrated, 6 mm. (3 lin.) broad, *pallid* then watery cinnamon.

*Odour weak.* There is a smaller form in pine woods. The stem is bulbous when short.

In woods. Epping Forest. Oct.

Inoloma.

In 'Hym. Eur.' Fries describes the smell as somewhat strong, but not foetid. Spores pruniform, dotted, 10-11 mk. *Q.* Name—*argentum*, silver. Silvered. *Fr. Monogr.* ii. p. 46. *Hym. Eur.* p. 360. *Grevillea*, vol. xiv. p. 38. *Ag. Krombh.* t. 2. f. 27.

**\*\* Gills, as well as the veil and stem, violaceous.**

**44. C. violaceus** Fr.—**Pileus** 7.5-15 cent. (3-6 in.) broad, dark violaceous, sometimes purplish-violet, truly fleshy, convex, then flattened, regular, obtuse, villous, the innate persistent villous down for the most part rimoso-squamulose, margin at first involute; flesh soft, deeper in colour. **Stem** 7.5-10 cent. (3-4 in.) long, 2.5 cent. (1 in.) thick, solid, stout, remarkably bulbous, spongy, soft, at first tomentose, then fibrillose, dark violaceous, internally violaceous-cinereous. Cortina woolly, azure-blue, then ferruginous with the shed spores. **Gills** somewhat adnate, firm, *distant*, connected by veins, broader than the flesh of the pileus, dark almost black violaceous, then coloured with the cinnamon spores, but again violaceous when these are rubbed off.

Inodorous. Very handsome. It is easily distinguished from its allies by the colour being always dark azure-blue externally and internally, by the villous-scaly pileus, and by the distant gills.

In woods and open ground. Frequent. Sept.-Nov.

Spores sphaeroid-ellipsoid, 12-13×7-8 mk. *K.* Edible, excellent, resembling *Agaricus campestris* in flavour. Name—*violaceous*, of a violet colour. *Fr. Monogr.* ii. p. 46. *Hym. Eur.* p. 360. *Sv. ätl. Sv. t.* 58. *Berk. Out.* p. 187. *C. Hbk.* n. 505. *S. Mycol. Scot.* n. 462. *Hussey i. t.* 12. *Ag. Sv. Bor. t.* 288. *Hedw. fil. Obs. t.* 4. *Bull. t.* 250?

**45. C. cyanites** Fr.—**Pileus** 7.5-12.5 cent. (3-5 in.) broad, at first dark blue, soon becoming pallid azure-blue or livid-fuscos, fleshy, soft, convex then flattened, obtuse, *silky, becoming even* and smooth, never floccoso-squamulose; flesh azure-blue then whitish, here and there reddish when broken. **Stem** 7.5-12.5 cent. (3-5 in.) long, more than 2.5 cent. (1 in.) thick at the base, 12-18 mm. ( $\frac{1}{2}$ - $\frac{3}{4}$  in.) at the apex, solid, very bulbous, fibrillose, azure-dark blue, lilac when touched, also azure-blue within, but reddish blood-colour on contact with the atmosphere, and *when compressed pouring out a watery vinous-reddish juice*. Cortina fibrillose, azure-blue. **Gills** rounded, *crowded*, thin, 6 mm. (3 lin.) and more broad, commonly *beautifully dark blue*.

A magnificent species. Var. *major*: pileus and stem compact, turning red more slowly, gills somewhat distant, cinereous-dark blue.

In wood. Reading.

Inoloma. Name—*κύανος*, dark blue. *Fr. Monogr.* ii. p. 47. *Hym. Eur.* p. 360. *Icon. t.* 152. *f.* 1. *B. & Br. n.* 1544.

**46. C. albo-violaceus** Fr.—**Pileus** 5–7.5 cent. (2–3 in.) broad, violaceous-white, fleshy, convex, *broadly umbonate* or rather gibbous, dry, not moist, beautifully *innately silky* and becoming even; flesh juicy, azure-blue-white. **Stem** 5–10 cent. (2–4 in.) long, solid, firm, *clavato-bulbous* or conico-attenuated, 1–2.5 cent. ( $\frac{1}{2}$ –1 in.) thick, externally and internally *white-violaceous*, white-villous, fibrillose above with the cortina, and often with the white *veil in the form of a zone* at the middle. **Gills** adnate, scarcely emarginate, 4–6 mm. (2–3 lin.) broad, somewhat distant, somewhat serrulated, of a peculiar *cinereous-violaceous* colour, at length cinereous-cinnamon with the spores.

Inodorous, insipid. The fibrils on the stem are longitudinally adpressed as in *Ag. geophyllus*. There is a larger form with the well-formed pileus obtuse.

In woods. Forres, &c. Sept.–Oct.

Spores sphaeroid-ellipsoid, 6–9 × 4–5 mk. *K.*; pruniform, 10 mk. *Q.* Name—*albus*, white; *violaceus*, violet. *Fr. Monogr.* ii. p. 49. *Hym. Eur.* p. 361. *Icon. t.* 151. *f.* 3. *S. Mycol. Scot. Supp. Scot. Nat.* 1883, p. 31. *Grevillea*, vol. xii. p. 55.

**47. C. camphoratus** Fr.—**Pileus** 5–7.5 cent. (2–3 in.) broad, at first lilac, then changing colour (becoming white, yellow, &c.) but not hygrophanous, truly fleshy, convex then flattened, obtuse, at first silky then becoming smooth; flesh azure-blue. **Stem** 7.5–12.5 cent. (3–5 in.) long, 1–2.5 cent. ( $\frac{1}{2}$ –1 in.) thick, solid, soft, bulbous or obclavate, when young with a woolly sheath, violet, *but internally white at the base*. Cortina fibrillose, azure-blue, at length cinnamon with the spores. **Gills** at first arcuate, commonly adnato-decurrent, but varying emarginate, thin, *crowded, at the first intensely azure-blue*, then becoming purple.

*Odour fætid and exceedingly penetrating, wholly peculiar*, and different from the goat smell of *C. traganus*.

In woods. Fine shade.

Spores ochraceous-cinnamon, *Fr.*; 9 × 6 mk. *K.* Name—*camphoratus*, strong-scented. *Fr. Monogr.* ii. p. 50. *Hym. Eur.* p. 362. *Icon. t.* 152. *f.* 2. *B. & Br. n.* 1128. *C. Hbk.*, n. 506.

\*\*\* *Gills or veil cinnamon, &c.*

**48. C. traganus** Fr.—**Pileus** about 7.5 cent. (3 in.) broad, very fleshy, convex then flattened, obtuse, dry, at first silky, becoming even, *lilac-purplish*, soon becoming pale, at length becoming smooth and yellowish internally and externally. **Stem** 7.5–12.5

cent. (3-5 in.) long, 1-2.5 cent. ( $\frac{1}{2}$ -1 in.) thick, the villous base *Inoloma*. however very bulbous, 2.5-5 cent. (1-2 in.) thick, solid, spongy, at first silky, then fibrillose, *violaceous then whitish, internally deep saffron-ochraceous*. Cortina continuous with the silky covering of the pileus, pallid violaceous, then cinnamon with the spores. **Gills** emarginate, broad, distant, distinct, firm, *saffron-ochraceous*, at length cinnamon, somewhat crenate at the edge.

Readily distinguished by its *very fætid odour, and by its thick distant gills being at the first saffron-ochraceous*.

In pine woods. Forres, &c. Aug.-Sept.

Spores sphaeroid-ellipsoid,  $8-10 \times 5-6$  mk. *K*. Name—*tragus*, a goat. With goat smell. *Fr. Monogr.* ii. p. 51. *Hym. Eur.* p. 362. *B. & Br. n.* 1499. *S. Mycol. Scot. n.* 463. *Ag. Schaff. t.* 56.

**Var. finitimus** Weinm.—**Pileus** silky, at length smooth, lilac, as is the stem, which is yellowish and mottled within, but not saffron-coloured nor brown. Smell not at all that of the typical form, but pleasant though peculiar, resembling that of gum just beginning to ferment.

Forres. Name—*finitimus*, nearly related. *Weinm. p.* 155. *B. & Br. n.* 1499. *S. Mycol. Scot. sub. n.* 463.

**49. C. tophaceus** Fr.—**Pileus** about 7.5 cent. (3 in.) broad, golden-tawny, opaque, not becoming pale, compactly fleshy, thin towards the margin, hemispherical, everywhere *villososquamulose*, varying slightly silky and shining; *flesh* soft, *white*, strong-smelling. **Stem** solid, bulbous, villososquamulose, tawny, the fibrillose veil of the same colour. **Gills** broadly emarginate, 12 mm. ( $\frac{1}{2}$  in.) and more broad, distant, of the same colour, at length tawny-cinnamon.

Commonly springing up in a dense band, somewhat caespitose, but also solitary and then larger, in *form, stature, and villososquamulose covering of pileus*, appearing like *C. violaceus*, only that species is of an entirely different colour. Odour not remarkable. It glitters from its *golden colour*.

In woods. King's Lynn.

Spores ellipsoid or subglobose, 6-8 mk. *C.B.P.*;  $7 \times 5$  mk. *W.P.* Name—*tophus*, tufa, a volcanic rock of an earthy texture. Of the colour. *Fr. Monogr.* ii. p. 52. *Hym. Eur.* p. 363. *Quel. Grev. t.* 109. f. 2.

\* **C. redimitus** Fr.—**Pileus** thinner, obtusely umbonate, shining, *golden, streaked with innate fibrils*; *flesh* white. **Stem** 2.5-5 cent. (1-2 in.) long, *slightly thickened at the base*, 12 mm. ( $\frac{1}{2}$  in.) thick, solid, internally soft, fibroso-striate, yellowish. **Gills** adnate, with a small decurrent tooth, light yellow then tawny.

In mixed wood. Glamis, 1875. Oct. Name—*redimio*, to wreath round,

Inoloma.

encircle. Perhaps of the fibrillose streaking. *Fr. Monogr.* ii. p. 52. *Hym. Eur.* p. 363. *Icon. t.* 153. f. 1. *B. & Br. n.* 1545. *S. Mycol. Scot. n.* 464.

**50. C. callisteus** Fr.—**Pileus** 5 cent. (2 in.) broad, *yellow-tawny*, not hygrophanous and not becoming pale, fleshy, *thin*, especially at the bent-in margin, convexo-expanded, scarcely umbonate, *rather smooth, silky towards the margin*, but commonly broken up into minute innate squamules; flesh yellowish-white. **Stem** 7.5–12.5 cent. (3–5 in.) long, clavato-bulbous, equally attenuated upwards, 12 mm. ( $\frac{1}{2}$  in.) and more thick at the base, solid, remarkably fibrillose or striate, *externally and internally firm and yellow, rhubarb-colour*, soft and spongy only at the base. Cortina marginal, fibrillose, fugacious, yellow-tawny. **Gills** wholly adnate, adnexed behind mutually and to the stem with flocci, plane, *somewhat distant*, distinct, 8 mm. (4 lin.) broad, thin, quite entire, yellow-tawny.

Odour none. *Very remarkable.* The figure of Saunders and Smith represents it admirably as regards stature and characters, but departs from it in the ferruginous colour.

In woods. Rare. July.

The colours are nearly those of *A. aureus*. *M. J. B.* Spores  $8 \times 6$  mk. *W. G. S.* Name—*κάλος*, beautiful. *Very beautiful.* *Fr. Monogr.* ii. p. 53. *Hym. Eur.* p. 363. *Berk. Out.* p. 187. *C. Hbk. n.* 507. *S. Mycol. Scot. n.* 455. *Saund. & Sm. t.* 3. *A. validus* Berk. *Eng. Fl.* v. p. 84.

**51. C. Bulliardi** Fr.—**Pileus** 5–7.5 cent. (2–3 in.) broad, *dark rufescent*, bay-brown-blood-colour becoming pale, fleshy, convex then expanded, obtuse, dry, commonly smooth, but also wholly fibrillose. **Stem** 5–7.5 cent. (2–3 in.) long, 12 mm. ( $\frac{1}{2}$  in.) thick, solid, fibrillose, variegated blood-red downwards as is also the ovate bulb, white at the apex. Cortina fugacious, whitish. **Gills** adnexed, with a decurrent tooth, 6 mm. (3 lin.) broad, somewhat crowded, purplish then ferruginous, crenulated at the (whitish) edge.

Flesh whitish, but that of the stem rufescent at the base. Readily distinguished by its *cinnabar bulb*. Stature small but firm. Strong smelling.

In mixed woods. Bristol, &c. Sept.

Name—after Bulliard. *Fr. Monogr.* ii. p. 53. *Hym. Eur.* p. 363. *Berk. Out.* p. 187. *C. Hbk. n.* 508. *S. Mycol. Scot. n.* 466. *Quel. t.* 9. f. 3 (departing from type). *Ag. Bull. t.* 431. f. 3.

**52. C. bolaris** Fr.—**Pileus** 2.5–5 cent. (1–2 in.) and more broad, light yellow-red, or pale, smooth or variegated with *innate, adpressed*, spot-like, *red scales*, rarely of one colour with continuous scales, fleshy, convexo-plane, obsoletely umbonate; flesh very

firm, white. **Stem** 5-7.5 cent. (2-3 in.) long, 6-10 mm. (3-5 lin.) *Inoloma*. thick, stuffed, sometimes hollow, hard, equal, pale, beautifully variegated with saffron-red, adpressed, fibrillose scales, white at the apex, sometimes continuously scarlet. *Cortina* fibrillose, fugacious, saffron-red. **Gills** commonly decurrent, arcuate, sometimes adnate, even emarginate, crowded, at first pale then dark cinnamon.

The stem varies straight, curved and elongated and flexuous. Very elegant. Sometimes solitary, sometimes somewhat caespitose.

In beech woods. Uncommon. Sept.-Nov.

Name—*βῶλος*, a clod; dim. *βωλάριον*; *βῶλινος*, made of clay, *i.e.*, of brick. Perhaps in allusion to the red colour. When young like a red clod. *Fr. Monogr.* ii. p. 54. *Hym. Eur.* p. 364. *Berk. Out.* p. 188. t. 19. f. 1. *C. Hbk.* n. 509. *S. Mycol. Scot.* n. 467. *Grevillea*, t. 79. *Ag. Pers. Ic. Pict.* t. 14. f. 1.

\*\*\*\* *Gills or veil dark, fuscous, olivaceous.*

**53. C. pholideus** Fr.—**Pileus** 5-10 cent. (2-4 in.) broad, fawn-colour, becoming pale, at length somewhat cinnamon, fleshy but thin, convex then flattened, somewhat umbonate and depressed round the umbo, *scaly* and squarrose *with innate, piloso-fasciculate, crowded, fuscous-blackish fibres*, which are fewer towards the margin; flesh pallid. **Stem** 7.5-10 cent. (3-4 in.) long, 6-12 mm. (3-6 lin.) thick, solid, attenuated upwards (sometimes shorter and clavato-bulbous), *scaly* as far as the somewhat annular *cortina* with fibrillose, crowded, somewhat concentric, fuliginous-black scales; above the universal *cortina* even and pallid violaceous, furnished with a spider-web *cortina* (double veiled). **Gills** somewhat emarginate, crowded, thin, 4-8 mm. (2-4 lin.) broad, *at first violaceous* then clay-colour, at length dark cinnamon with the spores.

Very distinct and very beautiful; readily recognised by the *gills being at the first violaceous, and by the pileus and stem being squarrose with fuliginous-blackish scales.*

In woods. Frequent. Sept.-Oct.

Spores subsphaeroid, 6-7 × 5-6 mk. *K.* Name—*φολίς*, a scale. *Scaly.* *Fr. Monogr.* ii. p. 55. *Hym. Eur.* p. 364. *Berk. Out.* p. 188. *C. Hbk.* n. 510. *S. Mycol. Scot.* n. 468. *Ag. Alb. & Schw.* t. 12. f. 1.

**54. C. sublanatus** Fr.—**Pileus** 7.5-10 cent. (3-4 in.) broad, fawn-colour, then becoming tan-fuscous, at length becoming ferruginous, slightly fleshy, campanulate then expanded, umbonate, clothed with *innate, floccose, fuscous squamules*; flesh comparatively thin, whitish. **Stem** 7.5 cent. (3 in.) long, solid,

Inoloma.

conico-elongated or clavato-bulbous, as much as 2.5 cent. (1 in.) thick at the base, clothed to the middle with fuscous down continued into a fibrillose cortina which does not form a zone, at the apex slightly violaceous, naked. **Gills** adnate, broader behind, scarcely crowded, 6 mm. (3 lin.) broad, *olivaceous-yellowish*, at length cinnamon.

The squamules on the pileus are less compact and not so much crowded as those of *C. pholideus*.

In woods. Rare. Oct.

Spores ovoid-pruniform, dotted, 10 mk. *Q*. Name—*lana*, wool. Somewhat woolly. *Fr. Monogr.* ii. p. 55. *Hym. Eur.* p. 364. *Berk. Out.* p. 188. *C. Hbk. n.* 511. *Ag. Sow. t.* 224. *Hussey ii. t.* 22.

**55. *C. arenatus* Fr.**—**Pileus** *pale yellowish-fuscous*, fleshy, convex, at first gibbous, *granulose with floccose squamules*. **Stem** clavato-attenuated, sheathed beyond the middle, fuscous-squamulose, even at the apex, pallid. **Gills** emarginate, ventricose, somewhat crowded, yellowish-cinnamon.

Closely related to *C. sublanatus*; when young also olive, not easily distinguished from forms of *C. pholideus* which have changed their colour, but the gills are never violaceous.

In woods. Uncommon. Aug.—Oct.

Spores 6-7 × 4-5 mk. *W.P.* Name—*arena*, sand. From the granulose pileus. *Fr. Hym. Eur.* p. 365. *Berk. Out.* p. 188. *C. Hbk. n.* 512. *S. Mycol. Scot. n.* 469. *Hussey i. t.* 72. *Ag. Bull. t.* 586.

**56. *C. penicillatus* Fr.**—**Pileus** 2.5 cent. (1 in.) or little more broad, ferruginous-fuscous, becoming tawny when dry, thin, slightly fleshy, convex, minutely umbonate, dry, *densely floccososcaly*, scales innate, dark, ferruginous-fuscous; flesh very thin, of the same colour as the pileus. **Stem** 5-7.5 cent. (2-3 in.) long, 4-6 mm. (2-3 lin.) thick, stuffed, *equal*, fragile, squamose almost to the apex with *adpressed, fuscous-ferruginous*, concentric *scales*, paler than the pileus, paler and adpressedly silky at the very apex. **Gills** separating, plane, somewhat crowded, 6 mm. (3 lin.) broad, dark brown.

Placed here on account of the innately squamulose pileus and its affinity with species in this section, but its stature is quite that of the *Dermocybæ*.

In pine woods. Cabalva. Sept.—Oct.

Spores pruniform, rough, 7-8 mk. *Q*. Name—*penicillum*, a painter's brush. Pencilled. *Fr. Monogr.* ii. p. 56. *Hym. Eur.* p. 365. *Grevillea, vol. viii. p.* 77.



**Tribe IV. DERMOCYBE** (δέρμα, skin; κῆβή, a head). **Pileus** Dermocybe. thinly and equally fleshy, at first silky with somewhat innate villous down, but becoming smooth when old, dry and not hygrophanous. Flesh watery when moist or coloured. **Stem** equal or attenuated, externally more rigid, elastic or brittle, internally stuffed or hollow. Veil single, fibrillose, forming a zone in *C. caninus*. A natural group, easily distinguished from *Inolomata* by the thinness and substance of the pileus and by the stem; and from the following subgenera by the pileus not being moist or hygrophanous, and by its short floccose or atomate covering, its brighter colour, &c. *Fr. Hym. Eur. p. 366.*

The species of this subgenus are very changeable, and not easily defined on account of the changeable colour of the gills. It comprehends two primary types; first that of *C. anomalus* with the flesh of the pileus white, and secondly that of *C. cinnamomeus* with the flesh scissile and coloured. *Fr. Monogr. ii. p. 57.*



XLIII. *Cortinarius (Dermocybe) cinnamomeus*. One-third natural size.

- \* Gills at first whitish or pallid.
- \*\* Gills at first violaceous, becoming purple.
- \*\*\* Gills brightly cinnamon, red, yellow. Stem and fibrillose cortina coloured. *Elegant.*
- \*\*\*\* Olivaceous, veil dingy pallid or fuscous. Pileus not torn into scales.

\* Gills at first whitish or pallid.

**57. C. ochroleucus** Fr.—**Pileus** 5 cent. (2 in.) broad, pale-white, fleshy, especially at the disc, broadly campanulate then expanded and somewhat gibbous, delicately and slightly silky, then becoming smooth, even; flesh firm, white. **Stem** 7.5 cent. (3 in.) long, 12 mm. (½ in.) thick, solid, ventricose, naked, white, fibrillose at the apex with the cortina. **Gills** broader behind, adnexed, then separating-free, 6 mm. (3 lin.) broad, crowded, clay-colour then ochraceous.

Odour none, taste remarkable, bitterish, not unpleasant. B. larger and more robust. Pileus 5–7.5 cent. (2–3 in.), somewhat obtuse, ochrey-white. Stem as much as 2.5 cent. (1 in.) thick, attenuated upwards. Gills adnate, pallid clay then cinnamon.

Dermocybe. In mixed woods. Rare. Oct.

Name—*ῥαῖος*, yellowish; *λευκός*, white. *Fr. Monogr.* ii. p. 57. *Hym. Eur.* p. 366. *Berk. Out.* p. 189. *C. Hbk.* n. 513. *S. Mycol. Scot.* n. 470. *Brig.* t. 31. f. 1-4. *Ag. Schæff.* t. 34.

**58. C. decumbens** Fr.—**Pileus** 2.5-4 cent. (1-1½ in.) broad, sometimes entirely *white*, sometimes yellowish, shining, by no means hygrophanous, fleshy, firm, convex then plane, gibbous then obtuse, even, smooth. **Stem** 2.5-5 cent. (1-2 in.) long, about 6 mm. (3 lin.) thick, stuffed, at length *hollow, ascending, clavato-bulbous* downwards, *smooth*, white. Cortina fugacious, white. **Gills** adnexed, crowded, 4 mm. (2 lin.) broad, thin, white then clay-colour, and at length ochrey-cinnamon with the spores.

Pileus silky-shining with a fibrillose pellicle, though smooth.

In woods. Epping.

Name—from the *decumbent* stem. *Fr. Monogr.* ii. p. 57. *Hym. Eur.* p. 366. *B. & Br.* n. 1775.

**59. C. tabularis** Fr.—**Pileus** 7.5 cent. (3 in.) and more broad, clay or fuscous-clay becoming pale, equally fleshy, not compact, convexo-plane, broadly gibbous, at length becoming remarkably plane, at first *veiled with* very thin, white *flocci*, silky near the margin, *then wholly smooth*; flesh white. **Stem** 5-7.5 cent. (2-3 in.) long, 6 mm. (3 lin.) and more thick, stuffed, at length hollow, tough, elastic, attenuated from the base or equal, erect, sometimes floccoso-scaly, sometimes even, somewhat smooth, white, becoming pale. Cortina white, fugacious. **Gills** somewhat *emarginate*, crowded, thin, 6 mm. (3 lin.) broad, whitish then clay-colour.

Sometimes the pileus is slightly concave, retaining water, and then the *flocci* are broken up and collected in the form of a zone round the margin. At the first the gills more rarely but slightly and very fugaciously incline to bluish-grey. A form has occurred with the gills at length ferruginous. Larger, paler, and smoother than *C. anomalus* to which it approaches.

In woods. Common. Sept.-Oct.

Name—*tabula*, a flat board or tablet. From the flat pileus. *Fr. Monogr.* ii. p. 58. *Hym. Eur.* p. 366. *Berk. Out.* p. 189. *C. Hbk.* n. 514. *S. Mycol. Scot.* n. 471. *Bull.* t. 431. f. 5.

**60. C. camurus** Fr.—**Pileus** 5-7.5 cent. (2-3 in.) broad, pallid fuscous-hoary, not hygrophanous, but becoming pale, pallid yellowish, umbo deeper in colour, very unequally fleshy, somewhat compact at the disc, but membranaceous almost to the middle, with a broad, obtuse, often oblique umbo, when dry rimosely incised; flesh white. **Stem** 7.5 cent. (3 in.) long, 8 mm. (4 lin.)

thick, somewhat hollow, equal, always ascending or twisted, even, smooth or fibrillose below, externally and internally white, naked and silvery-shining at the apex. Gills variously adnexed on account of the oblique pileus, sometimes broader behind and wholly adnate, sometimes almost free, crowded, 6 mm. (3 lin.) broad, thin, grey-clay-colour, then watery cinnamon and somewhat fuscous, quite entire and of the same colour at the edge. Dermocybe.

Odour somewhat rancid, but taste not remarkable. Different from all neighbouring species in being *remarkably fragile*.

In mixed woods. England. Glamis. Oct.

Name—*camurus*, crooked. Of the stem. *Fr. Monogr.* ii. p. 59. *Hym. Eur.* p. 367. *Icon. t.* 154. f. 1. *B. & Br. n.* 1546. *S. Mycol. Scot. n.* 472. *Bull. t.* 431. f. 4.

**61. C. diabolicus** Fr.—**Pileus** 2.5 cent. (1 in.) and more broad, at first *fuscous and crusted with grey*, then smooth and fuscous-yellowish, fleshy, thin, hemispherical, obtuse, then gibbous, dry, fragile, at length cracked. **Stem** 7.5 cent. (3 in.) long, commonly thin, 4 mm. (2 lin.) (but occurring as much as 10 mm., 5 lin.) thick, stuffed, attenuated downwards, smooth, pale, bluish-grey at the apex. Cortina fugacious. **Gills** adnate, separating, somewhat emarginate, somewhat crowded, 4 mm. (2 lin.) and more broad, firm, very pale bluish-grey then soon whitish, at length clay-colour.

Inodorous. A singular species, placed here on account of its colours. When old it might be readily taken for *C. raphanoides*.

In mixed woods. Uncommon. Aug.—Oct.

Name—"Called *diabolicus* because it holds a doubtful place among all the sections of this subgenus." *Fr. Fr. Monogr.* ii. p. 60. *Hym. Eur.* p. 367. *Berk. Out. p.* 189. *C. Hbk. n.* 515. *S. Mycol. Scot. n.* 473.

**\*\* Gills at first violaceous, becoming purple.**

**62. C. caninus** Fr.—**Pileus** 7.5–10 cent. (3–4 in.) broad, variable in colour, fuscous, brown, &c., equally fleshy, not compact, firm, not cracked, convex then becoming plane, obtuse, at first incrustated or hoary with silky villous down round the margin; when full grown *becoming smooth*; flesh white, at length somewhat yellowish. **Stem** 7.5 cent. (3 in.) and more long, 12 mm. (½ in.) thick, stuffed, spongy internally, at length hollow, thickened at the base, somewhat bulbous, often white-villous, pale-white, violaceous at the apex. **Gills** emarginate, somewhat distant, thin, 6–8 mm. (3–4 lin.) broad, bluish-grey or purplish then cinnamon.

The pileus is commonly *brick-rufescent* when old even becoming tawny

Dermocybe. when dry. The veil in the perfect state is double, the inferior one fibrillose or scaly clothing the greater part of the stem, and terminating near the apex in an annular (white or fuscous) zone, above which the stem is slightly villous and cortinate. Allied to *C. tabularis* as to stature, and distinct from *C. anomalus* in the stem being sheathed, veiled and adpressedly fibrous, somewhat annularly zoned above, and in the pileus being fuscous then rufescent.

In mixed woods. Common. Sept.—Nov.

Spores subsphæroid, 8–10 × 7–8 mk. *K.* Name—*canis*, a dog. Of doubtful meaning. Applied to various common plants. *Fr. Monogr.* ii. p. 60. *Hym. Eur.* p. 368. *Berk. Out.* p. 189. *C. Hbk.* n. 516. *S. Mycol. Scot.* n. 474. *Saund. & Sm.* t. 15. *Ag. Buxb.* C. 4. t. 22. *Bull.* t. 544. f. 1.

**63. *C. myrtilinus* Fr.—Pileus** 5–7.5 cent. (2–3 in.) broad, *pure fuliginous*, hoary-silky with dense fibrils, never rufescent, fleshy, thin, tough, gibbous then becoming plane; flesh when moist *watery fuscous*, when dry white, violaceous at the apex of the stem. **Stem** 5 cent. (2 in.) or a little more long, 6–8 mm. (3–4 lin.) thick, stuffed, externally tough, slightly bulbous, white-silky, whitish. Cortina scarcely manifest. **Gills** *adnate, somewhat distant*, of a beautiful *amethyst-azure-blue*, scarcely changing colour, never becoming purple.

The pileus is not at the first so convex as that of *C. anomalus*. The colour and habit are so like those of *Ag. nudus*, that those who say that that species is furnished with a veil seem really to have had this before them.

In mixed woods. Glamis, 1875. Rannoch. Sept.

Name—*myrtus*, myrtle. Of colour (*myrteus*), chestnut-brown. *Fr. Monogr.* ii. p. 61. *Hym. Eur.* p. 368. *B. & Br.* n. 1547. *S. Mycol. Scot.* n. 475. *Bolt.* t. 147.

**64. *C. anomalus* Fr.—Pileus** 2.5 cent. (1 in.) and more broad, fuliginous then rufescent, becoming hoary with separating fibrils, at length yellowish, but not hygrophanous, fleshy, thin, remarkably convex, then expanded and gibbous; flesh watery when moist, white when dry, not scissile. **Stem** 5–7.5 cent. (2–3 in.) and more long, 6 mm. (3 lin.) thick, stuffed then hollow, attenuated from the base, slightly sheathed, *fibrillose or somewhat scaly*, violaceous above, whitish below, of the same colour internally, at length becoming pale, somewhat yellow. **Gills** sometimes *adnate*, sometimes *emarginate*, with a decurrent tooth, crowded, thin; more or less violaceous, bluish-grey-purple, at length cinnamon.

Very thin, inodorous, more or less gregarious. Cortina coloured. There is a more slender variety in pine woods, with the stem thin, equal, smooth, and flexuous.

In woods. Common. Aug.—Oct.

The colour (even of the gills) is very changeable. Spores ellipsoid-sphæroid, Dermocybe. 8-9 × 6-7 mk. *K.* Name—*ἀ, ἀμαλός*, even. Irregular, anomalous. *Fr. Monogr.* ii. p. 62. *Hym. Eur.* p. 369. *Icon. t.* 154. f. 2. *Berk. Out.* p. 190. *C. Hbk. n.* 517. *S. Mycol. Scot. n.* 476. *Ag. Bull. t.* 431. f. 2.

**65. *C. spilomeus* Fr.—Pileus** 2.5 cent. (1 in.) broad, rufescent or clay-colour, not hygrophanous, somewhat fleshy, convex then expanded, gibbous, becoming smooth. **Stem** 5 cent. (2 in.) long, thin, 2-4 mm. (1-2 lin.) thick, fistulose, somewhat equal, white-lilac and beautifully *variegated with rufous or tawny scales*, furnished with a white cortina at the apex. **Gills** adnate or emarginate, crowded, narrow, thin, quite entire, bluish-grey or violaceous, becoming pale, at length watery cinnamon.

Very elegant, commonly cæspitose. The scales on the stem are like those of *C. photideus* but with the colour of those of *C. bolaris*.

In woods. Uncommon.

Spores subsphæroid, 8-9 × 7-8 mk. *K.* Name—*σπίλος*, a spot. From the variegated marking of the stem. *Fr. Monogr.* ii. p. 63. *Hym. Eur.* p. 369. *Icon. t.* 154. f. 3. *Berk. Out.* p. 190. *C. Hbk. n.* 518. *Ag. Brig. t.* 28. f. 4-6.

\*\*\* *Gills brightly cinnamon, &c.*

**66. *C. miltinus* Fr.—Pileus** 2.5-4 cent. (1-1½ in.) broad, bay-brown-cinnamon or when moist dark cinnamon, when dry shining, brick-colour, disc somewhat bay-brown, fleshy, thin, convex or lens-shaped then expanded, obtuse or broadly gibbous, even, polished-smooth, scissile at the disc; flesh watery when moist, tan when dry. **Stem** 5-7.5 cent. (2-3 in.) long, 4-6 mm. (2-3 lin.) thick, *fistulose*, very tough, somewhat cartilaginous, equally attenuated upwards, often twisted, white-tomentose at the base, *cinnamon* or reddish, *with red fibrils*, often villous at the apex with the red cortina. **Gills** adnate, 2-3 mm. (1-1½ lin.) broad, almost linear, plane, crowded, thin, beautiful reddish-cinnamon then ferruginous.

Odour none. Although the pileus is moist in rainy weather and becomes pale when dry, it cannot be called hygrophanous. An elegant and remarkable species, the type of the group. The habit is that of preceding species, but the colours are those of *C. cinnamomeus*.

In mixed woods. Ledbury.

Name—*μῖλτος*, red earth. From the colour. *Fr. Monogr.* ii. p. 64. *Hym. Eur.* p. 369. *Grevillea*, vol. xii. p. 42. *Quel. Grev. t.* 110. f. 3.

**67. *C. cinnabarinus* Fr.—Pileus** 5-7.5 cent. (2-3 in.) broad, *scarlet-red*, truly fleshy, campanulate, then flattened, obtuse or very obtusely umbonate, silky, then becoming smooth and shin-

Dermocybe. ing, or obsoletely scaly; the firm flesh paler. **Stem** 4-5 cent. ( $1\frac{1}{2}$ -2 in.) long, 6-8 mm. (3-4 lin.) and more thick, solid, equal, sometimes however bulbous, fibrillose or striate, *scarlet-red*, reddish-brick-colour internally. Cortina fibrillose, lax, cinnabar. **Gills** wholly adnate, somewhat decurrent, 6 mm. (3 lin.) broad, somewhat distant, connected by veins, unequal and darker at the edge, dark blood-colour when bruised.

Odour of radish. Readily distinguished from all others by its *splendid scarlet colour*, and from *C. sanguineus* by its short solid and firm stem, its broad pileus, and *somewhat distant gills*. Stem never becoming yellow.

In beech woods. Street, &c. Oct.

Name—*cinnabaris*, dragon's blood. From the colour. *Fr. Monogr.* ii. p. 64. *Hym. Eur.* p. 370. *Icon. t.* 154. f. 4. *S. Mycol. Scot. Supp. Scot. Nat.* 1882, p. 217. *Quel. Grev. t.* 110. f. 4.

**68. C. sanguineus** Fr.—**Pileus** 2.5-4 cent. ( $1-1\frac{1}{2}$  in.) broad, *blood-colour*, becoming slightly pale when dry, fleshy, thin, convex then plane, obtuse, occasionally depressed, silky or squamulose; flesh reddish, paler. **Stem** 5-7.5 cent. (2-3 in.) long, 4-6 mm. (2-3 lin.) thick, stuffed then hollow, equal (rather attenuated than thickened at the base), here and there flexuous, with fibrils of the same colour, almost darker than the pileus. Cortina arachnoid, fugacious, red blood-colour. **Gills** adnate, crowded, 4-6 mm. (2-3 lin.) broad, quite entire, dark blood-colour.

Wholly *dark blood-colour*, the stem when compressed pouring forth bloody juice. Odour of radish. Thinner than species nearest to it. The spores are ochraceous on a white ground, somewhat ferruginous on a black ground.

In woods, chiefly pine. Frequent. Sept.-Oct.

Spores  $6 \times 4$  mk. *W.G.S.* Name—*sanguis*, blood. From the colour. *Fr. Monogr.* ii. p. 65. *Hym. Eur.* p. 370. *Berk. Out.* p. 190. *C. Hbk. n.* 519. *S. Mycol. Scot. n.* 477. *Ag. Wulf. in Jacq. Coll.* 2. t. 15. f. 3. *Sow. t.* 43. *Bolt. t.* 36. *Klotsch Bor. t.* 385. *Krombh. t.* 2. f. 28-30.

**69. C. anthracinus** Fr.—**Pileus** dark *chestnut* or brown-fuscous, fleshy at the umbonate disc, otherwise thin, convex then expanded, umbo persistent, becoming fibrillose-even or smooth; flesh very dark, of the same colour as the pileus. **Stem** 5 cent. (2 in.) long, 4 mm. (2 lin.) thick, fistulose, equal, fibrillose, of an intense blood-colour, fuscous towards the base. **Gills** adnate, crowded, *deep red* or fiery in colour, blood-red when bruised.

Very elegant and most distinct; smaller in all its parts than those nearest to it.

In woods. Coed Coch.

Spores elongate-pruniform, 12-14 mk. *Q.* Name—*ἀσθραξ*, burning coal. Dermocybe. From the colour of the gills. *Fr. Monogr.* ii. p. 308. *Hym. Eur.* p. 370. *B. & Br. n.* 1269. *Quel. Grev. t.* 111. f. 1.

**70. C. cinnamomeus** Fr.—**Pileus** 1-10 cent. ( $\frac{1}{2}$ -4 in.) broad, somewhat cinnamon, fleshy, thin, obtuse, umbonate, silky or squamulose, with innate yellowish fibrils, at length becoming smooth. **Stem** stuffed then hollow, thin, *equal, yellowish as well as the flesh and veil.* **Gills** adnate, broad, crowded, shining.

There are innumerable forms sufficiently constant; the essential marks which are common to all are these: 1) *Pileus thin, when flattened obtusely umbonate, silky with yellowish villous down*, but when more fully grown often becoming smooth and then bright cinnamon, but the colour is variously changed. 2) *Stem wholly equal, stuffed then hollow, yellowish, fibrillose with the cortina, which is of the same colour.* 3) *Flesh scissile, yellowish.* 4) *Gills adnate, crowded, thin, broad, always shining.* 5) *Spores dark ochraceous.* The colour of the gills varies *blood-red, red cinnamon, saffron-tawny, golden, light yellow, &c.*

In mixed woods. Common. Aug.-Nov.

Name—*cinnamomum*, cinnamon. From the colour. *Fr. Monogr.* ii. p. 65. *Hym. Eur.* p. 370. *Berk. Out.* p. 190. *C. Hbk. n.* 520. *S. Mycol. Scot. n.* 478. *Ag. Krombh. t.* 71. f. 12-15. *Letell. t.* 618. *Brigant. t.* 30. f. 1-4. *Bolt. t.* 150. *Sow. t.* 205.

**71. C. uliginosus** Berk.—**Pileus** not exceeding 5 cent. (2 in.) in diameter, bright red-brown (almost *brick-red*), campanulato-conical, then expanded, very strongly umbonate, silky, sometimes streaked; flesh yellow-olive then cinnamon. **Stem** flexuous, paler than the pileus. **Gills** distant, adnate, with a tooth, yellow then olive, then cinnamon.

Remarkable for its very strong, but scarcely acute umbo.

In boggy woods among *Sphagnum*. King's Cliffe, &c.

Name—*uligo*, marshy ground. From its habitat. *Berk. Out.* p. 190. *C. Hbk. n.* 521. *S. Mycol. Scot. n.* 479. *Fr. Hym. Eur.* p. 371.

**72. C. orellanus** Fr.—**Pileus** orange-tawny, fleshy, obtusely umbonate, villosa-squamulose or fibrillose, *the flesh, which is similar in colour to the pileus, reddening.* **Stem** solid, firm, somewhat equal, striato-fibrillose and, as well as the *cortina, tawny.* **Gills** adfixed, broad, somewhat distant, at length opaque.

It differs from *C. cinnamomeus*: 1) *stem solid, firmer, striato-fibrillose, and, as well as the cortina, tawny;* 2) the ground colour of the firmer pileus also tawny, the fibrillose covering orange; 3) the similar flesh reddening; 4) gills broader, firmer, and somewhat distant. *C. cinnabarinus*, with which it has been confounded, is quite different in colours, and especially in that of the stem being the same as the pileus, not yellowish.

In mixed woods. Coed Coch, &c. Sept.-Oct.

Dermocybe. Spores pruniform,  $10 \times 4-5$  mk. *C.B.P.*;  $7 \times 4$  mk. *W.P.* Name—meaning not apparent. *Fr. Monogr.* ii. p. 66. *Hym. Eur.* p. 371. *B. & Br. n.* 1270. *S. Mycol. Scot. n.* 480. *Ag. Bull. t.* 598. Perhaps *Fungh. Linn. v. t.* 6. f. 19.

**73. C. infucatus Fr.**—**Pileus** 2.5–4 cent. (1–1½ in.) broad, yellow, not hygrophanous, fleshy, comparatively small, convex, obtuse, silky when dry, even; flesh whitish. **Stem** 7.5 cent. (3 in.) long, equally attenuated upwards from the clavate base (12 mm., ½ in. thick), 4–6 mm. (2–3 lin.) thick at the apex, solid, even, fibrillose, externally and internally pallid light-yellow. Cortina manifest, yellow, then darkened with the cinnamon spores. **Gills** adnate, crowded, thin, 2 mm. (1 lin.) broad, almost linear, tawny then cinnamon.

Remarkable for its bright yellow colour. It approaches *C. percomis* and *C. callisteus*; the former is readily distinguished by its viscid pileus and yellow flesh, the latter with more difficulty by its robust stature, by the pileus being smooth then slightly cracked with minute squamules, but especially by its firmer broad distant gills.

In woods. Haywood Forest, Hereford. Oct.

Name—*fusco*, to paint. From its bright colour. *Fr. Monogr.* ii. p. 309. *Hym. Eur.* p. 372. *Icon. t.* 155. f. 2. *Grevillea, vol. xii. p.* 42.

\*\*\*\* *Olivaceous*, veil dingy pallid, &c.

**74. C. cotoneus Fr.**—**Pileus** 7.5 cent (3 in.) broad, everywhere clothed with a *velvety*, dense, beautifully olivaceous covering, fleshy, but not compact, lax, campanulate then expanded, obtuse, somewhat repand, fragile when old; flesh thin for its breadth, soft, olivaceous-pale. **Stem** 7.5 cent. (3 in.) long, 12 mm. (½ in.) thick, solid, stout, soft, bulbous, somewhat fibrillose, olivaceous-pallid. *Veil* persistent, woven into a fuscous zone towards the apex of the stem. **Gills** adnate, separating, somewhat crowded, 4–6 mm. (2–3 lin.) broad, olivaceous then cinnamon, of the same colour and quite entire at the edge.

Handsome, soft to the touch, opaque.

In oak woods. Clifton.

Name—*κότινος*, wild olive. From the colour. *Fr. Monogr.* ii. p. 68. *Hym. Eur.* p. 372. *B. & Br. n.* 1960. *Quel. Grev. t.* 111. f. 5.

**75. C. raphanoides Fr.**—**Pileus** 2.5–5 cent. (1–2 in.) broad, fuscous-olivaceous, when fuller grown changing colour, becoming tawny, slightly fleshy, *campanulate then expanded*, obtusely umbonate, often undulated, *silky-fibrillose*, when fuller grown becoming smooth, flesh pallid, almost of the same colour. **Stem** about



5-7.5 cent. (2-3 in.) long, equally attenuated upwards, 6-10 mm. Dermocybe. (3-5 lin.) thick, rarely equal, stuffed, but in no wise furnished with a bark, fibrillose, opaque, *pale*. Veil filamentous, *pallid olive*. Gills adnate, slightly ventricose, scarcely crowded, somewhat olivaceous, at length cinnamon and somewhat ferruginous, edge paler.

Odour of radish, taste bitter. Protean like *C. cinnamomeus*. It varies with the stem elongated and twisted. The gills change colour with the state of the atmosphere, so that it is often difficult to recognise it.

In beech and fir woods. Frequent. Sept.-Oct.

Name—*raphanus*, radish. From the smell. *Fr. Monogr.* ii. p. 69. *Hym. Eur.* p. 373. *Berk. Out.* p. 191. *C. Hbk.* n. 522. *S. Mycol. Scot.* n. 481. *Mich.* t. 75. f. 2? *Quel. Grev.* t. III. f. 6.

**76. C. venetus** Fr.—**Pileus** 4-5 cent. (1½-2 in.) broad, when young green, then greenish-yellow, when dry more yellow, but not hygrophanous, fleshy, hemispherical, obtuse, equal, velvety or tomentose, the very thin erect down persistent; flesh yellowish-pallid. **Stem** 5-7.5 cent. (2-3 in.) long, 6-8 mm. (3-4 lin.) thick, stuffed or hollow upwards, equal, firm, curved, of the same colour as the pileus or a little paler, externally remarkably fibrilloso-silky, greenish-yellow. Cortina fibrillose, green. **Gills** adnate, connected by veins, somewhat distant, very broad, in the form of a segment, darker than the pileus, *olivaceous*.

Gregarious. *Most distinct*. The stem is often yellow-villous at the base. The yellow covering of the pileus readily distinguishes it from former species.

In woods. Rannoch. Oct.

Spores ovoid, punctate, 10 mk. *Q.* Name—*venetus*, sea-coloured. *Fr. Monogr.* ii. p. 70. *Hym. Eur.* p. 374. *Icon.* t. 155. f. 4. *B. & Br.* n. 1548. *S. Mycol. Scot.* n. 482.

**Tribe V. TELAMONIA** (τελαμών, lint). *Fr. Syst. Myc.* i. p. Telamonia. 210. Pileus moist, hygrophanous, at the first smooth or sprinkled with superficial whitish fibres of the veil. Flesh thin throughout or becoming so abruptly at the margin (not equally attenuated), scissile. *Stem annulate below from the universal veil or peronate with scales*, somewhat cortinate at the apex, hence the veil is somewhat double. *To this subgenus are referred some species, intermediate between this and the preceding one, on account of their double veil, the universal one forming a slight ring, and their pileus not being silky, &c.* *Fr. Hym. Eur.* p. 374.

Telamonia.


I. PLATYPHYLLI (πλατύς, broad; φύλλον, a leaf). Gills very broad, rather thick, more or less distant. Stem spongy and wholly fibrous.

\* Stem and cortina white or whitish.

\*\* Stem and gills violaceous. Cortina commonly white-violaceous, but universal veil white. Very distinguished.

\*\*\* Stem and veil reddish or yellow. Gills tawny or cinnamon, neither violaceous nor becoming brown.

\*\*\*\* Stem becoming fuscous, veil fuscous or dirty, gills dark-coloured. (Inolomata and Dermocybes are often sheathed and ringed, but the pileus is not hygrophanous, and is at the first scaly or silky.)



XLIV. *Cortinarius* (Telamonia) *armillatus*. One-fifth natural size.

II. LEPTOPHYLLI (λεπτός, thin; φύλλον, a leaf). Gills narrow, thin, more or less crowded. Pileus thin. Stem externally more rigid, somewhat cartilaginous, stuffed or hollow, often attenuated downwards.

\* Stem whitish, pallid, not floccoso-scaly.

\*\* Stem inclining to violet.

\*\*\* Stem and pileus tawny, ferruginous.

\*\*\*\* Stem floccoso-scaly, and, as well as the pileus, fuscous.

#### I.—PLATYPHYLLI.

\* Stem and cortina white or whitish.

77. *C. macropus* Fr.—Pileus 7.5 cent. (3 in.) broad, brick-colour, at length becoming ferruginous, fleshy, comparatively thin, convex then flattened, margin at the first inflexed, obtuse, dry, hoary with very small squamules, then becoming smooth; flesh cinereous. Stem 7.5–15 cent. (3–6 in.) long, 1–2.5 cent. ( $\frac{1}{2}$ –1 in.) thick, solid, somewhat equal, stout, fibrillose, dingy whitish, then of the same colour as the pileus. Veil distant from the pileus, woven into a narrow ring, white. Gills adnexed, very broad (1–2.5 cent.,  $\frac{1}{2}$ –1 in.), distant, even at the sides, sometimes crenated, sometimes quite entire, pallid then watery cinnamon.

Pileus not compact.

In woods. Leigh Wood, Bristol. Oct.

Spores pruniform, finely punctate, 8 mk. *Q.* Name—μακρός, long; ποῦς, a foot. Long-stemmed. *Fr. Monogr.* ii. p. 71. *Hym. Eur.* p. 374. *Bucknall, Brist. Nat. Soc. Proc.* vol. iv. t. 4.

78. *C. bivelus* Fr.—Pileus 5–10 cent. (2–4 in.) broad, tawny-ferruginous, here and there spotted or darker at the disc, not truly hygrophanous, somewhat equally fleshy, convexo-plane, always obtuse, soft, bibulous, smooth or slightly silky only round the

margin, commonly shining, rarely opaque, sometimes rimulose. *Telamonia*.

**Stem** solid, fleshy-spongy, sometimes bulbous, sometimes longer, equally attenuated, more frequently curt, thick, dingy white, internally somewhat ferruginous. The exterior veil villous, sheathing, white, terminating in a *spurious and fugacious ring*, above which the thin cortina is vanishing. **Gills** adnate or somewhat emarginate, at first somewhat crowded, then somewhat distant, more or less broad, bright tawny-cinnamon.

*The pileus is pierced as with pores*, a feature which is peculiar to most of the *Telamonie*, and which seems to have its origin in larvæ. Growing in troops, strong smelling, taste pleasant. The stem is in no wise more rigid externally; like that of the somewhat similar *C. subferrugineus*. There is a variety in which the pileus is campanulate and lax.

In woods. Coed Coch. Glamis. Sept.

Spores ellipsoid-sphæroid,  $8-10 \times 5-6$  mk. *K.* Name—*bis*, *velum*, with double veil. *Fr. Monogr.* ii. p. 72. *Hym. Eur.* p. 375. *Icon. t.* 156. f. 1. *B. & Br. n.* 1271. *S. Mycol. Scot. n.* 483. *Quel. Grev. t.* 111. f. 7.

**79. C. bulbosus** Fr.—**Pileus** about 7.5 cent. (3 in.) broad, *date-brown when moist*, fuscous-brick when dry, unequally fleshy, *campanulato-expanded*, obtuse or broadly *gibbous*, even, or *fibrillose-squamulose* towards the margin from the torn epidermis; flesh of the same colour when damp, only pallid, whitish when dry, more compact at the disc. **Stem** 5-7.5 cent. (2-3 in.) long, solid, bulbous, *paler than the pileus*, thinly sheathed and fugaciously ringed with the white veil. **Gills** adnate, somewhat distant, broad, at the very first dark, then brown-cinnamon, but never becoming violet.

The pileus is generally *smooth*, but here and there torn into fibrils towards the margin, like *C. armillatus* with which it is remarkably analogous. Stem somewhat saffron-yellow at the base and internally. It differs from the *Inolomata* in the hygrophanous pileus, and from the *Hygrocybes* in the gills.

In woods. Rare.

Name—*bulbus*, a bulb. From the bulbous stem. *Fr. Monogr.* ii. p. 73. *Hym. Eur.* p. 375. *Berk. Out.* p. 191. *C. Hbk. n.* 523. *S. Mycol. Scot. n.* 484. *Ag. Sow. t.* 130.

**80. C. urbicus** Fr.—**Pileus** 5 cent. (2 in.) broad, *clay-whitish*, fleshy, *convexo-plane*, obtuse, always *smooth*, but pitted when larger; flesh firm, whitish. **Stem** 5-7.5 cent. (2-3 in.) long, as much as 12 mm. ( $\frac{1}{2}$  in.) thick, solid, equal, zoned above the middle with a narrow white ring, when young villous above the ring. **Gills** emarginate, crowded, thin, broad, watery ferruginous.

**Telamonia.** Manifestly allied to *C. bivellus*, &c. The gills are not of the bright tawny colour of these. Easily distinguished from *C. bivellus* by its smoothness and colour; especially remarkable for the *clay-whitish colour* of the pileus.

In grassy places. Cabalva. Sept.-Oct.

Name—*urbs*, city. Found in frequented places. *Fr. Monogr.* ii. p. 72. *Hym. Eur.* p. 375. *Grevillea*, vol. viii. p. 77. *Quel. Grev. t.* 112. f. 1.

**81. *C. licinipes* Fr.—Pileus** 5–7.5 cent. (2–3 in.) broad, very pale yellow, tan-pallid when dry, fleshy-membranaceous, campanulate then convex and flattened, obtusely umbonate, and at length depressed round the umbo, even, smooth; flesh hygrophanous. **Stem** 10 cent. (4 in.) and more long, 6–8 mm. (3–4 lin.) thick, stuffed then hollow, at length fragile, equal, often flexuous, pale-white, white-villous at the base, otherwise clothed with *shining floccoso-plumose scales* which fall off with age, even above the distant membranaceous ring. **Gills** adnate and very broad behind, as much as 12 mm. ( $\frac{1}{2}$  in.), somewhat crowded, quite entire, watery cinnamon.

Fragile. Odour none. Var. short-stemmed, growing in drier places. *Bull. t.* 600. f. X. W. T.

In woods. Locality not recorded.

Name—*licinium*, lint; *pes*, a foot. Woolly-stemmed. *Fr. Monogr.* ii. p. 83. *Hym. Eur.* p. 376. *B. & Br. n.* 1549.

**\*\* Stem and gills violaceous, &c.**

**82. *C. torvus* Fr.—Pileus** 7.5–10 cent. (3–4 in.) broad, *brick-colour*, fleshy but thin with exception of the disc, convex, margin at first bent inwards, then wholly flattened, obtuse, somewhat fragile, even or broken up when dry, pierced with dotted points when old, at first *sprinkled with hoary squamules and fibrils*, and marbled with the same hoariness when it becomes smooth; flesh whitish when dry, dingy. **Stem** 7.5–12.5 cent. (3–5 in.) long, 1–2.5 cent. ( $\frac{1}{2}$ –1 in.) thick, solid, pierced by larvæ, and at length spongy, when young short and bulbous, then elongated and somewhat equal, when perfect *sheathed to the middle and ringed with the adnate veil*, above which it is more equal and pale violaceous, when young furnished with a villous cortina, then fibrillose; *below the white spreading ring whitish*, fibrillose and in fullest vigour floccoso-scaly, white-villous at the base. **Gills** somewhat adnate, *thick, distant*, plane, fragile, very broad (6–12 mm., 3–6 lin.), *at first violaceous, soon purplish-umber*, at length dark cinnamon, edge quite entire, at length veined at the base.

Striking, commonly robust. It is difficult to describe the ground colour of

the pileus; date-brown, brick-colour, copper-brown. There is a variety (in Telamonia. drier weather) with the gills rufescent-flesh-colour. Small forms changed in colour occur. The pileus like that of *C. scutulatus*, *elator*, &c. varies pitted-wrinkled and ribbed from the centre.

In mixed woods. Frequent. Aug.-Oct.

Name—*torvus*, savage. Growing in wild places. *Fr. Monogr.* ii. p. 74. *Hym. Eur.* p. 376. *Icon. t.* 157. f. 1. *B. & Br. n.* 1353. *C. Hbk. n.* 524. *S. Mycol. Scot. n.* 485. *Bull. t.* 600. f. *Q.R.S. Kalch. t.* 21. f. 1.

**83. *C. impennis* Fr.—Pileus** 5-10 cent. (2-4 in.) broad, *at first almost umber, then brick-colour*, changing colour and dingy, somewhat equally *fleshy*, very *obtuse, convex, smooth*, when young silky only round the margin, at length cracked; flesh pallid. **Stem** 5-10 cent. (2-4 in.) long, 1-2.5 cent. ( $\frac{1}{2}$ -1 in.) thick, solid, *cylindrical*, scarcely bulbous, *not clothed with scales*, pale, *becoming violet at the apex* and internally azure-blue, girt towards the apex with a white zone formed of the veil, and furnished with a cortina (of the same colour). **Gills** at first adnate, then emarginate, distant, rather thick, *at first intensely and brightly violaceous and somewhat purplish*, but immediately changing colour and becoming watery ferruginous.

The pileus *does not become hoary*. Approaching *C. torvus* in colours, but much inferior, certainly different *in the stem not being sheathed with the veil, in the pileus being smooth*, &c. In a young state the colour is almost that of *Ag. butyraceus*.

Among dead leaves. Bomere.

Name—*penna*, a feather. Without feathers. Smooth, contrasted with *C. plumiger*. *Fr. Monogr.* ii. p. 75. *Hym. Eur.* p. 376. *Icon. t.* 157. f. 2. *B. & Br. n.* 1880.

**84. *C. plumiger* Fr.—Pileus** fuscous, somewhat olivaceous when moist, brick-tan when dry, hygrophanous, slightly fleshy, *conical when young*, then campanulate, with a broad obtuse very prominent umbo, when more expanded as much as 7.5 cent. (3 in.), and often cracked, dry, clothed with *dense white floccoso-plumose scales*, which are sometimes erect and squarrose, sometimes (after showers) adpressed and silky. **Stem** 7.5-10 cent. (3-4 in.) long, 2.5 cent. (1 in.) thick at the base, 10 mm. (5 lin.) at the apex, solid, remarkably clavate, pale, internally watery whitish. *Veil floccoso-scaly on the stem, somewhat in the form of a ring at the apex, shining white*. **Gills** adnate, scarcely crowded broad, *at first violaceous*, soon of a watery then of a pure cinnamon, edge quite entire, of the same colour or clay.

Handsome.

Telamonia. In mixed woods. Glamis, 1875. Sept.

Name—*plumiger*, feather-bearing. From the scales. *Fr. Monogr.* ii. p. 76. *Hym. Eur.* p. 377. *B. & Br.* n. 1550. *S. Mycol. Scot.* n. 486. *Quel. Grev. t.* 112. f. 2.

**85. C. scutulatus** Fr.—**Pileus** comparatively small, at first *purple-umber* or brick-fuliginous, very hygrophanous, brick-colour when dry, somewhat fleshy, at first ovato-globose, then campanulato-hemispherical, obtuse, at first white-silky round the margin, then naked, but *rivulose in the form of innate squamules*, varying lacunose-wrinkled; flesh thin, violaceous. **Stem** sometimes 5–7.5 cent. (2–3 in.) long, 4–6 mm. (2–3 lin.) thick, sometimes 12.5–15 cent. (5–6 in.) long, 12 mm. ( $\frac{1}{2}$  in.) thick, solid and *rigid*, cylindrical or bulbous at the very base, *externally and internally intensely violaceous*, at length becoming fuscous, white-villous at the base, somewhat rooted, occasionally ascending, sheathed and somewhat ringed with the white veil. **Gills** adnate, rarely emarginate, more or less distant, 6 mm. (3 lin.) broad, violaceous then purple, white-serrated at the edge when young, at length cinnamon.

*Odour strong of radish.* Assuming various forms. *Veil white*; when in fullest vigour there is not only a *superior ring*, but there are also many zones from the floccose scales being arranged concentrically as in *C. triumphans*; commonly with a simple ring above and clothed with sparse fibrils, occasionally quite denuded. There are, moreover, variations with the stem hollow at the apex and the pileus umbilicate—with the pileus umbonate, more even, &c.

In moist woods. Foxley, Cabalva. Sept.–Oct.

Name—*scutula*, diamond, lozenge-shaped. From the markings on the pileus. *Fr. Monogr.* ii. p. 76. *Hym. Eur.* p. 377. *Icon. t.* 158. f. 2. *Grevillea*, vol. xii. p. 77.

**86. C. evernius** Fr.—**Pileus** 5–10 cent. (2–4 in.) broad, *purple-bay-brown*, brick-colour when dry and becoming isabelline-hoary when old, very hygrophanous, thin, almost *fleshy-membranaceous*, conico-campanulate then flattened, obsoletely umbonate, when young here and there adpressedly silky with the fibrillose white veil, but commonly smooth, *at length* rimosely incised and *torn into fibrils*, very fragile; flesh very thin, of the same colour. **Stem** 7.5–15 cent. (3–6 in.) long, 12 mm. ( $\frac{1}{2}$  in.) and more thick, stuffed, soft, *equal or attenuated downwards*, slightly striate, *violaceous*, becoming pale, *squamulose* and obsoletely zoned *with the adpressed veil*. **Gills** adnate, ventricose, *very broad* (1–2.5 cent.,  $\frac{1}{2}$ –1 in.), *distant, violaceous-purple*, becoming pale, at length cinnamon.

*Odour not remarkable. Changing colour very much.* It is analogous with

*C. elatior* in stature, but there is plainly no affinity between them from the *Telamonia*. absence of a glutinous veil, and from the very different colours.

In mixed woods. Frequent. Sept.—Oct.

Spores subsphaeroid-ellipsoid,  $8-12 \times 5-7$  mk. *K.* Name—*ἐκέρως*, (*ἐρως*, sprout) sprouting. Growing tall. *Fr. Monogr.* ii. p. 77. *Hym. Eur.* p. 377. *Berk. Out.* p. 191. *C. Hbk.* n. 525. *S. Mycol. Scot.* n. 487. *Sow.* t. 125 (departing very much from the type). *Quel. Grev.* t. 112. f. 3.

**87. *C. quadricolor* Fr.—Pileus** 5–7.5 cent. (2–3 in.) broad, pallid yellow, then somewhat tawny, shining when dry, slightly fleshy, conical then flattened, umbonate, smooth, at length pierced or spotted, radiato-striate at the margin; flesh thin, of the same colour. **Stem** 7.5 cent. (3 in.) long, thin, 4–6 mm. (2–3 lin.) thick, stuffed then hollow, equal, flexuous, slightly rigid, fibrilloso-striate with the adpressed veil, violaceous-whitish, the veil above collapsing in the form of a zoned oblique white ring, at length vanishing. **Gills** adnate, distant, broad, 6–8 mm. (3–4 lin.), but thin, white-serrated at the edge, sometimes dark violaceous, sometimes purplish then cinnamon.

Very much allied to *C. hinnuleus*, but the colours are different.

In woods. Coed Coch. Blaize Castle, Bristol. Oct.

Name—*quadricolor*, of four colours. *Fr. Monogr.* ii. p. 84. *Hym. Eur.* p. 378. *B. & Br.* n. 1663. *Bucknall, Brist. Nat. Soc. Proc.* vol. iv. t. 5. f. 2. *Ag. Schaff.* t. 303.

\*\*\* Stem and veil reddish or yellow.

**88. *C. armillatus* Fr.—Pileus** 7.5–12.5 cent. (3–5 in.) broad, red-brick-colour, truly fleshy, but not very compact, at first cylindrical, soon campanulate, at length flattened, dry, at first smooth, soon innately fibrillose or squamulose; flesh dingy pallid. **Stem** 7.5–15 cent. (3–6 in.) long, 12 mm. ( $\frac{1}{2}$  in.) thick, solid, firm, remarkably bulbous (bulb 2.5 cent., 1 in. thick, villous, whitish) and fibrillose at the base, when old striate and rufescent-pallid, internally isabelline. Exterior veil woven, red, arranged in 2–4 distant cinnabar zones encircling the stem; partial veil continuous with the upper zone, arachnoid, reddish-white. **Gills** adnate, slightly rounded, distant, at first pallid cinnamon, at length very broad (12 mm.,  $\frac{1}{2}$  in.), dark ferruginous, almost bay-brown.

Odour of radish. A very striking species. From the pileus not being hygrophanous, at the first smooth and at length torn into fibrils or squamulose, it might easily be taken for a species of *Inoloma*. The cortina itself is paler than the zones. It differs from all others in these zones.

In mixed woods. Frequent. Aug.—Oct.

*Telamonia*. The rings are usually somewhat oblique. Spores ellipsoid, 10-12 × 6-7 mk. *K.*; pruiniform, nucleate, rough, 10-12 × 5-7 mk. *C.B.P.* Name—*armilla*, a ring. Ringed. *Fr. Monogr.* ii. p. 78. *Hym. Eur.* p. 378. *Icon. t.* 158. f. 1. *Berk. Out. p.* 192. *B. & Br. n.* 1354. *C. Hbk. n.* 526. *S. Mycol. Scot. n.* 488.

**89. *C. hæmatochelis* Fr.**—*Pileus fuscous-brick-colour* (pallid), fleshy, thin, gibbous, silky-fibrillose. **Stem** solid, thickened but not bulbous, attenuated upwards, *encircled with a rufous zone*, becoming fuscous internally. **Gills** adnate, *crowded*, somewhat narrow, pallid cinnamon.

In woods. Coed Coch. Oct.

Name—*αἷμα*, blood; *χέλυς*, the chest. Girt with a red zone. *Fr. Hym. Eur. p.* 378. *B. & Br. n.* 1273, 1354. *Husseyi. t.* 19. *Ag. Bull. t.* 527. f. 1. *Paul. t.* 111.

**90. *C. limonius* Fr.**—*Pileus* 7.5-10 cent. (3-4 in.) broad, but often smaller, *tawny-lemon-yellow*, ochraceous yellow and opaque when dry, very hygrophanous, truly fleshy, convexo-plane, *obtuse*, even and *smooth* when moist, rimosely incised when dry; flesh of the same colour, not scissile. **Stem** 7.5 cent. (3 in.) long, 12 mm. ( $\frac{1}{2}$  in.) thick, fleshy, solid (but often pierced by larvæ), equal or sometimes attenuated, sometimes slightly thickened, *at the base*, where it is at length *deep saffron*, *floccoso-scaly with the light-yellow veil*, and here and there with a floccose ring at the apex, of the same colours as the pileus. **Gills** adnate, more rarely emarginate, distant, for a long time yellow to light-yellow, at length tawny-cinnamon.

Odour none. There is scarcely a manifest affinity with any other species than *C. psammcephalus*. The floccose ring is often wanting, but from the scales of the stem it is manifestly a species of *Telamonia*.

In pine woods. Rare.

Name—lemon-coloured (*Citrus Limonum*). *Fr. Monogr.* ii. p. 80. *Hym. Eur. p.* 379. *Icon. t.* 159. f. 1. *Berk. Out. p.* 192. *C. Hbk. n.* 527. *S. Mycol. Scot. n.* 489. *Quel. Grev. t.* 112. f. 4. *Ag. Holmsk. Ot. 2. t.* 40.

**91. *C. helvolus* Fr.**—*Pileus* 5-7.5 cent., 2-3 in. (in smaller specimens 2.5-5 cent., 1-2 in.) broad, dark tawny-cinnamon, very pale yellow when dry, slightly fleshy, *convexo-plane*, *obtuse* or obtusely umbonate, smooth, becoming even, *margin, which is at the first furnished with a cortina, regularly bent inwards*. **Stem** 5-20 cent. (2-8 in.) long, firm, solid, rarely pierced, somewhat equal, not tense and straight, fibrillose, girt above with an annular, narrow, ferruginous-margined zone formed of the woven veil. **Gills** *remarkably emarginate*, 8 mm. (4 lin.) broad, distant,



thick, somewhat veined at the base, opaque, tawny, then dark cinnamon. *Telamonia*.

The stem in the *largest* form is 20 cent. (8 in.) long, almost 2.5 cent. (1 in.) thick, attenuated at the base; *intermediate*, 7.5 cent. (3 in.) long, 8 mm. (4 lin.) thick; *smaller*, 5 cent. (2 in.) long, attenuated from the base, 4-6 mm. (2-3 lin.) thick at the apex. Flesh slightly tawny. Stem at length fuscous-ferruginous externally and internally.

In mixed woods. Coed Coch. Glamis. Sept.-Oct.

Name—*helvolus*, pale yellow. *Fr. Monogr.* ii. p. 85. *Hym. Eur.* p. 379. *B. & Br. n.* 1355. *S. Mycol. Scot. n.* 490. *Bull. t.* 531. *f.* 1 (except white flesh).

**92. *C. hinnuleus* Fr.—*Pileus*** pallid tawny-cinnamon, becoming pale, shining when dry, fleshy-membranaceous, campanulato-expanded, obtuse or obtusely umbonate, sometimes depressed in the centre, even, smooth, often pierced with dotted points; flesh of the same colour. **Stem** 2.5-10 cent. (1-4 in.) long, stuffed, but commonly pierced by larvæ, equal or *attenuated downwards*, rigid, dingy tawny and sometimes fuscous, but *white-silky with the adpressed silky veil* and white-zoned above with the woven veil. **Gills** more or less emarginato-adnexed, distant, but *thin*, the edge, which is of the same colour, acute and quite entire, 8-10 mm. (4-5 lin.) broad (semiovate when smaller), connected by veins, plane, saffron-tawny.

Varying much in stature. Stem sometimes short 2.5 cent. (1 in.) long, sometimes 5 cent. (2 in.) long, 4-6 mm. (2-3 lin.) thick, sometimes elongated 7.5-10 cent. (3-4 in.) and flexuous. It departs from all the others in this group in the *white veil*. The above is the earlier form in birch wood. *B.* later among poplar. Stem at length hollow, wholly fibrous, internally dark tawny, externally pale with the silky fibrils, ring obsolete; but *the pileus* (somewhat membranaceous, conico-expanded, acutely or obtusely umbonate, darker, yellowish when dry and at length rimosely torn) *is silky round the margins*. Gills adnate, distinct, opaque, slightly tawny then cinnamon. *Very small form*, stem 12 mm. (½ in.), gills very broad.

In mixed woods. Common. Sept.-Oct.

Name—*hinnuleus*, fawn. Fawn-coloured. *Fr. Monogr.* ii. p. 86. *Hym. Eur.* p. 380. *Berk. Out.* p. 192. *C. Hbk. n.* 528. *S. Mycol. Scot. n.* 491. *Sow. t.* 173. *Quel. Grev. t.* 113. *f.* 1.

**93. *C. gentilis* Fr.—*Pileus*** 1-2.5 cent. (½-1 in.) broad, tawny-cinnamon, *yellow when dry*, very hygrophanous, slightly fleshy, conico-expanded, then flattened, acutely umbonate, rimosely incised, varying somewhat silky; flesh thin and of the same colour. **Stem** 7.5 cent. (3 in.) and more long, 2-4 mm. (1-2 lin.) thick, stuffed then hollow, slender, equal or attenuated at the base, often curved, fibrillose, of the same colours as the pileus when moist and dry, never becoming pale. *Veil forming an annular zone*,

Telamonia. oblique, sometimes arranged in many zones, sometimes floccososcaly below the ring, *yellow*. **Gills** adnate, *thick, very distant*, plane, connected by veins, quite entire, tawny-cinnamon.

Gregarious. The stem is commonly more tense and straight than others of the same stature. The gills are of the same colour as the pileus in the primary state; they do not change colour when dry, like the pileus and stem, wherefore the fungus is remarkably of the same colour when moist, but when dry of different colours, the pileus and stem being yellow. Spores have been observed somewhat clay-coloured on a black ground.

In woods, chiefly pine. Frequent. Aug.-Oct.

Spores subsphaeroid-ellipsoid,  $8-9 \times 5-6$  mk. *K.*; pruniform, nucleate, slightly rough,  $10 \times 6$  mk. *C.B.P.* Name—*gentilis*, national, as opposed to exotic. *Fr. Monogr.* ii. p. 87. *Hym. Eur.* p. 380. *Icon. t.* 159. f. 2. *Br. Bath Transact.* 1870, p. 72. *C. Hbk. n.* 529. *S. Mycol. Scot. n.* 492. *Grevillea, t.* 84. f. 3.

**94. C. helvelloides** Fr.—**Pileus** 1-2.5 cent. ( $\frac{1}{2}$ -1 in.) broad, ferruginous, becoming tawny when dry, somewhat membranaceous, convex then flattened, umbonate (obtusely or acutely), smooth, somewhat striate when moist, when more fully grown cracked and squarrose. **Stem** 5-7.5 cent. (2-3 in.) long, 2-4 mm. (1-2 lin.) thick, fistulose, equal, very much *undulated* and flexuous, somewhat ferruginous and without any whiteness. *Veil yellowish, silky round the margin of the pileus*, and somewhat in the form of a ring *at the apex* of the stem, above which the stem is *white-silky* and glittering. **Gills** adnate, *very thick and exceedingly distant*, rather broad, distinct, *at first violaceous-umber* (never purplish), *edge white-floccose*, soon however of one colour umber-cinnamon, at length ferruginous.

Quite inodorous. The pileus varies fibrillose when young; sometimes bulate, rarely conical: it varies also in colour.

In mixed woods. Uncommon. Aug.-Oct.

Name—*Helvella-like*. *Fr. Monogr.* ii. p. 87. *Hym. Eur.* p. 380. *Icon. t.* 139. f. 3. *B. & Br. n.* 929. *C. Hbk. n.* 530. *S. Mycol. Scot. n.* 493.

\*\*\*\* *Stem becoming fuscous, &c.*

**95. C. punctatus** Fr.—**Pileus** hoary-umber (tan), somewhat membranaceous, conico-convex, smooth, at length even, pierced. **Stem** somewhat fistulose, equal, slender, undulated, fibrilloso-striate, *yellow-fuscous, girt with a pallid fuscous zone from the fugacious veil*. **Gills** adnate, very distant, quite entire, brown-cinnamon.

Primary form: pileus 12 mm. ( $\frac{1}{2}$  in.) broad; stem 5-7.5 cent. (2-3 in.) long, scarcely beyond 2 mm. (1 lin.) thick; gills almost 6 mm. (3 lin.) broad. Very

much allied to *C. gentilis*, distinct in the colour of all its parts, and in the Telamonia. convex, obtuse or obtusely umbonate pileus. Commonly also thinner, very flexuous, like *C. helvelloides*. In pine woods it occurs *larger*, somewhat cinnamon, pileus at length fibrillose, gills sinuato-adnexed and less distant.

In woods. Locality not recorded.

Name—*punctus*, a puncture. Pierced. *Fr. Hym. Eur. p. 382. Monogr. ii. p. 91. B. & Br. n. 1551.*

## II.—LEPTOPHYLLI.

\* *Stem whitish, pallid, &c.*

**96. C. triformis** Fr.—Pileus 5–7.5 cent. (2–3 in.) broad, *remarkably hygrophaneous, fawn-colour when young, then yellowish or honey-colour*, isabelline when dry, unequally fleshy, thin at the margin, convex then plane, obtuse or slightly gibbous, superficially fibrillose or becoming smooth, at length pierced-dotted, always even, *opaque*. **Stem** 7.5 cent. (3 in.) long, 12 mm. ( $\frac{1}{2}$  in.) thick, somewhat bulbous, stuffed, spongy internally, fragile, rather smooth, pallid, ringed upwards with the woven veil, *the ring distant and white*. **Gills** adnate, ventricose, somewhat emarginate, 8 mm. (4 lin.) broad, connected by veins, somewhat distant, *thin, watery honey-colour*, then watery cinnamon.

The pileus varies in colour, at first brownish or livid-yellowish, dingy tan when dry. The gills are of one colour.

In woods. Alresford, Hants.

Name—*tres*, three; *forma*, form. From its assuming three distinct forms, of which the above (*Schæfferi*) is the type. *Fr. Monogr. ii. p. 73. Hym. Eur. p. 382. Schæff. t. 247? Grevillea, vol. xiv. p. 38.*

\*\* *Stem inclining to violet, &c.*

**97. C. periscelis** Fr.—Pileus 5 cent. (2 in.) broad, *lilac, white-silky*, hygrophaneous, campanulate then convex, umbo fleshy, otherwise somewhat membranaceous. **Stem** 7.5–10 cent. (3–4 in.) long, 4–6 mm. (2–3 lin.) thick, equal, straight, fibrillose, of the same colour as the pileus, somewhat ringed with the *woven fuscous veil, becoming fuscous when dry*, white-villous at the base. **Gills** adnate, crowded, narrow, pallid then dark ferruginous.

Peculiar, rather perhaps belonging to the fuscous series.

In bogs and under beech. Bowood.

Name—*περισκέλις*, a garter. From the ring. *Fr. Hym. Eur. p. 383. Berk. Out. p. 193. C. Hbk. n. 531.*

**98. C. flexipes** Fr.—Pileus 1–2.5 cent. ( $\frac{1}{2}$ –1 in.) broad, at first

*Telamonia* very dark date-brown-fuscos, also inclining to violaceous, but becoming pale, very pale yellow in dry weather, tan when old, slightly fleshy, at first conical and acute, then expanded and acutely umbonate, at length depressed round the umbo, hoary-fibrillose, at length denuded, torn when old; flesh thin, of the same colour as the pileus. **Stem** 10 cent. (4 in.) long, 4 mm. (2 lin.) thick, stuffed, then pierced, equal, flexuous, *floccoso-scaly* below the woven sufficiently manifest white ring, pallid, *violaceous* throughout or at least *at the apex*. **Gills** adnate, *slightly distant, broad, purple or umber-violaceous*, then at length cinnamon, whitish at the edge.

When old almost like *C. evernius*, with which it agrees in the variation of colour.

In woods. Frequent. Sept.—Oct.

Name—*flexus*, bent; *pes*, a foot. From the flexuous stem. *Fr. Monogr.* ii. p. 84. *Hym. Eur.* p. 384. *S. Mycol. Scot. Supp. Scot. Nat.* 1881, p. 36. *Grevillea*, t. 113, f. 3.

**99. C. flabellus** Fr.—**Pileus** 18 mm. ( $\frac{3}{4}$  in.) broad, *olivaceous-fuscos*, tan when dry, somewhat membranaceous, commonly acutely umbonate, at first conical, then flattened, at first covered with white superficial separating scales, silky when dry, and at length rimosely incised, torn into fibrils; flesh very thin, paler. **Stem** 2–4 mm. (1–2 lin.) thick, sometimes short, 5 cent. (2 in.) long, in the typical form elongated (7.5–10 cent., 3–4 in.), stuffed then hollow, equal, undulated and flexuous, *floccoso-scaly*, pallid, becoming violet at the apex. *Veil white*, inferior, giving rise to the scales on the stem, terminating in a ring, which is sometimes perfect and entire, sometimes woven and oblique. **Gills** adnate, *crowded, linear, narrow, dark violaceous*, then cinnamon, at length ferruginous.

Growing in troops. *Odour strong*, somewhat of radish. Ring often wanting.

In woods. Coed Coch.

According to Fries's earlier description the umbo is commonly obtuse and vanishing. Name—*flabellus*, a small fan. Meaning not apparent. *Fr. Monogr.* ii. p. 85. *Hym. Eur.* p. 384. *B. & Br. n.* 1881.

\*\*\* *Stem and pileus tawny, ferruginous.*

**100. C. psammocephalus** Fr. *Wholly tawny-cinnamon*.—**Pileus** 2.5 cent. (1 in.) broad, fleshy, thin, convex then plain and at length umbonate and revolute, *broken up into minute furfuraceous squamules*. **Stem** 2.5 cent. (1 in.) long, 4 mm. (2 lin.) thick, stuffed, at

length hollow, somewhat attenuated, *sheathed with the continuous Telamonia. squamulose veil*, from which proceeds a fibrillose cortina. **Gills** sinuato-adnate, crowded, 4 mm. (2 lin.) broad, at length darker, umber-cinnamon.

Odour none. Internally it is of the same colour (not white as in Bulliard's fig.); the pileus and stem become pale and somewhat golden when dry. No distinct ring.

In mixed woods. Frequent. Sept.—Oct.

Name—*ψάμμος*, sand; *κεφαλή*, head. From the furfuraceous squamules on the pileus. *Fr. Monogr.* ii. p. 80. *Hym. Eur.* p. 384. *Berk. Out.* p. 193. *C. Hbk. n.* 532. *S. Mycol. Scot. n.* 494. *Bull. t.* 531. f. 2.

**101. C. incisus** Fr.—**Pileus** *tawny-ferruginous, opaque*, slightly fleshy, variable in form, sometimes very acutely, sometimes obsoletely umbonate, *convexo-expanded*, naked when young, *then* (especially in dry weather) *torn into fibrils or scales*, but even and shining when scorched by the sun. **Stem** sometimes curt, 2.5 cent. (1 in. long), 2–4 mm. (1–2 lin.) thick, sometimes elongated 5–10 cent. (2–4 in.), somewhat stuffed, flexuous, wholly fibrous in moist places, fibrillose, not polished externally. **Gills** adnate, *somewhat distant*, cinnamon-ferruginous.

Not much smell. Growing in troops, somewhat *cæspitose*. The pileus is commonly tawny-ferruginous, but when young also date-brown, olivaceous-fuscous, &c. It differs from *G. gentilis*, &c., in the colour being darker, ferruginous, and especially in the somewhat zoned *white veil*.

In woods. Loughborough.

Spores ellipsoid or sphaeroid-ellipsoid, uniguttate, 9–12 × 6 mk. *K.* Name—*incido*, to cut into. From the cracked pileus. *Fr. Monogr.* ii. p. 88. *Hym. Eur.* p. 384. *Icon. t.* 160. f. 1. *B. & Br. n.* 1272. *Ag. Bull. t.* 586. f. 2.

**102. C. iliopodius** Fr.—**Pileus** 2.5–5 cent. (1–2 in.) broad, cinnamon, tan when dry, slightly fleshy, conical then expanded and for the most part acutely umbonate, at the first silky with hoary fibrils, becoming smooth when full grown. **Stem** here and there short, commonly elongated, stuffed, flexuous, *equal*, elastic, sheathed to the middle with the white veil which becomes even and silky, and cortinato-ringed where the sheathing ends, naked and fibrilloso-striate at the apex, internally saffron-cinnamon. **Gills** adnate, slightly distant, thin, cinnamon.

Very protean, and there are individual forms which can scarcely be defined. It is distinguished, however, from others by its *opaque cinnamon colour*, that of the stem often becoming fuscous, by the *stem being externally somewhat cartilaginous and paler*, by the *white veil*, and by the *gills being more crowded and thin*.

In mixed woods. Common. July–Nov.

Telamonia.

Name—*ἰλὺς*, mud; *ποὺς*, a foot. From the dirty stem. *Fr. Monogr.* ii. p. 88. *Hym. Eur.* p. 385. *Berk. Out.* p. 193. *C. Hbk.* n. 533. *S. Mycol. Scot.* n. 495. *Ag. Bull.* t. 586. f. 2. *A.B.*, t. 578 partly.

\*\*\*\* *Stem floccoso-scaly, &c.*

**103. C. hemitrichus** Fr.—**Pileus** 2.5–7.5 cent. (1–3 in.) broad, dark fuscous, fuscous-tan when dry, umbo commonly persistently dark, somewhat fleshy, *convexo-expanded*, sometimes acutely, sometimes obtusely umbonate or wholly obtuse (in largest specimens even umbilicate on account of the hollow apex of the stem), sometimes wholly, sometimes *round the margin elegantly white-fibrillose-curved with erect flocci*, the fibrils however superficial, even, never torn. **Stem** 4–6 cent. (1½–2½ in.) long, at the very first hollow, equal, thick, firm, scarcely ever flexuous, pallid fuscous, white-flocculose below the medial ring which is woven, shining white, not rarely membranaceous and reflexed. **Gills** adnate, but ventricose at the base, hence they appear rounded, very crowded, 6 mm. (3 lin.) broad, quite entire, clay-colour, at length cinnamon.

Odour none, taste not unpleasant. Pileus stiff, scissile. It becomes adpressedly fibrillose when the flocci are flattened, and becomes smooth when these are wiped off by showers. The colour resembles that of *C. castaneus* when dry. In smaller forms the fibrils of the pileus are shorter, and the ring less conspicuous.

In mixed woods. Uncommon. Sept.–Oct.

Spores sphaeroid-ellipsoid,  $7 \times 5$  mk. *K.* Name—*οπίξ*, hair. Partially hairy. *Fr. Monogr.* ii. p. 89. *Hym. Eur.* p. 385. *Icon.* t. 160. f. 2. *B. & Br.* n. 930. *C. Hbk.* n. 534. *S. Mycol. Scot.* n. 496.

**104. C. stemmatus** Fr.—**Pileus** 4–5 cent. (1½–2 in.) broad, *date-brown* when moist, becoming pale when dry, slightly fleshy, convex then flattened, obtuse, fragile, *hoary-silky round the margin* when moist, *fibrillose* when dry. **Stem** 5–7.5 cent. (2–3 in.) long, stuffed then fistulose, equal or slightly attenuated at the base, often curved, soft, externally and internally ferruginous-date-brown, paler and becoming silky-even at the apex. **Gills** adnate, crowded, 4 mm. (2 lin.) broad, narrower behind, date-brown, opaque.

Distinguished by the *pileus being hoary-silky round the margin, and by the very crowded, narrow, date-brown gills*. It occurs under two forms: the *earlier* with the stem floccoso-squamulose and somewhat ringed; and the *later* with the stem naked, the veil being wiped off. Allied to *C. uraceus*, but in that species there is no trace of a woven veil even in its earliest stages.

In moist woods. Lyne, Sussex.

Name—*στέμμα*, a wreath. From the silky marginal band. *Fr. Monogr.* ii.

p. 89. *Hym. Eur.* p. 385. *Icon. t.* 160. f. 3. *Grevillea*, vol. xii. p. 68 *Telamonia*. (ringless form).

**105. C. paleaceus** Fr.—**Pileus** *fuscous* when damp, dingy when dry, very hygrophanous, somewhat membranaceous, conical then expanded, acutely or obtusely umbonate, at first *silky* with superficial squamulose white villous down (veil), becoming smooth, opaque; flesh of the disc of the same colour, almost none at the sides. **Stem** 7.5 cent. (3 in.) long, 2 mm. (1 lin.) thick, fistulose, slender, toughish, undulated on the surface, externally and internally fuscous, paler when young, white-villous at the base, here and there squamulose with white flocci, and white-ringed at the apex. **Gills** adnate, broad, *crowded*, at first *pallid-whitish*, at length cinnamon.

Odour weak. Commonly acutely umbonate. It differs from *C. iliopodius* in the stem not being tawny within, and in the veil being floccoso-scaly not becoming even.

In beech woods. Cabalva. Sept.—Oct.

Name—*palea*, chaff. From the scales. *Fr. Monogr.* ii. p. 92. *Hym. Eur.* p. 386. *Icon. t.* 160. f. 4. *Grevillea*, vol. viii. p. 77. *S. Mycol. Scot. Supp. Scot. Nat.* 1883, p. 31. *Quel. Grev. t.* 113. f. 5, t. 114. f. 4.

**Tribe VI. HYGROCYBE** (ὕγρός, moist; κῆβή, head). **Pileus** Hygrocybe. smooth or only covered with white superficial fibrils, not viscous, but moist when fresh, changing colour when dry, flesh very thin or scissile; rarely more compact at the disc. **Stem** slightly rigid, not sheathed; cortina thin, fibrillose, rarely collapsing and forming an irregular zone on the stem. *Quite distinct from Telamonia, but only by the veil.* *Fr. Hym. Eur.* p. 386.

**I. FIRMIORES.** Pileus somewhat fleshy, convex or campanulato-convex then expanded, obtuse or at length gibbous, *margin at first incurved.* Stem (for the most part) attenuated upwards.

\* *Stem white, cortina of the same colour.*

\*\* *Stem and gills commonly inclining to violet.* In various species of the former section there is a fugitive tinge of violet at the apex of the stem.

\*\*\* *Stem and somewhat obsolete veil yellow or rufous.*



XLV. *Cortinarius (Hygrocybe) castaneus.* One-third natural size.

Hygrocybe. \*\*\*\* *Stem inclining to fuscous*; cortina of a pallid dirty colour or white (not yellow); gills dark.

II. TENUIORES. Pileus somewhat membranaceous, conical then expanded, umbonate, umbo acute or more rarely obtuse and vanishing (the opposite is the way of the Firmiores), *margin at first straight*. Stem somewhat equal or attenuated at the base. (*The thinness of the pileus is only relative to its size.*)

\* *Stem white.*

\*\* *Stem inclining to violet or reddish.*

\*\*\* *Stem yellowish, commonly becoming pale.*

\*\*\*\* *Stem inclining to fuscous.*

### I.—FIRMIORES.

\* *Stem white, cortina of the same colour.*

106. **C. subferrugineus** Fr.—Pileus about 7.5 cent. (3 in.) broad, *ferruginous* or watery cinnamon, *more or less hygrophanous*, when dry sometimes shining and tawny, sometimes becoming pale, unequally fleshy, abruptly more compact at the disc, convex then expanded, obtuse, here and there flexuous, firm but not rigid, even, smooth; flesh scissile, dingy, isabelline-white. **Stem** variable in stature, commonly 7.5 cent. (3 in.) long, 12 mm. ( $\frac{1}{2}$  in.) thick, solid, more or less bulbous, attenuated upwards, adpressedly fibrillose, externally rigid and somewhat cartilaginous, pallid, internally soft, saffron-yellow at the base. Veil entirely fibrillose and marginal only, not sheathing, very fugacious. **Gills** remarkably emarginate, 6 mm. (3 lin.) *broad, more or less crowded, opaque*, at first pallid, soon watery then dark ferruginous.

Among heaped rotting pine-leaves the stem is stout, very inflated, spongy. Odour that of the *Inolomata*, taste unpleasant. There are many forms definite as to colour, but scarcely to be defined by description. It differs from *C. armeniacus* in the opaque colour, in the fleshy obtuse pileus, in the stem not being elastic, in the veil not collapsing, &c., and from *C. bivelus* in the simple veil, the smooth and hygrophanous pileus (watery ferruginous when moist, becoming tan-ferruginous when dry), &c.

In woods. Coed Coch. Sept.

Name—*ferrugo*, rust-colour. *Somewhat ferruginous*. Fr. Monogr. ii. p. 93. Hym. Eur. p. 387. B. & Br. n. 1275. Quel. t. 113. f. 6. Ag. Batsch f. 186 (not good).

107. **C. armeniacus** Fr.—Pileus 5–10 cent. (2–4 in.) broad, tawny-cinnamon when moist, ochraceous when dry, rigid, slightly fleshy, campanulate then convex and flattened, broadly and obtusely umbonate, even, smooth, not viscid, here and there slightly striate at the margin; flesh somewhat of the same colour, scissile, **Stem** commonly 5–7.5 cent. (2–3 in.) long, 12 mm. ( $\frac{1}{2}$  in.) thick, stuffed, *conico-attenuated*, fibrillose, white, *externally rigid and*



*somewhat cartilaginous*, internally spongy, hence elastic, *wholly* Hygrocybe. *white*. Cortina simple, white, *somewhat sheathing*, collapsing and commonly leaving an adpressed (not annular) zone on the stem. **Gills** adnate, at length slightly rounded, always crowded, rather broad, at first pallidly then tawny cinnamon, shining.

Handsome in pine woods. Firm, bright, changeable. Pileus soon destitute of white fibrils. There is a remarkable variety (*falsarius*) in which the pileus is paler, light yellowish, white when dry. On heaped rotting leaves in moist places the stem becomes very much swollen, among damp mosses it is longer and more slender.

In pine woods. Uncommon. Oct.

Spores ellipsoid, nucleate,  $4-5 \times 2-2\frac{1}{2}$  mk. *C.B.P.* Name—from likeness in colour to Apricot (*Prunus Armeniaca*). *Fr. Monogr.* ii. p. 94. *Hym. Eur.* p. 387. *Berk. Out.* p. 193. *C. Hbk.* n. 535. *S. Mycol. Scot.* n. 497. *Ag. Schæff.* t. 81.

**108. C. privignus** Fr.—**Pileus** about 5 cent. (2 in.) broad, the ground colour fuscous, but becoming *hoary-pale with a very thin white film*, pallid-tan when dry, slightly and unequally fleshy, convex then flattened (even reflexed and undulated), obtusely umbonate, without striæ, dry, but remarkably hygrophanous; flesh hygrophanous. **Stem** about 7.5 cent. (3 in.) long, 6 mm. (3 lin.) thick, stuffed then hollow, sometimes attenuated from the base, sometimes equal, most frequently twisted, silvery-pale, here and there white-silky with the veil. **Gills** adnate, broad, not crowded, distinct, at first watery, then opaque cinnamon, serrated and white-fimbriate at the edge when perfect.

*Very fragile*, strong smelling, but scarcely acrid. The most remarkable of the group.

In pine woods. Woolhope Foray, 1885. Oct.

Spores pruiniform, punctate, 8 mk. *Q.* Name—*privignus*, a step-son. Relation not apparent, but the plant being fragile occupies a singular position in the group. *Fr. Monogr.* ii. p. 95. *Hym. Eur.* p. 388. *Grevillea*, vol. xiv. p. 38.

**109. C. duracinus** Fr.—**Pileus** 5 cent. (2 in.) and more broad, when moist watery brick-colour, *tan* when dry, always *opaque*, fleshy, plane with a broad obtuse umbo, and with a singular elevated ridge round the margin caused by the margin being at first sharply and regularly bent inwards to the breadth of half a line, at which stage it is clothed with white silkiness, and afterwards becoming flattened, when it is naked and even; cuticle hard, rigid, fragile. **Stem** 7.5 cent. (3 in.) long, 12 mm. ( $\frac{1}{2}$  in.) thick, *solid*, hard, somewhat bulbous but *rooted at the attenuated stem-like base*, externally even, *smooth*, and *rigid*, with a *thick cartilaginous* rigid separable *cuticle* (the fragments of

Hygrocybe. which when it breaks up become revolute), pale-white. Cortina appearing only as a silky narrow white zone round the very margin of the pileus. **Gills** adnate, moderately crowded, 4-6 mm. (2-3 lin.) broad, *watery cinnamon*, even at the sides.

No trace of a fibrillose cortina has been observed in any stage of its growth. In pine woods the stem is not bulbous, but attenuated downwards.

In old pastures beside stumps. Dun, 1876. Scone, 1877. Oct.

Name—*durus*, hard; *actnus*, berry. Hard. From the coating of the pileus and stem. *Fr. Monogr.* ii. p. 96. *Hym. Eur.* p. 388. *B. & Br. n.* 1776. *S. Mycol. Scot. n.* 498. *Quel. Grev. t.* 115. f. 1.—*Schæff. t.* 221.

**110. C. dilutus** Fr.—**Pileus** about 5 cent. (2 in.) broad, at first bay-brown, even watery brick-colour, *tan when dry*, in either state *opaque*, somewhat fleshy, convex then expanded, umbonate (umbo thin, vanishing), smooth when the veil is rubbed off. **Stem** 5-7.5 cent. (2-3 in.) long, 4-8 mm. (2-4 lin.) thick, stuffed then hollow, slightly attenuated from the base, externally not cartilaginous - polished, white - silky, then becoming smooth, opaque, whitish. **Gills** deeply emarginate, *very ventricose*, 6-8 mm. (3-4 lin.) broad, *crowded*, *pale cinnamon*, not shining.

Inodorous. Veil white, often collapsing into spots on the stem or forming spurious zones, and on the pileus forming a white-silky zone round the margin. *Wholly soft and fragile*, and thus readily distinguished from the much more robust *C. armeniacus* with which it agrees in colours, so that it was formerly taken for a thinner form of that species.

In woods. Rare.

Not recorded since the time of Bolton. Name—*dilutus*, diluted, watery. Of the colour. *Fr. Monogr.* ii. p. 98. *Hym. Eur.* p. 389. *Berk. Out. p.* 194. *C. Hbk. n.* 536. *Grevillea, t.* 85. f. 2. var. *Ag. Bolt. t.* 10.

**\*\* Stem and gills commonly inclining to violet.**

**111. C. saturninus** Fr.—**Pileus** 5-7.5 cent. (2-3 in.) even 10-12.5 cent. (4-5 in.) broad, *dark bay-brown* somewhat umber when damp, soon becoming pale, brick-colour, changing colour very much, *unequally fleshy*, campanulate when young, then expanded, obtuse, even, smooth (or when young superficially *silky round the margin* with the veil); flesh violaceous then whitish, thin at the circumference. **Stem** 5-7.5 cent. (2-3 in.) long, 1-2.5 cent. ( $\frac{1}{2}$ -1 in.) thick, occasionally shorter and bulboso-ventricose, firm, stuffed, internally spongy, *thickened downwards*, fibrillose, *deep violet*, but becoming white. Cortina inferior, abundant, white. **Gills** rounded-adfixed, very broad (8 mm., 4 lin.), crowded, thin, fragile, *purplish* when young, at length watery ferruginous.

Somewhat caespitose. It varies in stature, sometimes much smaller. Cor- Hygrocybe.  
tina remarkably fibrillose, lax, not forming a zone. Manifestly allied to *C. armeniacus*; they might be easily confounded when this is old and changed in colour.

In grassy places. Woolhope Foray, 1885. Oct.

Name—*saturnine*, gloomy. Of the dark colour. *Fr. Monogr.* ii. p. 99. *Hym. Eur.* p. 390. *Grevillea*, vol. xiv. p. 39.

**112. C. imbutus** Fr.—**Pileus** 7.5 cent. (3 in.) broad, toast-brown then pale yellowish, *somewhat unequally fleshy, convex*, obtuse, even, smooth, obsoletely hoary-fibrillose towards the margin. **Stem** 7.5 cent. (3 in.) long, not 2.5 cent. (1 in.) thick, *solid, equal*, sometimes twisted, becoming even, scarcely fibrillose, whitish, pale violaceous at the apex, of the same colour internally. Cortina white, adhering to the margin of the pileus and to the apex of the stem, fugacious. **Gills** rounded, *somewhat distant*, 6 mm. (3 lin.) *broad*, with narrower and shorter ones intermixed, *at the first dark bluish-grey* or violaceous-cinereous, *never purplish*, then watery cinnamon.

Flesh dingy, manifestly violaceous only at the apex of the stem. It approaches *C. subferrugineus* in the same way as *C. saturninus* approaches *C. armeniacus*.

. In woods. Dinmore.

Name—*imbuo*, to soak, stain. Perhaps from the dyed flesh. *Fr. Monogr.* ii. p. 100. *Hym. Eur.* p. 390. *Grevillea*, vol. viii. p. 109.

**113. C. castaneus** Fr.—**Pileus** *fuscous-chestnut, scarcely becoming pale when dry*, but *shining*, umbo becoming black, paler at the circumference, slightly fleshy, but firm, almost *pliant*, campanulate then flattened, not or only obtusely umbonate, often irregular, even, smooth. **Stem** about 2.5 cent. (1 in.) long, 4–6 mm. (2–3 lin.) thick, *somewhat fistulose* (when young also stuffed), *tough, externally cartilaginous*, slightly fibrillose with the veil, otherwise smooth, even, somewhat equal, *pallid violaceous* or *pallid rufescent*, unchangeable when dry. Cortina entirely fibrillose, scanty, white. **Gills** adnate, but varying emarginate, *crowded*, thin, 4–6 mm. (2–3 lin.) broad, edge whitish, violaceous then ferruginous.

Gregarious, even caespitose, polymorphous, always very small. The pileus is often obsoletely white-silky round the margin with the cortina. Spores dark ferruginous.

In woods, &c. Common. July-Oct.

Odour none; taste like *Marasmius oreades*. *M. F. B.* Name—*castaneus*, chestnut. *Fr. Monogr.* ii. p. 101. *Hym. Eur.* p. 391. *Berk. Out.* p. 194. *C. Hbk. n.* 537. *S. Mycol. Scot. n.* 499. *Ag. Bull. t.* 268.

Hygrocybe.

\*\*\* *Stem and somewhat obsolete veil yellow or rufous.*

**114. C. colus** Fr.—**Pileus** 2.5–5 cent. (1–2 in.) broad, *brown-rufescent*, brick-colour paler and shining when dry, somewhat fleshy, small in comparison with the tall stem, campanulate when young, then convex, obtusely umbonate or wholly obtuse, well-formed; flesh of the same colour, dingy whitish when dry. **Stem** 10 cent. (4 in.) long, 6–8 mm. (3–4 lin.) thick below, only 4 mm. (2 lin.) at the apex, stuffed, almost solid, *equally attenuated upwards, encircled with the blood-red mycelium* at the base, which is sometimes rooted, stiff, naked, not cartilaginous but remarkably *fibrillose longitudinally* and throughout, scissile, tense and straight, *coloured with fibrils of the same colour as the pileus*, but paler than the pileus. Cortina entirely fibrillose, tawny-reddish. **Gills** adnate, scarcely sinuate, 6 mm. (3 lin.) broad, plane, firm, tough, rather thick, scarcely crowded, veined, dark cinnamon, paler when young.

Spores ochrey-cinnamon. *Mycelium fiery-saffron-yellow.* *Hym. Eur.*

In woods. Blaize Castle, Bristol. Oct.

Spores pruiniform, 8–9 mk. *Q.* Name—*colus*, a distaff. From the remarkably fibrillose stem. *Fr. Monogr. ii. p. 102.* *Hym. Eur. p. 391.* *Bucknall, Fung. Bristol, pt. vii. p. 3.*

**115. C. renidens** Fr.—**Pileus** 2.5–5 cent. (1–2 in.) broad, *feruginous-tawny* when damp, ochraceous when dry, here and there becoming pale only at the disc, slightly fleshy, firm, convexo-plane, *obtuse* or gibbous, very smooth, *shining*; flesh thin, scissile, paler. **Stem** 4–7.5 cent. (1½–3 in.) long, 6–8 mm. (3–4 lin.) thick, *firm*, stuffed, exactly *equal*, externally somewhat cartilaginous, wholly dissolvable however into fibrils, of the same colour as the pileus, yellowish-pale, then tawny. Cortina laxly *fibrillose*, fugacious, yellow. **Gills** adnate, but also separating-free, somewhat crowded, entire, about 6 mm. (3 lin.) broad, at first pallid cinnamon then tawny.

Spores dark ochraceous. Odour weak, by no means that of radish. Remarkably hygrophaneous. Its nature is that of *C. armeniacus*, but it is smaller, and is readily distinguished by its brighter colour, and especially by the *yellow cortina*. The pileus is sometimes umbilicate from the hollow apex of the stem.

In shady woods. Highbeach, Epping.

Name—*renideo*, to shine back. Glistening. *Fr. Monogr. ii. p. 103.* *Hym. Eur. p. 392.* *Icon. t. 162. f. 1.* *Grevillea, vol. xi. p. 70.*—*Paul. t. 54. f. 1, 2?*

\*\*\*\* *Stem inclining to fuscous, &c.*

Hygrocybe.

**116. C. uraceus** Fr.—**Pileus** 2.5–5 cent. (1–2 in.) broad, *when moist umber* or brown, sometimes inclining to olive, somewhat shining, *when dry* (when young brick-colour) commonly *tan* or isabelline, *slightly fleshy*, conical then campanulate, then expanded and umbonate or obtuse, when moist even, smooth, when dry at length somewhat fibrillose; flesh *fuscous*, darker in the stem. **Stem** 5–10 cent. (2–4 in.) long, 4–8 mm. (2–4 lin.) thick, slightly tubular, stuffed then hollow, cylindrical, exactly equal, firm, *fibrilloso-striate* (the striæ paler), but becoming even, *fuscous*, naked and becoming pale at the apex, sometimes inclining to olive, at length becoming wholly fuscous-black, internally of the same colour. *Cortina superior*, fibrillose, *fuscous*, but rarely conspicuous. **Gills** adnate, ventricose, firm, 6 mm. (3 lin.) and more broad, distant, *cinnamon-brown*, the edge sometimes quite entire and of the same colour, sometimes white and fimbriato-serrated.

The slight margin of the pileus is incurved. There are variations which cannot be noted.

In pine woods. Dinmore.

Spores pruiniform, granulated, 8–9 mk. *Q.* Name—*ῥαξ*, a mouse. Mouse-coloured. *Fr. Monogr.* ii. p. 112. *Hym. Eur.* p. 393. *Icon. t.* 162. f. 3. *Grevillea*, vol. viii. p. 77.

**117. C. jubarinus** Fr.—**Pileus** 4–7.5 cent. (1½–3 in.) broad, *splendidly tawny-cinnamon and shining*, somewhat fleshy, campanulato-flattened, obsoletely umbonate, when larger and old depressed in the centre, often repand and undulated, at length reflexed, fragile, even and smooth at the disc, when young silky towards the margin with the veil, when old innately fibrillose under a lens. **Stem** 5 cent. (2 in.) long, when larger 12 mm. (½ in.), when smaller 4–6 mm. (2–3 lin.) thick, at first stuffed, then hollow, *fibrilloso-striate*, *externally and internally pale tawny*, paler at the base and (naked) apex. **Gills** adnate, *somewhat distant*, 2–6 mm. (1–3 lin.) broad, distinct, *tawny-cinnamon*, the edge which is quite entire of the same colour.

At first sight very like *C. cinnamomeus*, but differing in the fugacious, fibrillose, *white veil*. The pileus is not velvety-villous as in *C. cinnamomeus*. It may also be confounded with *C. incisus* on account of its colours, but it differs from that species in the surface being scarcely hygrophanous, always polished and entire, although the pileus is rimoso-incised at the margin.

In woods. Coed Coch.

Name—*jubar*, radiance. Shining. *Fr. Monogr.* ii. p. 105. *Hym. Eur.* p. 393. *B. & Br. n.* 1883.—*Bull. t.* 431. f. 1.

Hygrocybe.

## II.—TENUIORES.

\* *Stem white.*

**118. C. rigens** Fr.—**Pileus** 2.5–5 cent. (1–2 in.) and more broad, *opaque and tan-clay-colour when damp, tan-whitish when dry, fleshy*, thin, when young campanulate and lax, then convex, obtuse or broadly gibbous, even, smooth; flesh slightly firm, white. **Stem** 5–10 cent. (2–4 in.) long, 4–10 mm. (2–5 lin.) thick, stuffed, furnished with a remarkably *cartilaginous bark*, elastic, *rigid and tough, rooted*, smooth, even, naked, *pale* when moist, white when dry. *Cortina scarcely manifest*. **Gills** *adnate, somewhat decurrent, distant*, very broad (6–10 mm., 3–5 lin.), distinct, watery clay-colour, then pallid cinnamon.

The stem varies much in stature, sometimes equal, thickened sometimes downwards sometimes towards the apex, which is then hollow, even fusiform.

In mixed woods. Uncommon. Oct.

Spores pruiniform, granulated, 8–9 mk. *Q.*; 7 × 5 mk. *W.P.* Name—*rigeo*, to be stiff. Stem rigid. *Fr. Monogr.* ii. p. 107. *Hym. Eur.* p. 395. *B. & Br. n.* 391. *C. Hbk. n.* 539. *S. Mycol. Scot. n.* 500.

**119. C. Reedii** Berk.—**Pileus** 2.5 cent. (1 in.) broad, *persistently brown*, conical then expanded and strongly umbonate, smooth, shining, *disc areolate*, margin splitting; flesh pallid. **Stem** 4–5 cent. (1½–2 in.) long, solid, fibrillose-striate, *slightly bulbous*, white. Veil fibrillose, evanescent. **Gills** ascending, attenuated behind, free, broad, ventricose, white or pallid, then cinnamon.

Tasteless and scentless. Allied to *C. leucopus*.

Among moss and beech mast. Hayes, Surrey. May.

Name—after Reed. *Berk. Out. p.* 194. *C. Hbk. n.* 538. *Fr. Hym. Eur. p.* 395. *Hussey ii. t.* 45.

**120. C. leucopus** Fr.—**Pileus** scarcely reaching 2.5 cent. (1 in.) broad, *very pale yellow*, tan when dry, *somewhat fleshy*, conical then expanded and umbonate, even, smooth, moist but not viscous. **Stem** 2.5–5 cent. (1–2 in.) long, 4–8 mm. (2–4 lin.) thick, stuffed then *hollow, rather thick*, equal or slightly attenuated upwards, soft, *shining white*, furnished with a cortina at the middle. **Gills** *adnexed*, separating, *crowded, ventricose*, thin, pallid then cinnamon, the edge which is quite entire of the same colour.

With the habit of *C. pluvius*, but not viscid.

In mixed woods. Frequent. Sept.–Oct.

Name—λευκός, white; πούς, a foot. White-stemmed. *Fr. Monogr.* ii. p. 108. *Hym. Eur.* p. 395. *Berk. Out.* p. 194. *C. Hbk. n.* 540. *S. Mycol. Scot. n.* 501. *Ag. Bull. t.* 533. f. 2. Hygrocybe.

**121. C. scandens** Fr.—*Pileus* when moist at first tawny-ferruginous, then honey-colour, tan when dry, umbo becoming tawny, somewhat membranaceous, slightly fleshy at the umbo, which is sometimes acute sometimes obtuse, conical when young, 12 mm. ( $\frac{1}{2}$  in.) broad, broader (2.5 cent., 1 in. and more) when flattened, smooth, slightly striate at the margin; flesh yellowish. *Stem* 7.5–10 cent. (3–4 in.) long, 4 mm. (2 lin.) thick, fistulose, thickened at the apex, always attenuated at the base, very thin, somewhat awl-shaped, flexuous, soft, even, yellowish, but at first sight, from the adpressed fibrils, and when dry shining whitish. Cortina thin, fibrillose, white. *Gills* adnate, thin, somewhat distant, narrow, 2–4 mm. (1–2 lin.) broad, attenuated in front, tawny-cinnamon, the edge which is quite entire of the same colour.

The colour varies in intensity with the age of the plant. The umbo is sometimes obsolete. The base of the stem is always white. Like *C. obtusus*.

In pine wood. Forres. Oct.

Name—scando, to climb. From the long-drawn wavy stem. *Fr. Monogr.* ii. p. 108. *Hym. Eur.* p. 396. *Icon. t.* 163. f. 1. *S. Mycol. Scot. Supp. Scot. Nat.* 1883, p. 31.

**\*\* Stem inclining to violet or reddish.**

**122. C. erythrinus** Fr.—*Pileus* 2.5–4 cent. (1–1½ in.) broad, bay-brown-rufous when moist, becoming tawny when dry, slightly fleshy, conical then convex, regular, umbonate, the obtuse or obsolete umbo darker, even, smooth; flesh when moist somewhat of the same colour. *Stem* 4–7.5 cent. (1½–3 in.) long, 4–6 mm. (2–3 lin.) thick, stuffed then hollow, equal, rarely thickened at the base, straight or ascending, smooth, but striate with the adpressed fibrils, shining, silvery-white, violaceous and often pruinose upwards. Cortina superior, fibrillose, white. *Gills* slightly adnexed, thin, somewhat distant, ventricose, at first pallid, then pale cinnamon.

In woods. Coed Coch. Oct.

Name—ερυθρός, red. *Fr. Monogr.* ii. p. 109. *Hym. Eur.* p. 396. *B. & Br. n.* 1665.

**123. C. decipiens** Fr.—*Pileus* 2.5 cent. (1 in.) or a little more broad, shining, not very hygrophanous, bay-brown, shining and brick-colour when dry, umbo always darker, fleshy-membranaceous, campanulato-expanded, acutely umbonate, at length depressed round the umbo, smooth, margin when full grown slightly

**Hygrocybe.** striate and at length innately torn. **Stem** 5-10 cent. (2-4 in.) long, 2-4 mm. (1-2 lin.) thick, stuffed, then fistulose, exactly equal, sometimes tense and straight, sometimes flexuous, but the surface not undulated, fibrillose, pallid, pale rufescent or with brick-coloured spots, somewhat brick-colour internally, externally *covered over with a pallid separable cuticle*. Cortina fibrillose, white, very fugacious. **Gills** adnate, more or less crowded, thin, 4-6 mm. (2-3 lin.) broad, brick-colour-ferruginous.

It approaches *C. castaneus*. It belongs to this series though the stem does not incline to violet, but is only reddish-white. There is a remarkable variety with paler pileus, flexuous and smooth stem, and less crowded gills.

In mixed woods. Frequent. Sept.

Name—*decipio*, to deceive. Beguiling, cheating. *Fr. Monogr.* ii. p. 109. *Hym. Eur.* p. 396. *B. & Br. n.* 1129. *C. Hbk. n.* 541. *S. Mycol. Scot. n.* 502. *Hoffm. Ic. t.* 9. f. 2. *Grevillea, t.* 114. f. 3.

**124. C. germanus** Fr.—**Pileus** almost 2.5 cent. (1 in.) broad, *remarkably hygrophanous, fuscous when moist*, clay-colour when dry, opaque, somewhat membranaceous, campanulate, obtusely umbonate when expanded, *fragile, somewhat silky*, but not torn into squamules and not striate. **Stem** 7.5 cent. (3 in.) long, 2 mm. (1 lin.) thick, *somewhat fistulose, equal*, but often twisted, smooth, silvery-pale, *somewhat lilac*. Cortina fibrillose, fugacious. **Gills** adnate, somewhat distant, broad, watery cinnamon.

Strong-smelling.

In pine woods. Logie, Forfarshire. Oct.

Name—*germanus*, full brother. Closely related to those next it. *Fr. Monogr.* ii. p. 109. *Hym. Eur.* p. 397. *S. Mycol. Scot. Supp. Scot. Nat.* 1882, p. 217. *Quel. Grev. t.* 114. f. 2.

\*\*\* *Stem yellowish, commonly becoming pale.*

**125. C. detonsus** Fr.—**Pileus** 2.5-5 cent. (1-2 in.) broad, *bright yellowish when moist*, tan when dry, *somewhat membranaceous*, conical then expanded, somewhat umbonate, slightly silky then rather smooth, striate to the middle when moist, even and *slightly silky when dry*; flesh thin, whitish. **Stem** 5-7.5 cent. (2-3 in.) long, about 4 mm. (2 lin.) thick, stuffed then fistulose, equal or attenuated upwards, *soft*, smooth, pale, *the yellowish colour deeply struck into the flesh*. **Gills** adnate, ventricose, distinct, somewhat distant, quite entire, unicolorous, *at first bright yellowish* then brick-cinnamon.

Fragile. The umbo is at first acute, gradually vanishing, sometimes darker. Stature almost that of *C. decipiens*. It occurs much larger with the stem yellowish-brick-colour.



Among moss in woods. Glamis, 1875. Sept.

Hygrocybe.

Name—*detondeo*, to shear. Shaven, smooth. *Fr. Monogr.* ii. p. 110. *Hym. Eur.* p. 397. *B. & Br. n.* 1552. *S. Mycol. Scot. n.* 503.

**126. C. obtusus** Fr.—**Pileus** 1-4 cent. ( $\frac{1}{2}$ -1 $\frac{1}{2}$  in.) broad, at the first bay-brown-ferruginous, soon cinnamon, when dry either (according to age) becoming pale-ochraceous or tan-whitish, somewhat membranaceous, conical then campanulate, at length expanded and obtusely umbonate, smooth, striate at the margin. **Stem** 5-10 cent. (2-4 in.) long, at first spongy-stuffed, soon hollow, comparatively thick, curved, flexuous (not undulated), attenuated at the base, fragile, sprinkled with adpressed white silky fibrils, otherwise rather smooth, tan-yellowish when moist, whitish when dry. **Gills** adnate, ventricose, somewhat distant, very broad, rather thick, the shorter ones narrower, connected by veins, tawny-cinnamon, white-fringed at the edge.

Spores somewhat ochraceous. Gregarious. Strong-smelling. *Cortina entirely fibrillose, whitish*; pileus when young here and there rendered hoary silky by the cortina, commonly smooth, but torn when dry. Umbo vanishing. It is very changeable in stature, but the colour is constant, though it changes so much with age and the state of the atmosphere that it requires long familiarity with the plant to enable one to distinguish individual specimens in different conditions.

In woods, chiefly pine. Frequent. Oct.

Name—*obtus*, obtuse. From the umbo. *Fr. Monogr.* ii. p. 111. *Hym. Eur.* p. 397. *Icon. t.* 163. f. 3. *B. & Br. n.* 1274. *S. Mycol. Scot. n.* 504.

**127. C. acutus** Fr.—**Pileus** honey-colour to very pale yellow and somewhat shining, tan or white when dry, with a silky appearance, somewhat membranaceous, conical then campanulate, then expanded, umbo acute and of the same colour, at length depressed round the umbo, striate when moist, obsoletely fibrillose round the margin when young, smooth when full grown; flesh very thin, of the same colour. **Stem** 7.5 cent. (3 in.) and more long, 2-4 mm. (1-2 lin.) thick, fistulose, slender, equal, flexuous, white-fibrillose, at length smooth, of the same colours as the pileus when moist and dry. *Cortina* adhering to the margin, fibrillose, white. **Gills** adnate, thin, crowded, more distant as the pileus becomes more expanded and sometimes free, lanceolate, ochraceous-cinnamon.

Sometimes scattered, sometimes growing in troops. The habit is that of *Galera*, but remarkable for its acute umbo.

In fir and mixed woods. Frequent. Sept.-Nov.

Spores ellipsoid-oblong, nucleate, 8-9  $\times$  3-4 mk. *C.B.P.* Name—*acutus*, acute. From the umbo. *Fr. Monogr.* ii. p. 112. *Hym. Eur.* p. 398. *Berk.*

Hygrocybe. *Out. p.* 195. *C. Hbk. n.* 542. *S. Mycol. Scot. n.* 505. *Quel. Grev. t.* 112. *f.* 5.

\*\*\*\* *Stem inclining to fuscous.*

**128. C. Junghuhnii** Fr.—**Pileus** about 2.5 cent. (1 in.) broad, *shining cinnamon* when damp, tawny when dry, slightly fleshy at the disc, otherwise thin, *convexo-plane, umbonate with a papilla*, when damp striate to the middle, *under a lens somewhat velvety with thin, sparse, white, persistent fibrils*. **Stem** 5–7.5 cent. (2–3 in.) long, 4 mm. (2 lin.) thick, stuffed then hollow, externally pale brick-colour, internally darker, externally *shining* and *adpressedly fuscous-fibrillose*, quite equal or attenuated and whitish at the base. Veil obsolete, inclining to fuscous. **Gills** adnate, ventricose, 4 mm. (2 lin.) broad, thin, veined at the base, saffron-brick-colour.

Odour none.

In woods. King's Cliffe. Aug.

Spores 8 mk. *B. & Br.* Name—after Junghuhn. *Fr. Monogr. ii. p.* 113. *Hym. Eur. p.* 398. *B. & Br. n.* 1130. *C. Hbk. n.* 543.

**129. C. milvinus** Fr.—**Pileus** 1–2.5 cent. ( $\frac{1}{2}$ –1 in.) broad, *olivaceous-fawn-colour*, becoming hoary-tan and opaque when dry, membranaceous with exception of the disc, conical then convex, obtusely or obsoletely umbonate, when full grown striate to the middle and *beautifully wreathed at the margin with white squamules*, somewhat silky when dry; flesh very thin, of the same colour, not scissile. **Stem** 5–7.5 cent. (2–3 in.) long, equal, 4 mm. (2 lin.) thick, *somewhat fistulose*, curved, but never flexuous, even, somewhat naked (not scaly), *pallid fuscous*, here and there with white silky spots. Veil obsolete, except in the form of the wreath encircling the margin of the pileus. **Gills** adnate, connected by veins, somewhat distant, very thin, scarcely ventricose, watery ferruginous, becoming olive.

Scented. A marked species, growing in troops after heavy rain.

In woods. Wrotham, Kent. Oct.

Name—*milvus*, a kite. Of the colour of a kite's back. *Fr. Monogr. ii. p.* 114. *Hym. Eur. p.* 399. *B. & Br. n.* 1553. *Quel. Grev. t.* 114. *f.* 6.

**130. C. fasciatus** Fr.—**Pileus** 1 cent. ( $\frac{1}{2}$  in.) broad, *somewhat membranaceous, conical* then expanded, *the acute umbo blackish*, brick-colour at the circumference, smooth, slightly silky and becoming pale when dry. **Stem** sometimes 5 cent. (2 in.) long, 2 mm. (1 lin.) thick, straight, sometimes elongated 7.5–10 cent.

(3-4 in.) and flexuous, fistulose, undulated on the surface, smooth, *Hygrocybe* not furnished with a cuticle but wholly fissile into fibrils, *pallid fuscous*, at length cinnamon-fuscous. **Gills** *adnate*, slightly ventricose, *thin*, but truly *distant*, distinct, 3 mm. (1½ lin.) broad, cinnamon.

Spores cinnamon. On account of its colours it is like *C. flexipes*, but in its whole nature it is nearest to *C. acutus*.

In pine woods. Dinmore, &c. Sept.-Oct.

Umbo very acute. *M. F. B.* Name—*fascia*, band; fillet. From the stem splitting into bundles of fibres. *Fr. Monogr.* ii. p. 114. *Hym. Eur.* p. 399. *Grevillea*, vol. viii. p. 78. *B. & Br. n.* 1884. *Quel. Grev. t.* 114. f. 5.

# GENUS VI.—*Gomphidius* (γόμφος, a wooden bolt or nail).

*Gomphidius*.

*Fr. Epicr.* p. 319.

Hymenophore decurrent on the stem. Gills *composed of a mucilaginous membrane*, scissile, continuous at the acute edge, pruinose with the blackish *fusiform spores*. Veil viscoso-floccose. *Growing on the ground, fleshy, putrescent, pileus at length turbinate; gills decurrent, distant, soft.*

A small genus, with great differences among the species, intermediate by its habit between *Cortinarius* and *Hygrophorus*. *Fr. Hym. Eur.* p. 399.

Universal veil glutinous, at first terminating on the stem in a floccose ring or fugacious cortina. The gills often admit of being detached and stretched out into a continuous membrane. *Fr. Monogr.* ii. p. 149.



XLVI. *Gomphidius viscidus*.  
One-fifth natural size.

**1. *G. glutinosus* Fr.**—**Pileus** 5-12.5 cent. (2-5 in.) broad, purple-fuscous, often mottled with black spots, fleshy, convex, obtuse, at length plane, even depressed, even, smooth, very glutinous; flesh thick, about 12 mm. (½ in.), soft, white. **Stem** 5-7.5 cent. (2-3 in.) and more long, about 12 mm. (½ in.) thick, solid, whitish, thickened and externally and internally yellow at the base, viscid with the veil, fibrillose or varying with black

Gomphidius. scales. Cortina often woven in the form of a ring, but soon fugacious. **Gills** deeply decurrent, distant, distinct, branched, quite entire, mucilaginous, 6–8 mm. (3–4 lin.) broad, at first whitish, then cinereous, clouded with the spores.

Trama none, wherefore the gills easily separate from the pileus. Taste watery, mouldy. Odour not marked.

In woods, chiefly pine. Common. July–Nov.

Spores oblong, uniguttate, 18–23 × 6–8 mk. *K.*; 16–17 × 6 mk. *W.G.S.* Name—*gluten*. Glutinous. *Fr. Monogr.* ii. p. 149. *Hym. Eur.* p. 399. *Berk. Out.* p. 196. *C. Hbk.* n. 579. *S. Mycol. Scot.* n. 506. *Ag. Schæff.* t. 36. *Sow. t. 7. Fl. Dan.* t. 1247. *Letell.* t. 547. *Krombh.* t. 62. f. 18–20. *Corda ap. Sturm.* xi. t. 51.

**Var. roseus.**—**Pileus** rose-colour. **Stem** white, attenuated and rosy flesh-colour internally at the base.

Very distinguished, constant, always smaller. Spores oblong, 20–22 × 6 mk. *K. Fr. Monogr.* ii. p. 150. *Hym. Eur.* p. 400. *C. Hbk.* n. 579. *S. Mycol. Scot.* n. 506. *Krombh.* t. 63. f. 13–17. *Saund. & Sm.* t. 8.

**2. G. viscidus** Fr.—**Pileus** 5–7.5 cent. (2–3 in.) and more broad, fuscous-rufous, compact, at first campanulate, then expanded, umbonate, slightly viscous, shining when dry; flesh yellowish. **Stem** 7.5–10 cent. (3–4 in.) and more long, 12 mm. (½ in.) thick, solid, equal or attenuated at the base which is rhubarb-coloured internally, scaly-fibrillose, not very viscous, yellowish. Cortina very evidently floccose, not glutinous, woven in the form of a ring, but readily falling off. **Gills** deeply decurrent, distant, the shorter ones adnexed to the longer, not truly branched, at first paler, somewhat olive, at length fuscous-purple, clouded with the spores.

Hymenophore descending between the gill plates. Odour not unpleasant.

In woods, chiefly pine. Frequent. July–Oct.

In *Hym. Eur.* Fries describes the gills as truly branched, while in *Monogr.* they are described as above. Name—*viscidus*, viscid. Spores 16–17 × 6 mk. *W.G.S. Fr. Monogr.* ii. p. 150. *Hym. Eur.* p. 400. *Berk. Out.* p. 196. *C. Hbk.* n. 580. *S. Mycol. Scot.* n. 507. *Ag. Linn.*—*Schæff.* t. 55. *Krombh.* t. 4. f. 5–7. *Pers. Ic. & descr.* t. 13. f. 1–3.

**3. G. gracilis** B. & Br.—**Pileus** 2.5 cent. (1 in.) broad, pale vinous brown, clothed with dingy fuliginous gluten, at length spotted with black. **Stem** 5 cent. (2 in.) long, 3 mm. (1½ lin.) thick, slender, flexuous, pale, sprinkled with minute white scales above, virgate below (with the remains of the slime), yellow at the base. **Gills** decurrent, arched, forked, thick, obtuse, of a watery dingy white.

The gluten on the pileus dries especially round the margin into black spots, Gomphidius, or forms a narrow irregular black border. The gills are clothed under a lens with short tomentose hairs. Distinguished from other species by its slender, thin, and delicate habit.

In fir woods. Frequent. July–Oct.

Spores oblong, elliptic,  $24 \times 9$  mk. with a nucleus at either end. *B. & Br.*; fusiform ellipsoid, 15 mk. *Q.* Name—*gracilis*, slender. *B. & Br.* n. 698. *Berk. Out.* p. 196. t. 12. f. 7. *S. Mycol. Scot.* n. 508. *Fr. Hym. Eur.* p. 400. *G. stillatus* *C. Hbk.* n. 581.

GENUS VII.—**Paxillus** (*paxillus*, a small stake). *Fr. Gen.* Paxillus.  
Hymenom. p. 8.

Hymenophore continuous with the stem, decurrent. *Gills* membranaceous, scissile, somewhat branched and here and there anastomosing behind, *distinct from the hymenophore and readily separating from it.* Spores dingy whitish or ferruginous. *Fleshy, putrescent, pileus at first with the margin involute then continuously and gradually unfolded and dilated, indeterminate.*

A natural genus, but its extent is not yet properly determined. There are several species of *Tricholoma* and *Clitocybe*, which apparently ought to be transferred to it. It is analogous to *Boletus* in respect of its gills easily separating from the pileus, but to *Polyporus* in respect of its indefinite growth. Most of the species seem to be edible. *Fr. Hym. Eur.* p. 400.

It comprehends two typical sections which are manifestly distinct, the one that which has *P. lepista* for its type, the other (Tapinia) that which has *P. involutus* for its type. The former always grow on the ground, but the Tapiniæ occur both on the ground and on trunks; the former have the pileus entire, the latter more or less dimidiate, or altogether lateral and sessile. *Fr. Monogr.* ii. p. 115.

*Trib. I. LEPISTA (P. lepista).* Pileus entire, central. Spores dingy, only in *P. panæolus* with a tendency to ferruginous.

*Trib. II. TAPINIA (ταπεινός, to depress).* Pileus commonly excentric or resupinate. Spores ferruginous.



XLVII. *Paxillus involutus.*  
One-fourth natural size.

Paxillus.

## I.—LEPISTA.

1. **P. giganteus** Fr.—Tan-white. **Pileus** fleshy, comparatively *thin*, depressed then plano-infundibuliform, soft, margin involute, then spreading, smooth, *sulcate in the form of small channels*. **Stem** solid, obese, smooth. **Gills** somewhat decurrent, very crowded, here and there branched and anastomosing, whitish then tan-colour.

Large. The pileus is smooth when moist, flocculose or rivulose when dry, easily splitting from the margin to the stem. *Hymenophore not horny*. Very remarkable, very distinct from *A. maximus*.

In meadows and woods. Uncommon. Sept.—Oct.

Pileus 4–14 inches broad, fleshy, often splitting at the margin, broadly infundibuliform, the base of the funnel sunk into the stem with no trace of an umbo, dirty white with an ochraceous tinge, minutely adpresso-squamulose to the naked eye, sometimes guttate; the whole surface under a lens clothed with a fine matted silkiness; margin grooved, the grooves shallow. Gills close, forked, yellow-white, as broad as the flesh of the pileus. Stem 2½–3 inches high, nearly 2 thick at the base, firm, fleshy, elastic, quite solid, sub-bulbous, sometimes attenuated upwards, minutely but conspicuously pubescent, when bruised dirty rufescent. Odour strong like that of *M. oreades*. *M. J. B.* Name—*giganteus*, of gigantic size. *Fr. Hym. Eur. p. 401. S. Mycol. Scot. n. 509. Ag. Sow. t. 244. Fr. Monogr. i. p. 118. Sv. atl. Sv. t. 86. Berk. Eng. Fl. v. p. 33. Out. p. 110. Letell. t. 682. Quel. t. 3. f. 3. A. maximus C. Hbk. n. 95.*

2. **P. lepista** Fr.—Pileus 7.5–10 cent. (3–4 in.) and more broad, dingy whitish, sometimes (young) compactly fleshy, sometimes (full grown) thin, plano-depressed, obtuse, *without striæ*, but *rimuloso-squamulose towards the circumference*, dry, the involute margin commonly undulato-flexuous; flesh white. **Stem** sometimes curt, 2.5 cent. (1 in.) or little more, attenuated downwards, sometimes elongated to 7.5–10 cent. (3–4 in.), equal, always blunt at the base, about 12 mm. (½ in.) thick and more, solid, compact, spongy-elastic, and at length pierced by larvæ and hollow, whitish, sometimes rufescent or inclining to fuscous, white-villous at the base. **Gills** *deeply decurrent, somewhat branched* but simple at the base, 4–6 mm. (2–3 lin.) broad, very crowded, quite entire, dingy white, *at length darker*.

*The cuticle of the stem is somewhat horny, continuous with the hymenophore and similar to it. Solitary and sporadic, very changeable. Odour meaty, somewhat rancid, approaching that of A. prunulus, to which it is somewhat allied.*

In woods. Slough.

Spores reddish, then becoming fuscous-pallid. *Fr.* Name—*lepista*, a pan. From its shape. *Fr. Monogr. ii. p. 116. Hym. Eur. p. 402. Icon. t. 164. f. 1. B. & Br. n. 1554. Ag. Sterb. t. 19. C.*

**3. P. paradoxus** Berk.—**Pileus** 5–7.5 cent. (2–3 in.) broad, deep *Paxillus*. rufous-umber or yellowish-brown, fleshy, compact, in some cases subexcentric, convex or pulvinate then plane, scarcely depressed, often lobed and sinuate at the margin, dry, adpressedly tomentose, not shining. **Stem** 4 cent. (1½ in.) and more long,  $\frac{3}{10}$ – $\frac{4}{10}$  in. thick and when bulbous thicker, solid, bulbous or attenuated at the base, fusiform, somewhat rooted, adpressedly fibrillose, otherwise even and smooth, *the cuticle most frequently breaking up into squarrosely revolute flaps*, yellow, sometimes with a dingy purplish tinge. **Gills** truly decurrent, distant, the alternate ones broader,  $\frac{1}{10}$ – $\frac{3}{10}$  in., entire at the edge, connected by veins at the base and *in some cases anastomosing to form pores towards the margin*, yellow then golden, reddish when bruised.

Stem variable in stature. *No trace of a veil.* Flesh soft, juicy, dingy white, slightly rufous under the cuticle of the pileus, becoming yellow at the base of the stem. Taste and odour scarcely any.

In woods. Wrekin, Salop, 1875. Sept.

Spores ovate, oblong, of medium size, lurid ochraceous. *Kalchb.* 16–17 × 6 mk., with a nucleus at each end. *W.P.* Name—*παράδοξον*, contrary to expectation or rule. From its having features contrary to normal *Agaricus*, under which Kalchbrenner placed it. *Berk. Ag. Kalchb. Fung. Hung. t. 16. f. 1. Fr. Hym. Eur. p. 244. Grevillea, vol. iv. p. 118.*

**4. P. panæolus** Fr.—**Pileus** 2.5–5 cent. (1–2 in.) broad, whitish, fleshy, thin, convexo-plane then somewhat depressed, even, smooth, moist, the thin margin involute; flesh becoming black. **Stem** 2.5 cent. (1 in.) or little more long, 6 mm. (3 lin.) thick, stuffed, striato-fibrillose, rufescent, thickened downwards. **Gills** slightly decurrent, crowded, narrow, at length *watery-ferruginous*.

The whole plant is at first whitish, then the stem is rufescent, and the gills in the end become cinnamon. The gills are separated from the hymenophore by a horny line, and readily separate from the pileus. Spores watery ferruginous, in which it approaches the *Tapiniae*.

\* **spilomæolus**.—**Pileus** *spotted as with drops*, and, as well as the slender stem, yellowish-white. Gills at length watery ferruginous, horny-grey at the base.

Among fir-leaves. Stoke Poges.

The spotted pileus and dingy spores at once distinguish it from any *Tricholoma* with which it might be confounded. The stem is sometimes incrassated at the base, sometimes quite equal. *B. & Br.* The var. only is British. Name—*παναίολος*, all-variegated. *Fr. Monogr. ii. p. 117, 310. Hym. Eur. p. 402. B. & Br. n. 1666. Hoffm. Ic. t. 10. f. 1.*

Paxillus.

## II.—TAPINIA.

**5. P. involutus** Fr.—**Pileus** 7.5–12.5 cent. (3–5 in.) broad, ochrey-ferruginous, fleshy, firm, convexo-plane then depressed, for the most part central, at the first pubescent, soon becoming smooth, somewhat viscid when moist, shining when dry, obtuse and villous round the margin which is at first closely involute, when more fully grown extenuated and acute; *flesh* compact, *pallid*. **Stem** 5–7.5 cent. (2–3 in.) long, 1–2.5 cent. ( $\frac{1}{2}$ –1 in.) thick, fleshy, solid, firm, thickened upwards, more rarely at the same time bulbous at the base, naked, dingy yellowish, commonly spotted. **Gills** decurrent, rather broad, branched and forming pores behind, when young arcuate, crowded, pallid, at once spotted when touched, at length extended, ferruginous.

Commonly large, obese, readily distinguished by the *pileus being villous round the very involute margin, by the gills anastomosing and forming pores at the base and being dingy-spotted when touched*. Pileus more rarely excentric, darker in colour in wet, and paler in dry weather. It varies lobed and scaly.

In woods. Very common. June–Nov.

Edible; highly esteemed in Russia. Spores ferruginous, *Fr.*; ellipsoid or sphaeroid-ellipsoid or oblong-ellipsoid, 8–16 × 6 mk. *K.*; 5 × 6 mk. *W.G.S.* Name—*involutus*, rolled inwards. From the involute pileus. *Fr. Monogr.* ii. p. 118. *Hym. Eur.* p. 403. *Sv. ätl. Sv. t.* 75. *Berk. Out. p.* 195. *t.* 12. *f.* 5. *C. Hbk. n.* 547. *S. Mycol. Scot. n.* 510. *Hoffm. Ic. t.* 10. *f.* 2. *Ag. Batsch f.* 61. *Ventur. t.* 42. *f.* 6, 7. *Schæff. t.* 72. *Bull. t.* 240, 576. *f.* 2. —*Klotzsch. Bor. t.* 391.

**6. P. leptopus** Fr.—**Pileus** 4–7.5 cent. ( $1\frac{1}{2}$ –3 in.) broad, fuscous-yellowish, always excentric or lateral, at length depressed but gibbous in the middle, fleshy, thin, dry, covered with dense down, soon *torn* up into dense villous scales, which are fuscous or yellowish; *flesh yellow*. **Stem** very short, scarcely reaching 2.5 cent. (1 in.), somewhat incurved, attenuated downwards, *yellow within*. **Gills** *simple* behind, decurrent but not anastomosing, *tense and straight*, crowded, very narrow, yellowish, then darker, but not spotted when touched.

Changeable in stature and form. A remarkable species, with the habit of *P. involutus*, from which it is readily distinguished by having the gills simple (not anastomosing) at the base.

On the ground about stumps, &c. Forres. Glen Tilt. Kenmore. Aug.

Name—λεπτός, thin; πούς, a foot. Thin-stemmed. *Fr. Hym. Eur. p.* 403. *Icon. t.* 164. *f.* 3. *B. & Br. n.* 1666\*. *S. Mycol. Scot. n.* 511. *A. filamentosus Fr. Monogr.* ii. 118, 311. *B. & Br. n.* 1276.

**7. P. atrotomentosus** Fr.—**Pileus** 5–10 cent. (2–4 in.) and more



broad, ferruginous, compactly fleshy, excentric, plano-infundibuliform and sometimes wholly lateral and ascending, dry, *rivuloso-granular* on the surface, sometimes also slightly tomentose, the thin margin involute; *flesh white*. **Stem** 5-7.5 cent. (2-3 in.) long, 1-2.5 cent. ( $\frac{1}{2}$ -1 in.) thick, solid, elastic, somewhat equal, not tuberous, curved-ascending, rooting, covered over with *dense*, soft *velvety down*, which is *umber-blackish* or inclining to violaceous. **Gills** adnate, scarcely decurrent, branched at the base, somewhat anastomosing, crowded, 6 mm. (3 lin.) broad, yellowish, easily separating from the sulcate hymenophore. Paxillus.

Commonly solitary. Robust, firm, often very large. The pileus is variable in form. The gills do not form pores like those of *P. involutus*. Sometimes lateral, but marginate behind.

On pine-stumps. Rare. Aug.-Nov.

Spores somewhat clay-colour, paler than those of *P. involutus*. *Fr.*; ellipsoid-sphæroid, subhyaline,  $4-6 \times 3-4$  mk. *K.* Name—*ater*, black; *tomentum*, down. *Fr. Monogr.* ii. p. 119. *Hym. Eur.* p. 403. *Berk. Out.* p. 195. *C. Hbk. n.* 548. *S. Mycol. Scot. n.* 512. *Ag. Batsch f.* 32. *Paul. t.* 33. *f.* 2, 3. *Nees. f.* 175.

**8. P. panuoides** Fr. Wholly dingy yellow.—**Pileus** about 4 cent. ( $1\frac{1}{2}$  in.) long, fleshy, *sessile or extended behind and at the first resupinate*, soon conchate dimidiate and obovate, at length broadly expanded undulato-lobed and often imbricated, surface pubescent then becoming smooth, somewhat rivulose; flesh equal but thin. **Gills** decurrent to the base, anastomosing behind, *branchæ*, crowded, *crisped, yellow*.

Very variable in size and changeable in form.

On sawdust, &c. Uncommon. Aug.-Oct.

Spores ellipsoid-sphæroid,  $4-6 \times 3-4$  mk. *K.*;  $3 \times 4$  mk. *W.G.S.* Name—*Panus*; εἶδος, appearance. *Panus*-like. *Fr. Monogr.* ii. p. 120. *Hym. Eur.* p. 404. *Berk. Out.* p. 196. *t.* 12. *f.* 6. *C. Hbk. n.* 549. *S. Mycol. Scot. n.* 513. *Merulius Sow. t.* 403.

**9. P. fagi** B. & Br.—Crisped, pallid upwards, orange beneath. **Gills** crisped, orange.

Remarkably gregarious. Forming a wide crisped mass of great beauty, very different in appearance from *P. panuoides*, which is confined to fir or sawdust.

On a beech-stump. Coed Coch.

Name—*fagus*, beech. *B. & Br. n.* 1961.

Hygro-  
phorus.

GENUS VIII.—**Hygrophorus** (ὕγρός, moist; φέρω, to bear).  
Fr. Gen. Hymen.

Hymenophore continuous with the stem and descending unchanged as a trama into the gills. Gills acute at the edge, clothed with the hymenium which turns into a waxy mass, and not membranaceous. Spores globose, white. *Growing on the ground, putrescent; pileus viscid or watery, gills often branched.* Fr. Hym. Eur. p. 405.



XLVIII. *Hygrophorus pratensis*.  
One-third natural size.

This genus differs from the central one of *Agaricus* in the manifest trama, the substance of which is similar to that of the pileus; from *Lactarius* and *Russula* by the trama not being vesicular, but somewhat floccose with granules intermixed; from *Cantharellus*, its nearest ally, by the sharp edge of the gills. The *Cortinarii*, *Paxilli*, and *Gomphidii*

are at once distinguished from it by their coloured spores and the changing colour of their gills, as well as by other marks. From all the other genera of Agaricini it is distinguished by a mark peculiar to itself—viz., by the hymeneal stratum of the gills changing into a waxy mass, which is at length removable from the trama. This altogether singular character is specially remarkable in *H. caprinus*, *coccineus*, *murinaceus*, &c. Hence the gills seem full of watery juice, but they do not become milky like those of the *Lactarii*. *Monogr.* ii. p. 121.

I. LIMACIUM (*limax*, a slug; slimy). Fr. Syst. Myc. p. 31. Universal veil viscid, with occasionally a floccose partial one, which is annular or marginal. Stem clothed with scales, or more frequently rough with dots above. Gills adnato-decurrent.

\* *White or yellowish white.*

\*\* *Reddish.*

\*\*\* *Tawny or light yellow.*

\*\*\*\* *Olivaceous-umber.*

\*\*\*\*\* *Fuscos-cinereous or livid.*

II. CAMAROPHYLLUS (καμάρα, a vault; φύλλον, a leaf; from the shape of the gills). Fr. Syst. Myc. i. p. 98. Veil none. Stem even, smooth or fibrillose, and not rough with dots. Pileus *firm*, opaque, *moist* in rainy weather, *not viscid*. Gills distant, arcuate.

\* *Gills deeply and at length obconically decurrent.*

\*\* *Gills ventricose, sinuato-arcuate, or plano-adnate.*

III. **HYGROCYPE** (ὕγρος, moist; κῆβή, a head). Fr. Syst. Myc. i. p. 101. Hygro-  
Veil none. Whole fungus thin, watery succulent, *fragile*. Pileus *viscid* phorus.  
when moist, *when dry shining*, rarely floccoso-scaly. Stem hollow, soft, not  
punctate. Gills soft. *Most of the species brightly coloured and shining.*

This group presents the type of the genus, and clearly differs from the  
*Agarici*; from the rest of the Hygrophori it might easily be separated as a  
genus by itself.

\* *Gills decurrent.*

\*\* *Gills adnexed, somewhat separating.*

#### I.—LIMACIUM.

\* *White or yellowish white.*

1. **H. chrysodon** Fr.—Pileus 5-7.5 cent. (2-3 in.) broad, *white*,  
shining when dry, but commonly yellowish with minute adpressed  
squamules at the disc, light yellow-flocculose at the involute *mar-*  
*gin*, fleshy, convex then plane, obtuse, viscid; flesh white, some-  
times reddish. **Stem** 5-7.5 cent. (2-3 in.) long, about 12 mm.  
(½ in.) thick, stuffed, soft, somewhat equal (sometimes, however,  
irregularly shaped or thickened at the base), white, with minute  
*light yellow squamules*, which are more crowded and arranged in  
the form of a ring *towards the apex*. **Gills** decurrent, distant,  
6 mm. (3 lin.) broad, thin, white, somewhat yellowish at the edge,  
sometimes crisped.

Odour not unpleasant. There is a manifest *veil*, not woven into a continu-  
ous ring, but *collected in the form of floccose squamules at the apex of the stem*  
*and the margin of the pileus*. Var. *leucodon* with white squamules.

In woods. Rare.

Name—*χρυσός*, gold; *ὀδόν*, a tooth. From the yellow tooth-like squamules.  
Fr. Monogr. ii. p. 123. Hym. Eur. p. 405. Berk. Out. p. 197. C. Hbk. n.  
550. Ag. Batsch f. 212. Gonn. & Rab. viii.-ix. t. 10. f. 1. Var. with shin-  
ing white pubescence B. & Br. n. 1356\*.

2. **H. eburneus** Fr. Wholly *shining white*.—Pileus fleshy,  
sometimes thin, sometimes somewhat compact, convexo-plane,  
somewhat repand, even, *very glutinous* in rainy weather, *margin*  
*soon naked*. **Stem** sometimes short, sometimes elongated, stuffed  
then hollow, unequal, *glutinous* like the pileus, *rough at the apex*  
*with dots in the form of squamules*. **Gills** decurrent, distant,  
veined at the base, 6-8 mm. (3-4 lin.) broad, tense and straight,  
quite entire.

Odour mild, not unpleasant. Very changeable. The veil is absent, unless  
the *very plentiful gluten* which envelops the stem be regarded as a universal  
veil; *margin of the young pileus* involute, only at the first *pubescent, soon*  
*naked*. The stem is soft internally, at length hollow, attenuated towards the  
base.

In woods and pastures. Frequent. Sept.-Oct.

Hygro-  
phorus.

Edible. Spores subsphaeroid, irregular, 5-6 mk. *K.*; 4×5 mk. *W.G.S.*; subellipsoid, 6×4 mk. *C.B.P.* Name—*ebur*, ivory. Ivory-white. *Fr. Monogr.* ii. p. 123. *Hym. Eur.* p. 406. *Berk. Out.* p. 197. *t.* 15. *f.* 4. *C. Hbk.* n. 551. *S. Mycol. Scot.* n. 514. *Gonn. & Rab. t.* 11. *f.* 5. *Ag. Bull. t.* 551. *f.* 2. *Viv. t.* 17. *Schæff. t.* 39. *Krombh. t.* 61. *f.* 11-14. *Buxb. iv. t.* 30. *f.* 2.

**3. H. cossus** Fr.—**Pileus** 4 cent. (1½ in.) broad, yellowish-white, disc somewhat ochraceous, fleshy, convexo-plane, obtuse, smooth, glutinous, shining when dry, *margin naked*; flesh white. **Stem** 5-7.5 cent. (2-3 in.) long, varying elongated, 4-10 mm. (2-5 lin.) thick, varying thin, stuffed, soft, somewhat equal, furfuraceous and rough with dotted points upwards, white or becoming tinged with yellow. **Gills** adnate, decurrent, distant, connected by veins, firm, white.

Veil none. It differs from *H. eburneus* in the *pileus being yellowish, in the margin being at the first naked, and especially in its unpleasant odour of goat-moth.*

In woods. Frequent. Sept.-Oct.

Name—from its odour of goat-moth (*Phalæna cossi*). *Fr. Monogr.* ii. p. 124. *Hym. Eur.* p. 406. *Berk. Out.* p. 197. *C. Hbk.* n. 552. *S. Mycol. Scot.* n. 516. *Ag. Sow. t.* 121.

**4. H. penarius** Fr.—**Pileus** tan-colour, opaque, fleshy especially when young, at first umbonate, then very obtuse, hemispherical then flattened, even, smooth, *commonly dry*, margin at first involute, exceeding the gills, undulated when flattened; flesh thick, hard, whitish, unchangeable. **Stem** curt, 4 cent. (1½ in.) or more long, about 12 mm. (½ in.) thick at the apex, *solid, compact*, hard, *attenuated at the base into a fusiform root*, ventricose to the neck, again attenuated upwards or wholly fusiform-attenuated, pale-white, smeared with tenacious easily dried slime, *scabrous*; flesh firm, but *externally more rigid*, cuticle somewhat fragile. Veil not conspicuous. **Gills** adnato-decurrent, acute behind, *distant, thick*, 6-8 mm. (3-4 lin.) broad, veined, tan inclining to pale.

Odour pleasant, taste sweet. The fusiform root is as long as the stem.

In mixed woods. Stoke Poges.

Edible. Name—*penus*, food. For food. *Fr. Monogr.* ii. p. 124. *Hym. Eur.* p. 406. *Sv. atl. Sv. t.* 48. *B. & Br. n.* 1778. *Ag. Schæff. t.* 238. *Sow. t.* 71.

**5. H. pulverulentus** B. & Br.—**Pileus** about 8 mm. (⅓ in.) broad, shining white, pulvinate, viscous, the involute margin tomentose. **Stem** 18 mm. (¾ in.) 2-4 mm. (1-2 lin.) thick, stuffed, nearly equal, attenuated at the very base, *wholly powdered*

with rose-coloured meal. **Gills** decurrent, thick, obtuse at the edge, whitish. Hygro-  
phorus.

Allied to *H. eburneus*, but the rose-coloured meal with which the stem is covered separates it from all other species.

Among pine-leaves. Glamis, 1876. Nov.

A small but very beautiful species. Name—*pulvis*, dust; pulverulent. *B. & Br. n.* 1667. *S. Mycol. Scot. n.* 515.

\*\* Reddish.

**6. H. erubescens** Fr.—**Pileus** 5–10 cent. (2–4 in.) and more broad, white becoming everywhere red, fleshy, gibbous then convexo-plane, viscid, *adpressedly dotted with squamules or becoming smooth*, sometimes wholly compact, sometimes thin towards the margin which is at the first naked; flesh firm, white. **Stem** sometimes short, robust, 5 cent. (2 in.) long, 2.5 cent. (1 in.) thick and attenuated upwards, sometimes elongated, 10 cent. (4 in.) long, equal or attenuated at the base, *solid, flexuous, with red fibrils, dotted with red upwards*. **Gills** decurrent, distant, *soft, white, with red spots*.

Veil none. The ground colour is white, as it is also internally, but it everywhere becomes red and the pileus often rosy-blood-colour. Handsome, growing in troops, commonly forming large lax circles.

In pine woods. Hereford.

Spores ellipsoid, very obtuse at both ends, 8–10 × 4–5 mk. *K.* Name—*erubescens*, to become red. *Fr. Monogr. ii. p.* 125. *Hym. Eur. p.* 407. *Sv. atl. Sv. t.* 65. *Quel. t. II. f. I.* *Grevillea, vol. vi. p.* 71. *Ag. Kalchbr. Ic. t.* 18. *f. 2.*

**7. H. glutinifer** Fr.—**Pileus** *rufescent*, whitish round the margin, fleshy, convexo-expanded, thin with exception of the disc which is at length broadly gibbous, *with a glutinous pellicle, disc wrinkled-dotted*. **Stem** *stuffed*, somewhat elastic, *ventricose downwards, with a viscous veil*, of the same colour as the pileus, white-squamulose at the apex. **Gills** arcuato-decurrent, rather thick, shining white.

Gregarious, mild, variable in stature.

In woods. Penzance.

Fries considers it different from *A. aromaticus* *Sow. t.* 144. Berkeley (*B. & Br. n.* 2019) thinks it is probably the same. Name—*gluten*, *fero*, to carry. Gluten-bearing. *Fr. Hym. Eur. p.* 407. *Epicr. p.* 322. *B. & Br. n.* 2019. *Ag. Bull. t.* 258, 539. *f. B.*

\*\*\* Tawny or light-yellow.

**8. H. arbustivus** Fr.—**Pileus** 5 cent. (2 in.) broad, slightly tawny-brick-colour, paler round the margin, opaque, fleshy, con-

Hygro-  
phorus.

vexo-plane, obtuse, somewhat repand, viscid, *streaked with innate fibrils*, but wholly smooth, even. **Stem** 4 cent. ( $1\frac{1}{2}$  in.) long, 12 mm. ( $\frac{1}{2}$  in.) thick, solid, equal, incurved, *with white mealy granules at the apex*, smooth at the base, pale white, externally more rigid and polished, internally somewhat spongy, hence elastic. **Gills** adnate, scarcely decurrent, distant, thick, white.

Odour and taste not unpleasant. It differs from all neighbouring species in the *mealy free* (not innate) *white granules at the apex of the stem*. In structure it is nearest to *H. pratensis*.

In woods. Colleyweston, &c. Dec.

Perhaps edible. Name—*arbustum*, a plantation. Found under trees. *Fr. Monogr.* ii. p. 126. *Hym. Eur.* p. 408. *B. & Br. n.* 932, 1013. *C. Hbk. n.* 556.

**9. H. discoideus** Pers.—**Pileus** 2.5–5 cent. (1–2 in.) broad, *pale yellowish inclining to pale, the disc always darker somewhat ferruginous*, slightly fleshy with exception of the firmer disc, on its first appearance campanulato-convex, margin inflexed, then rather plane and somewhat obtuse, and at length depressed at the disc, even, smooth, *very glutinous*; flesh of the same colour as the gills. **Stem** 4–5 cent. ( $1\frac{1}{2}$ –2 in.) long, 6–10 mm. (3–5 lin.) thick, stuffed, equal or attenuated upwards, very *soft* (texture fibrous, separable), *flocculose, viscid*, pale-white, *with white dots at the apex*. Veil entirely glutinous, rarely terminating somewhat in the form of a ring on the stem. **Gills** at first adnate, somewhat crowded, then decurrent, distant, moderately thin, soft, pale, yellowish-white.

Gregarious, here and there forming rings. In later autumn it occurs with the pileus clay-coloured, disc ferruginous, stem elongated, at length hollow.

In grassy places. Laxton Park, Norths. Oct.

Stem dotted all over with viscid granules. *B. & Br.* Spores ellipsoid-sphaeroid,  $5\text{--}6 \times 4\text{--}5$  mk. *K.* Name—*δίσκος*, *εἶδος*. Disc-shaped. *Fr. Monogr.* ii. p. 127. *Hym. Eur.* p. 408. *B. & Br. n.* 1558. *Ag. Pers.*—*Gonn. & Rab.* viii.-ix. t. 10 f. 4.

\*\*\*\* *Olivaceous-umber.*

**10. H. limacinus** Fr.—**Pileus** 4–6 cent. ( $1\frac{1}{2}$ –2½ in.) broad, *disc umber then fuliginous*, paler round the margin, fleshy, convex then flattened, obtuse, smooth, viscid; flesh rather firm, white. **Stem** 5–7.5 cent. (2–3 in.) long, 12 mm. ( $\frac{1}{2}$  in.) thick, *solid*, firm, ventricose, *viscous*, flocculose, fibrilloso-striate, *roughened with squamules at the apex*. **Gills** adnate then decurrent, somewhat distant, thin, *white inclining to cinereous*.

Veil entirely viscous, not floccose. Intermediate between *H. olivaceo-albus*

and *H. agathosmus*, differing from the former in the stem being squamulose at the apex, and from the latter in the *manifest veil* being cortinate. Hygro-  
phorus.

In woods among damp leaves. St Leonards. Oct.

Name—*limax*, a slug; slimy. *Fr. Monogr.* ii. p. 129. *Hym. Eur.* p. 409. *B. & Br. n.* 1277. *Saund. & Sm. t.* 28. *Ag. Scop.*

**11. *H. olivaceo-albus* Fr.—Pileus** 2.5–5 cent. (1–2 in.) broad, *olivaceous-fuscous*, becoming pale, especially towards the margin, which is at the first naked, fleshy, *at first acorn-shaped, then expanded, umbonate*, and at length depressed round the umbo, wholly even, smooth, *glutinous*; flesh thin, white. **Stem** about 7.5 cent. (3 in.) long, 6 mm. (3 lin.) thick, *solid*, equal or attenuated at the very base, *squamulose and spotted with the fuscous veil*, which is in the form of an adnate sheath, *viscid*, but the ground colour is white; veil terminating near the apex in the form of a ring above which the stem is even, shining white. **Gills** decurrent, *distant*, broad, simple, connected by veins at the base, *always shining white*.

Margin somewhat striate when old. Scattered throughout the summer in rainy weather, not enduring cold, and hence very different in its nature from *H. hypothejus*, which is allied.

In woods, chiefly pine. Frequent. Aug.–Oct.

Spores  $4 \times 7$  mk. *W.G.S.* Name—from the *olivaceous* pileus and *white* gills. *Fr. Monogr.* ii. p. 128. *Hym. Eur.* p. 410. *Berk. Out.* p. 198. *C. Hbk. n.* 558. *S. Mycol. Scot. n.* 517. *Ag. Brig. t.* 5. f. 1-3.

**12. *H. hypothejus* Fr.—Pileus** 2.5–5 cent. (1–2 in.) broad, *at first smeared with olivaceous gluten*, cinereous, when the gluten disappears, becoming pale and yellowish, orange or rarely (when rotting) rufescent, fleshy, *thin*, convex then depressed, *obtuse*, even, somewhat streaked; flesh thin, white then becoming light yellow. **Stem** 5–10 cent. (2–4 in.) long, 4–6 mm. (2–3 lin.) and more thick, *stuffed*, equal, *even, viscous*, but rarely spotted with the veil, at length hollow. Partial *veil* floccose, at the first *cortinate and annular, soon fugacious*. **Gills** decurrent, *distant*, distinct, at first pallid (even whitish), soon *yellow*, sometimes flesh-colour.

Very protean, changeable in colour and variable in size. Stem not scabrous. There is no trace of the veil when the plant is full grown. Appearing after the first cold autumn nights, and lasting even till snow.

In pine woods. Frequent. Sept.–Nov.

Spores subellipsoid,  $8-9 \times 4-5$  mk. *K.*;  $12 \times 4$  mk. *W.G.S.* Name—*ὕψος*; *θειον*, sulphur. From the sulphur-colour under the gluten. *Fr. Monogr.* ii. p. 128. *Hym. Eur.* p. 410. *Berk. Out.* p. 198. *C. Hbk. n.* 557. *S. Mycol. Scot. n.* 518. *Ag. Krombh. t.* 72. f. 24, 25. *Gonn. & Rab. t.* 10. f. 5. *Sow. t.* 8. *Alb. & Schw. t.* 10. f. 4. *Buxb. C. iv. t.* 21.

Hygro-  
phorus.

**13. H. cerasinus** Berk.—**Pileus** 4–6 cent. ( $1\frac{1}{2}$ – $2\frac{1}{2}$  in.) broad, pale *umber* then grey, fleshy, convex, broadly umbonate, often more or less wavy, sometimes depressed, viscid, shining when dry, *margin minutely tomentose*. **Stem** 2.5–5 cent. (1–2 in.) long, solid, white, attenuated below, punctato-squamulose above. **Gills** decurrent, broad, white, tinged with pink, sometimes forked, very distant.

Smell like that of cherry-laurel leaves. Somewhat gregarious. The minute white down at the margin of the pileus is divided into little linear heaps by the pressure of the gills in the early stage of growth.

In woods, fir, &c. Rare. Oct.

Name—from its smell of cherry-laurel (*Prunus laurocerasus*). Berk. *Eng. Fl.* v. p. 12. *Out.* p. 197. C. *Hbk.* n. 553. S. *Mycol. Scot.* n. 519. Fr. *Hym. Eur.* p. 410.

\*\*\*\* *Fuscos-cinereous or livid.*

**14. H. fusco-albus** Fr.—**Pileus** about 5 cent. (2 in.) broad, fuscous then cinereous, fleshy, moderately firm, convexo-plane, even, smooth, viscous, margin *white-floccose*. **Stem** 5–7.5 cent. (2–3 in.) long, 8–12 mm. (4–6 lin.) thick, solid, equal, *when dry white-floccose at the apex*. **Gills** decurrent, broad, rather thick, snow-white.

Stature and veil those of *H. chrysodon*, but the veil is somewhat annular, floccose not squamulose.

Among moss. Gwrwch, 1881.

Remarkable for its distinct floccose veil. M. *J.B.* Name—from the *fuscous* pileus and *white* gills. Fr. *Hym. Eur.* p. 410. B. & Br. n. 1962. Ag. *Fung. Linn.* v. t. 6. f. 1.

**15. H. agathosmus** Fr.—**Pileus** 4–6 cent. ( $1\frac{1}{2}$ – $2\frac{1}{2}$  in.) broad, livid-grey, unicolorous, *dotted with minute, raised, crowded, viscous, pellucid, whitish papillæ*, fleshy, convex then plane, gibbous, viscous, margin at first involute and villous, soon unfolded and naked, at length reflexed and undulated; flesh soft, watery, whitish. **Stem** 5–7.5 cent. (2–3 in.) long, 6–10 mm. (3–5 lin.) and more thick, solid, firm, at length soft (also hollow), equal or slightly thickened downwards, *somewhat fibrilloso-striate*, not viscous, *roughened as if mealy upwards with white punctiform squamules*, which at length become cinereous. Partial veil not conspicuous. **Gills** decurrent, distant, soft, 6–8 mm. (3–4 lin.) broad, quite entire, somewhat veined at the base, shining white.

A very remarkable species. Odour chiefly when older sweet, of anise. Lasting into late autumn. The pileus when luxuriant is much larger and



flexuous. The stem varies curved, flexuous, &c. The squamules take the place of the veil. Hygro-  
phorus.

In fir woods. Forres. Glamis. Sept.-Nov.

Spores sphaeroid-ellipsoid, hyaline,  $8 \times 5$  mk. *K.* Name—ἀγαθός, good; ὀσμὴ, scent. From the agreeable scent. *Fr. Monogr.* ii. p. 129. *Hym. Eur.* p. 411. *S. Mycol. Scot.* n. 520. *Gonn. & Rab.* viii.-ix. t. 11. f. 4.

**16. H. mesotephros** B. & Br.—**Pileus** about 2.5 cent. (1 in.) broad, white, *disc brown*, convex, somewhat hemispherical, viscid, striate, the extreme margin often remaining quite even; flesh white, *hygrophanous*. **Stem** about 5 cent. (2 in.) long, 4 mm. (2 lin.) thick, stuffed with a fibrillose pith, slender, flexuous, attenuated at the base, white, viscid, *floccoso-granulated at the apex*. **Gills** shortly decurrent, moderately broad, ventricose, rather distant, pure white.

A very delicate species, allied to *H. fusco-albus*, but with a very different habit. In age the lower part of the stem is slightly stained, but by no means squamose.

In woods. Rare. Bowood.

Spores 9 mk. long. *B. & Br.*; elliptical, pruniform, 10-12 mk. *Q.* Name—μέσος, middle; τέφρος, ash-coloured. From the dark disc. *B. & Br. Ann. Nat. Hist. ser. 2. vol. xiii. t. 15. f. 2. Berk. Out. p. 198. C. Hbk. n. 555. Fr. Hym. Eur. p. 411.*

**17. H. livido-albus** Fr.—**Pileus** *livid, unicolorous*, fleshy, thin, obtuse, *even, smooth*, viscid, margin naked. **Stem** stuffed, slender, equal, rather even. **Gills** decurrent, distant, distinct, shining white.

It differs from *H. eburneus* not only in the livid colour but also in the slightly viscid pileus as well as stem being even, smooth. It approaches the *Camarophylli*.

In woods. Street. Oct.

Name—from the *livid* pileus and *white* gills. *Fr. Hym. Eur. p. 412. Monogr. ii. p. 131. B. & Br. n. 1357. Ag. Fl. Dan. t. 1907. f. 2.*

## II.—CAMAROPHYLLUS.

\* *Gills deeply and at length obconically decurrent.*

**18. H. caprinus** Fr.—**Pileus** 5-10 cent. (2-4 in.) and more broad, at first *blackish-fuliginous*, or blackish, at length cinereous-fuliginous, but varying azure-blue, &c., fleshy, at first conico-convex, often with a prominent acorn-shaped umbo, then flattened or depressed, obtuse, somewhat repand, at the first slightly viscid, *streaked with fibrils*; flesh soft, fragile, thin towards the margin which is at length repand. **Stem** 7.5-10 cent. (3-4 in.) long, 4

Hygro-  
phorus.

cent. (1½ in.) and more thick, solid, equal or attenuated downwards, at length soft, often hollow and fragile at the apex, *externally in a remarkable degree longitudinally fibrillose*, fuliginous, pubescent and whitish at the base. Veil none. **Gills remarkably decurrent, exceedingly distant, very broad, thick**, quite entire, at first white, at length waxy-soft, becoming glaucous.

A most distinct species, in habit to be compared only with *Ag. elixus*. The gills are sometimes altogether eaten away by slugs.

In pastures. Bath. Glamis. Oct.

Spores ellipsoid-sphaeroid, 6-8 × 4-5 mk. *K.*; ellipsoid, guttate, 10 mk. *Q.* Name—*caper*, a goat. From the streaking of the fibrils. *Fr. Monogr.* ii. p. 131. *Hym. Eur.* p. 412. *B. & Br. n.* 1278. *S. Mycol. Scot. n.* 521. *Ag. Scop.—Krombh. t.* 72. *f.* 21-23.

**19. H. leporinus** Fr.—**Pileus** 2.5-5 cent. (1-2 in.) broad, yellow-rufescent, varying fuscous, equally fleshy, convex, gibbous, equal, *fibrilloso-floccose*, opaque. **Stem** short, stuffed, firm, rigid, attenuated commonly downwards, more rarely upwards, *fibrillose*, pallid. **Gills** decurrent, pale yellowish.

Its appearance is wholly that of *H. pratensis*.

In woods and on downs. Rare. Sept.-Oct.

Spores pale umber. *M. f. B.* Name—*lepus*, a hare. From the colour. *Fr. Hym. Eur.* p. 412. *Berk. Out. p.* 199. *C. Hbk. n.* 559. *S. Mycol. Scot. n.* 522. *Ag. Schæff. t.* 313.—*Batt. t.* 9. *f. B.*

**20. H. nemoreus** Fr.—**Pileus** somewhat orange, *equally fleshy*, convex then expanded, gibbous and at length depressed, somewhat smooth. **Stem** stuffed, firm, *squamulose, fibroso-striate*, attenuated at the base. **Gills** decurrent, thick, distant, somewhat of the same colour.

In woods. Stoke Poges. Dec.

Spores white. *B. & Br.*; ovate-spherical, punctate, 5-6 mk. *Q.* Name—*nemus*, a grove. Found in wooded groves. *Fr. Hym. Eur. p.* 413. *B. & Br. n.* 1668.

**21. H. pratensis** Fr.—**Pileus** 2.5-5 cent. (1-2 in.) and more broad, somewhat pale yellowish, *compactly fleshy especially at the disc, thin towards the margin*, convex then flattened, *almost turbinate* from the stem being thickened upwards, even, smooth, moist (but not viscous) in rainy weather, when dry often rimosely incised, here and there split regularly round; flesh firm, white. **Stem** 4-5 cent. (1½-2 in.) long, 12 mm. (½ in.) and more thick, *stuffed*, internally spongy, externally polished-evened and firmer, *attenuated downwards, even, smooth*, naked. **Gills remarkably**

*decurrent*, at first *arcuate*, then *extended in the form of an inverted cone*, very distant, thick, firm, brittle, connected by veins at the base, very broad in the middle, of the same colour as the pileus. Hygro-  
phorus.

Very protean. Veil none. The flesh of the pileus is formed as it were of the stem dilated upwards. The typical form resembles the *Cantherelli*. Everywhere becoming light yellow-tawny, but varying with the stem and gills pale-white.

In pastures. Common. Aug.—Nov.

Edible; very little distinctive flavour. Spores sphaeroid-ellipsoid, 6-10 × 4-6 mk. *K.*; 5 × 4 mk. *W.G.S.* Name—*pratense*, meadow. Found in meadows. *Fr. Monogr.* ii. p. 132. *Hym. Eur.* p. 413. *Sv. atl. Sv. t.* 30. *Berk. Out.* p. 199. *C. Hbk.* n. 560. *S. Mycol. Scot.* n. 523. *Hussey* ii. t. 90. *Ag. Pers.*—*Fl. Dan.* t. 1735. f. 1. *Grev.* t. 91. *Krombh.* t. 43. f. 7-10. *Bull.* t. 587. Forms departing from the type: *Bolt.* t. 56. *Sow.* t. 141. *Pers. Myc. Eur.* t. 28. f. 1. *Brig.* t. 22.

\* **H. cinereus** Fr.—The thinner pileus and gills *cinereous*, margin at length striate. Stem white.

Otherwise as in *H. pratense*. Spores ovoid-spherical, 6 mk. *Q.* Name—from the *cinereous* colour. *Fr. Hym. Eur.* p. 413. *Sv. atl. Sv. t.* 30 (two upper fig.) *B. & Br.* n. 1561. *S. Mycol. Scot.* n. 523.

\* **H. pallidus** B. & Br.—Pileus pallid, infundibuliform, margin undulated, deflexed. Stem dilated upwards, fibrilloso-striate. Gills distant, decurrent, branched, pallid.

Coed Coch. Oct. *B. & Br.* n. 1356\*. *Fr. Hym. Eur.* p. 413.

22. **H. foetens** Phill.—Pileus about 2.5 cent. (1 in.) broad, dark brown, somewhat fleshy, convex then becoming plane, smooth, at length broken up into squamules. Stem 2.5-7.5 cent. (1-3 in.) long, 4 mm. (2 lin.) thick, stuffed, attenuated downwards, shining, clothed with transversely arranged fibrous scales, paler than the pileus. Gills decurrent, distant, rather thick, somewhat of the same colour as the pileus or paler, somewhat glaucous-pruinose.

Fragile. Very fetid, nauseous. The odour is very similar to that of *Thelephora fastidiosa*. Not at all viscid.

On the ground. Shrewsbury, 1878, 1884. Nov.

Nearly related to *H. micans*. B. & Br. Name—*foetens*, stinking. *Phill. Grevillea*, vol. vii. p. 74. *B. & Br.* n. 1780.

23. **H. virgineus** Fr. Wholly white.—Pileus fleshy, convex then plane, obtuse, moist, at length depressed, cracked into patches, floccose when dry. Stem curt, stuffed, firm, attenuated at the base, externally becoming even and naked. Gills decurrent, distant, rather thick.

Flesh sometimes equal, sometimes abruptly thin. Commonly confounded

Hygro-  
phorus.

with *H. niveus*, but it is more difficult to distinguish it from white forms of *H. pratensis*. It is distinguished chiefly by its smaller stature, by the colour being constantly white, sometimes becoming pale, by the *obtuse pileus* being scarcely turbinate, *at length cracked into patches and floccose when dry*, and by the gills being thinner, &c.

In pastures. Common. Aug.—Nov.

Taste like *Marasmius oreades*. *M. J. B.* Edible; delicious when broiled or stewed. Name—*virgo*, a virgin. From the purity of the colours. *Fr. Monogr.* ii. p. 133. *Hym. Eur.* p. 413. *Berk. Out.* p. 199. *C. Hbk. n.* 561. *S. Mycol. Scot. n.* 524. *Price f.* 41 (ochrey-white). *Ag. Wulf. in Jacqu. Miscell.* 2. t. 15. f. 1. *Sow. t.* 32. *Grev. t.* 166. *Gonn. & Rab. t.* 10. f. 3. *Bull. t.* 188. *Batt. t.* 12. f. *H.—Batsch f.* 200 (a very small form).

**24. *H. ventricosus* B. & Br.** White.—**Pileus** 5–7.5 cent. (2–3 in.) broad, fleshy, unequal, convex. **Stem** 6 cent. (2½ in.) long, 12 mm. (½ in.) thick in the middle, solid, at length partially hollow, attenuated at the base and apex. **Gills** deeply decurrent, narrow, sometimes forked.

Often tinged with red from the growth of a little *Fusisporium*.

Among grass. Coed Coch.

Name—*venter*, the belly. From the ventricose stem. *B. & Br. n.* 1777.

**25. *H. niveus* Fr.** Wholly white.—**Pileus** scarcely reaching 2.5 cent. (1 in.) broad, *somewhat membranaceous*, and without a more compact disc, hence truly *umbilicate*, campanulate then convex, smooth, striate and viscid when moist, not cracked when dry; flesh thin, everywhere equal, white, hygrophanous. **Stem** 5 cent. (2 in.) or a little more long, 2–4 mm. (1–2 lin.) thick, *fistulose, equal*, even, smooth, tense and straight. **Gills** decurrent, *distant, thin*, scarcely connected by veins, arcuate, quite entire.

Thinner, *tougher*, and later than *H. virgineus*, &c. Being hygrophanous the pileus is *shining white* when dry. Very tender forms occur.

In pastures. Common. Sept.—Oct.

Name—*niveus*, snow-white. *Fr. Monogr.* ii. p. 133. *Hym. Eur.* p. 414. *Berk. Out.* p. 199. *C. Hbk. n.* 562. *S. Mycol. Scot. n.* 525. *Ag. Scop.—Schæff. t.* 232. *Krombh. t.* 25. f. 1-3.

**26. *H. Wynnii* B. & Br.** Lemon-yellow, hygrophanous.—**Pileus** thin, umbilicate, or somewhat infundibuliform, striate. **Gills** decurrent, narrow, thin.

Fœtid when decayed, losing much of its lemon colour when it parts with its moisture. Resembling *Ag. mollis* Bull., but on a smaller scale.

On chips, old stumps, &c. Coed Coch, &c.

Name—after Mrs Lloyd Wynne. *B. & Br. n.* 1781, 1962\*. *Sm. Gard. Chron. Oct.* 1878, p. 476. *fig.* Clitocybe xanthophylla, *Bresadola Fung. Trid. fig.*

**27. H. russo-coriaceus** B. & Br.—**Pileus** about 12 mm. ( $\frac{1}{2}$  in.) Hygro-  
phorus. broad, ivory-white, fleshy, convex, slightly viscid. **Stem** 18 mm.—2.5 cent. ( $\frac{3}{4}$ –1 in.) long, not 2 mm. (1 lin.) thick, slender, *solid*, thickened upwards, smooth, pure white. **Gills** decurrent, arched, broad, thick, very few and distant, with a few shorter ones intermixed.

Known at once by its persistent *delightful odour*, like that of Russian leather or *Potentilla atro-sanguinea*.

In pastures. Rare. Oct.

Spores  $11 \times 6$  mk. *W.G.S.* Name—from its odour of Russian leather. *B. & Br. n.* 332. *Berk. Out. p.* 199. *C. Hbk. n.* 563. *S. Mycol. Scot. n.* 526. *Fr. Hym. Eur. p.* 414. *Saund. & Sm. t.* 28. *f.* 2.

**28. H. micaceus** B. & Br.—**Pileus** 12–8 mm. ( $\frac{1}{2}$ – $\frac{1}{3}$  in.) broad, hemispherical, at first light yellow then becoming cinereous, wrinkled, micaceous. **Stem** 18 mm. ( $\frac{3}{4}$  in.) long, 2 mm. (1 lin.) thick, solid, granulated, light yellow then brown downwards. **Gills** decurrent, pallid umber.

Mycelium white. The whole plant turns dark brown when dry. When young it looks like a small specimen of *Leotia lubrica*.

On clayey soil. Coed Coch, 1878. Oct.

Name—*mica*, grain, granule. From the micaceous pileus. *B. & Br. n.* 1779.

**\*\* Gills ventricose, sinuato-arcuate or plano-adnate.**

**29. H. fornicatus** Fr.—**Pileus** 2.5 cent. (1 in.) high and broad, white or pallid livid, fleshy, but thin, *campanulate then expanded*, obsoletely umbonate, somewhat repand, viscid, *even, smooth*. **Stem** 5–7.5 cent. (2–3 in.) long, 6–8 mm. (3–4 lin.) thick, *solid* or hollow at the very apex, compact, firm, tough, equal, somewhat undulated, smooth, shining white. **Gills** *sometimes almost free*, sometimes slightly adnexed with a small decurrent tooth (*sinuato-adnexed*), *thick, ventricose, distant*, distinct, exceeding the margin, white.

The structure of the stem and pileus is that of *H. pratensis*, but it is wholly white, and the attachment of the gills is plainly different. It has occurred with the stem fuscous-squamulose at the base.

In pastures. Holme Lacy, Herefordshire. Batheaston. Oct.

Name—*fornix*, a vault. Arched. *Fr. Monogr. ii. p.* 133. *Hym. Eur. p.* 414. *B. & Br. n.* 1420. *Krombh. t.* 25. *f.* 4, 5.

**30. H. distans** Berk.—**Pileus** about 5 cent. (2 in.) broad, white, with a silky lustre, here and there stained with brown, somewhat fleshy, plane or depressed, viscid. **Stem** white above, *cinereous*

Hygro-  
phorus.

below, and attenuated, not spotted. **Gills** decurrent, *few, very distant*, somewhat ventricose, pure white then tinged with cinereous, interstices obscurely rugose.

Often umbilicate. Remarkable for the few and distant gills.

In woods. Rare. King's Cliffe.

Name—from the *distant* gills. *Berk. Out. p. 200. t. 13. f. 1. C. Hbk. n. 564. Fr. Hym. Eur. p. 415.*

**31. H. Clarkii** B. & Br.—**Pileus** *livid-cinereous*, convex, somewhat umbonate, viscous, margin even. **Stem** hollow, of the same colour as the pileus. **Gills** adnate, broad, when larger nearly 12 mm. ( $\frac{1}{2}$  in.) broad, distant, thick, white.

Fragile.

In pastures. Street. Perth Fungus Show. Sept.—Oct.

Name—after J. A. Clark. *B. & Br. n. 1358. S. Mycol. Scot. n. 527. Fr. Hym. Eur. p. 415.*

**32. H. metapodius** Fr.—**Pileus** 4–7.5 cent. ( $1\frac{1}{2}$ –3 in.) broad, cinereous-fuscos, *compactly fleshy*, convex then plane, obtuse, at first even and slightly shining, *then silky and squamulose*, irregular; flesh 12 mm. ( $\frac{1}{2}$  in.) thick, pallid-grey, reddish when broken and at length becoming black. **Stem** 2.5–5 cent. (1–2 in.) long, 12 mm. ( $\frac{1}{2}$  in.) and more thick, stuffed, *attenuated downwards*, unequal, ascending, smooth, *cinereous, reddish internally*. **Gills** variously adnate (for the most part arcuato-decurrent behind, or broadly emarginate), distant, thick, veined, about 6 mm. (3 lin.) broad, grey-white.

*Odour of new meal*, taste mild. In respect of its flesh being compact, reddish when broken and at length becoming black, and its thick juicy gills, it is very analogous with *Russula nigricans*. There is a var. *paradoxa* appearing among taller mosses; stem elongated to 10 cent. (4 in.), wholly fusiform, pileus smaller, regular and villous, gills obconico-decurrent as in *H. pratensis*.

In old pastures. Street, 1871. Glamis, 1876. Aug.—Nov.

Name—from the stem (*πούς*) being as it were upside down (*μετά*). *Fr. Monogr. ii. p. 135. Hym. Eur. p. 415. B. & Br. n. 1359. S. Mycol. Scot. n. 528.*

**33. H. ovinus** Fr.—**Pileus** 5 cent. (2 in.) broad, *fuscous, slightly fleshy, campanulate then expanded*, somewhat umbonate, at first slightly viscid and even, then dry and *squamulose*, at length revolute, undulated, rimosely incised; flesh only 4 mm. (2 lin.) thick, rigid-fragile. **Stem** about 5 cent. (2 in.) long, 6 mm. (3 lin.) thick, *somewhat stuffed, somewhat equal* or slightly thickened at both ends, curved or twisted, compressed, smooth, slightly shin-

ing, pallid. **Gills** arcuato-adnate, decurrent with a tooth, distant, thick, as much as 6 mm. (3 lin.) broad, connected by veins and occasionally divided, grey then rufescent, the thin edge quite entire. Hygro-  
phorus.

Very much allied to *H. metapodius*, odour almost the same. The flesh is the same except in being thinner and rigid-fragile. In a larger var. the pileus becomes black and the stem fuscous-black.

In pastures. Uncommon. Sept.

Name—*ovis*, a sheep. Woolly, squamulose. *Fr. Monogr.* ii. p. 135. *Hym. Eur.* p. 415. *Berk. Out.* p. 200. *C. Hbk. n.* 565. *S. Mycol. Scot. n.* 529. *Hussey ii. t.* 50. *Ag. Bull. t.* 580.

**34. H. subradiatus** Fr.—**Pileus** varying in colour, white, livid, disc fuscous, &c., somewhat membranaceous, slightly fleshy at the disc from the apex of the stem being dilated, otherwise quite membranaceous, but moderately tough and somewhat irregularly shaped, convex then expanded, obsoletely umbonate, hygrophanous when moist, *radiato-striate* from the translucent gills. **Stem** 5 cent. (2 in.) long, 4 mm. (2 lin.) thick, *fistulose, equal*, attenuated at the base, twisted, ascending, *smooth*, pale, white at the base. **Gills** plane but deeply decurrent with a tooth, ventricose, somewhat thin, distant, connected by veins, white.

Allied to the thinner forms of *H. pratensis*, but differing very widely in the attachment of the gills.

In old pastures. Glamis, 1874. Sept.

Name—*sub*, and *radiatus*, rayed. From the character of the striæ. *Fr. Monogr.* ii. p. 136. *Hym. Eur.* p. 416. *B. & Br. n.* 1670. *S. Mycol. Scot. n.* 530. *Ag. Schum.*

\* **H. lacmus** Fr.—**Pileus** 2.5 cent. (1 in.) broad, *lilac* then becoming pale, fragile, slightly fleshy at the disc, otherwise membranaceous, plano-depressed, occasionally at the same time umbonate, unequal at the circumference, *radiato-striate when moist*, when dry even, shining, commonly smooth, but occasionally fibrillose or squamulose on the disc. **Stem** elongated to 5 cent. (2 in.) and more, stuffed then hollow, often twisted, smooth, somewhat shining, narrowed at the base and apex. **Gills** plano-decurrent, thin, 2 mm. (1 lin.) broad, connected by veins, somewhat distant, cinereous.

In the same habitats as *H. subradiatus*. In pastures and heathy places. Coed Coch. Loch Maree, &c. Sept.-Oct. Name—*lac*, varnish. Shining. *Fr. Monogr.* ii. p. 136. *Hym. Eur.* p. 416. *B. & Br. n.* 1780. *S. Mycol. Scot. n.* 530. *Fl. Dan. t.* 1731. f. 1.

**35. H. irrigatus** Fr.—**Pileus** 2.5-5 cent. (1-2 in.) broad, *livid*,

Hygro-  
phorus.

becoming dingy white as the gluten separates, slightly fleshy, fragile, campanulate, expanded, obtuse, even, smooth, *viscous*. **Stem** 5-7.5 cent. (2-3 in.) long, 2-4 mm. (1-2 lin.) thick, fistulose, flexuous, *tough*, equal, smooth, *very viscous*, *livid*. **Gills** *adnate*, with a *decurrent tooth*, somewhat distant, distinct, fragile, rather thick, scarcely connected by veins, either wholly white or grey.

Very variable, sometimes much larger. The gills are less juicy than those of *H. unguinosus*, with which it must not be confounded.

In grassy places and fir woods. Laxton, Norths. Glamis. Oct.-Nov.

Name—*irrigio*, to wet. From its *watery* texture. *Fr. Monogr.* ii. p. 137 (*irriguus*). *Hym. Eur.* p. 416. *Icon. t.* 168. f. 3. *B. & Br. n.* 1280. *S. Mycol. Scot. n.* 531. *Ag. Pers.*

**36. H. Houghtoni** B. & Br.—**Pileus** 4-5 cent. (1½-2 in.) broad, bright coloured, convex, at length depressed in the centre, striate, *very viscous*. **Stem** 5 cent. (2 in.) and more long, 6 mm. (¼ in.) thick, transversely undulated, very viscous, tawny-light yellow. **Gills** *decurrent*, thin, pale yellowish.

Odour foxy. The stem is sometimes tinged above with blue. *The gelatinous coat is extremely thick*, and at length separates and forms a cup in the centre.

Among grass. Preston, Salop, 1872. Oct.

Name—after Rev. William Houghton. *B. & Br. n.* 1360. *Fr. Hym. Eur.* p. 416.

### III.—HYGROCYPE.

#### \* *Gills decurrent.*

**37. H. Colemannianus** Blox.—**Pileus** 2.5-5 cent. (1-2 in.) broad, reddish *umber*, paler when dry except in the centre, somewhat fleshy, at first somewhat campanulate, at length expanded, strongly umbonate, when moist striate and slightly viscid. **Stem** 2.5 cent. (1 in.) or more long, 2-4 mm. (1-2 lin.) thick, brittle, fibrous, nearly equal, *somewhat silky*, white, very slightly tinged with umber. **Gills** deeply *decurrent*, rather broad, distant, interstices veined and rugose, umber but paler than the pileus.

In grassy places. Twycross, Warwickshire.

Spores 6 × 8 mk. *B. & Br.* Name—from W. H. Coleman. *Blox.—Berk. Out. p.* 200. *B. & Br. n.* 701. *C. Hbk. n.* 566. *Fr. Hym. Eur. p.* 417.

**38. H. sciophanus** Fr.—**Pileus** 1-4 cent. (½-1½ in.) broad, *very hygrophanous with a peculiar deep tawny or rather vivid*



*brick but opaque colour*, becoming pale when dry, becoming white when older, thin, slightly fleshy at the disc, otherwise membranaceous, hemispherical or obtusely campanulate then expanded, always obtuse, slightly viscid, even at the disc, remotely and pellucidly *striate to the middle*, regular, torn when older. **Stem** 4-7.5 cent. (1½-3 in.) long, hollow, wholly equal, always thin, scarcely more than 2 mm. (1 lin.) thick, flexuous, even, smooth, slippery, never tense and straight, *tawny yellowish*, but paler than the pileus, even becoming white. **Gills** *attenuato-adnate*, at the first slightly ascending, then plano-decurrent, distant, commonly distinct, of the same colour as the pileus, or sometimes inclining to rose-colour. Hygro-  
phorus.

Very elegant, densely gregarious, fragile. In habit and opaque tawny colour it is like *H. pratensis*, but is thin, fragile, juicy, with the pileus somewhat repand, and the gills thin and plano-decurrent. The stem is never tense and straight as in *H. conicus*. There is a var. with the *pileus date-brown*.

In mossy places. Perth.

Spores very pale clay-coloured. *B. & Br.* Name—σκιά, shade; φαίνω, to appear. From its shaded colour. *Fr. Monogr.* ii. p. 137. *Hym. Eur.* p. 417. *Icon. t.* 167. f. 1. *B. & Br. n.* 1560. *S. Mycol. Scot. n.* 532. *Ag. Fl. Dan. t.* 1845. f. 2.

**39. H. lætus** Fr.—**Pileus** reaching 2.5 cent. (1 in.) broad, *tawny, shining*, not becoming pale, slightly fleshy, almost membranaceous at the margin, convex then flattened, obtuse, slippery, even or slightly pellucid-striate round the margin; flesh thin, of the same colour as the pileus and paler. **Stem** 5-7.5 cent. (2-3 in.) long, thin, about 4 mm. (2 lin.) thick, hollow or rather fistulose, equal, *tough*, even, very smooth, slippery, *tawny*. **Gills** *somewhat decurrent*, distant, *thin*, somewhat connected by veins, for the most part *flesh-coloured*, but varying whitish, fuliginous, &c.

The stem is here and there undulated and darker at the apex, in which it approaches *H. psittacinus*. It departs from all the others in this group in the toughness, especially of the stem. Appearing in troops.

In pastures. Frequent.

Spores nearly globose, 8 mk. *B. & Br.*; ovoid pruniform, 8 mk. *Q.* Name—*lætus*, joyful, pleasing. From the bright colour. *Fr. Monogr.* ii. p. 138. *Hym. Eur.* p. 417. *Icon. t.* 167. f. 2. *B. & Br. n.* 702. *Berk. Out. p.* 200. *C. Hbk. n.* 567. *S. Mycol. Scot. n.* 533. *Ag. Pers.*

**40. H. vitellinus** Fr.—**Pileus** 1-2.5 cent. (½-1 in.) broad, *citron-egg-yellow, becoming white when dry*, very thin, slightly fleshy only at the disc, otherwise almost membranaceous, smooth, viscid, *margin plicato-striate*. **Stem** 5 cent. (2 in.) long, scarcely

Hygro-  
phorus.

2 mm. (1 lin.) thick, fistulose, equal, even, smooth, *pallid light yellow, becoming white when dry*, very fragile, here and there flexuous. **Gills** deeply decurrent, somewhat distant, egg-yellow.

It is certainly different from *H. ceraceus*, which is somewhat like it, in the colour being wholly bright citron-yellow, whitish when dry, in the margin of the pileus and in the deeply decurrent gills.

In pastures. Penzance.

Name—*vitellus*, yolk of egg. Egg-yellow. *Fr. Monogr.* ii. p. 312. *Hym. Eur.* p. 417. *Icon. t.* 167. *f.* 3. *B. & Br. n.* 2020.

**41. H. ceraceus** Fr.—**Pileus** about 2.5 cent. (1 in.) broad, *waxy-yellow, shining*, slightly fleshy, thin, but slightly firm, convexo-plane, obtuse, slightly pellucid-striate, viscid. **Stem** 2.5–5 cent. (1–2 in.) and more long, about 4 mm. (2 lin.) thick, *hollow, often unequal*, flexuous and at length compressed, even, smooth, of the same colour as the pileus, never darker at the apex. **Gills** *adnato-decurrent, broad, almost triangular*, distinct, yellow.

Fragile; easily distinguished from others by its waxy (not changeable) colour. Growing in troops. The gills are by no means emarginato-adnexed and ventricose as in the similar, but more watery and sulphur-coloured *H. chlorophanus*. Like neighbouring species it varies very much in stature.

In woods and pastures. Common. Sept.–Oct.

Spores sphaeroid-ellipsoid,  $7 \times 4$  mk. *K.*;  $6 \times 4$  mk. *W.G.S.* Name—*cera*, wax. From the colour. *Fr. Monogr.* ii. p. 138. *Hym. Eur.* p. 417. *Berk. Out. p.* 201. *C. Hbk. n.* 568. *S. Mycol. Scot. n.* 534. *Ag. Wulf, in Jacq. Coll. ii. t.* 15. *f.* 2. *Sow. t.* 20.

**42. H. coccineus** Fr.—**Pileus** 2.5–5 cent. (1–2 in.) and more broad, *at first bright scarlet, then soon changing colour and becoming pale*, slightly fleshy, convex, then plane and often unequal, *obtuse*, at first viscid and even, *smooth*, not floccoso-scaly; flesh of the same colour as the pileus. **Stem** 5 cent. (2 in.) long, 6–8 mm. (3–4 lin.) thick, *hollow*, then *compressed* and rather even, not slippery, *scarlet upwards, always yellow at the base*. Gills wholly adnate, *decurrent with a tooth*, plane, distant, connected by veins, watery-sect as if fatty, when full grown *purplish at the base, light yellow in the middle, glaucous at the edge*.

Flesh of the pileus descending into the gills and forming a trama of the same colour. Fragile. Varying in stature, easily mistaken for some of the following species which are of the same colour. Pileus at length becoming yellow.

In pastures. Very common. Aug.–Nov.

Spores sphaeroid-ellipsoid,  $6-8 \times 4-5$  mk. *K.*;  $4 \times 8-7$  mk. *W.G.S.* Name—*coccineus*, of scarlet colour. *Fr. Monogr.* ii. p. 139. *Hym. Eur. p.* 417.

*Berk. Out. p. 201. C. Hbk. n. 569. S. Mycol. Scot. n. 535. Hussey i. t. Hygro-*  
*61. Price f. 57. Ag. Schæff. t. 302. Bull. t. 570. f. 2 partly. Fl. Dan. phorus.*  
*t. 715. Sow. t. 381 (scarlet fig.) Batt. t. 19. f. B.*

**43. H. miniatus** Fr.—**Pileus** scarcely reaching 2.5 cent. (1 in.) broad, at first *vermilion* then *becoming pale and opaque*, slightly fleshy, convex, obtuse, then *umbilicate*, at first even and smooth, then *squamulose*. **Stem** about 5 cent. (2 in.) long, 2 mm. (1 lin.) thick, *somewhat stuffed, equal, round*, even, smooth, shining, *vermilion*. **Gills** adnate, not decurrent, plane, distant, distinct, rather thick, *yellow* or sometimes light yellow-vermilion.

Very fragile. The gills are firmer than in *H. coccineus*. Intermediate between *H. coccineus* and *H. turundus*.

In open places, pastures, &c. Common. July–Oct.

Spores 5 × 8 mk. *W.G.S.* Name—*minium*, vermilion. *Fr. Monogr. ii. p. 139. Hym. Eur. p. 418. Berk. Out. p. 201. C. Hbk. n. 570. S. Mycol. Scot. n. 536. Quel. t. 10. f. 5. Ag. Fl. Dan. t. 1009. Harz. t. 4. f. 1.*

**44. H. turundus** Fr.—**Pileus** 1–2.5 cent. ( $\frac{1}{2}$ –1 in.) broad, *yellow or tawny, variegated with cinereous-fuscos squamules*, slightly fleshy, fragile, convex then expanded, *umbilicate* or depressed, often elegantly crenate at the margin. **Stem** 5–7.5 cent. (2–3 in.) long, about 4 mm. (2 lin.) thick, rigid-fragile, stuffed then fistulose, *round, equal, even* and polished, *tawny-reddish*. **Gills** at first slightly then deeply *decurrent*, distant, juicy, distinct, at first *white* then becoming yellow, the acute edge of the same colour.

Very protean. The pileus is commonly dry, soon broken up into fuliginous flocci, but it occurs during rain at the first viscid and even, and in this condition it must be carefully distinguished (by the colour of the golden pileus and of the gills) from *H. miniatus*, to which it is allied though very dissimilar. It varies with the somewhat veined gills of the *Cantharelli*.

On peat soil. Farragon, Perthshire, 1877. Aug.

2000 feet. Name—*turunda*, lint. Flocculose. *Fr. Monogr. ii. p. 139. Hym. Eur. p. 418. B. & Br. n. 1670. S. Mycol. Scot. n. 537.*

**Var. mollis** B. & Br. Golden.—**Pileus** 12–18 mm. ( $\frac{1}{2}$ – $\frac{3}{4}$  in.) broad, nearly plane, at length slightly depressed, clad with soft, short, radiating hairs of the same colour. **Stem** 2.5–3 cent. (1–1 $\frac{1}{4}$  in.) long, 2–4 mm. (1–2 lin.) thick, stuffed, equal. **Gills** decurrent, arcuate, narrow, distant.

In moist mossy ground. Rare. July–Sept. *B. & Br. n. 1279. S. Mycol. Scot. n. 537. Fr. Hym. Eur. p. 418.*

**45. H. mucronellus** Fr.—**Pileus** 6–8 mm. (3–4 lin.) broad, when moist a) scarlet b) yellow, when dried changing colour

Hygro-  
phorus.

very much, becoming hoary, slightly fleshy, when smaller *acutely conical*, when larger *campanulate* and obtuse, then expanded, when moist pellucidly striate, when dried somewhat silky, but never squamulose. **Stem** about 5 cent. (2 in.) long, 2 mm. (1 lin.) thick, fistulose, thin, somewhat attenuated and *white at the base*, otherwise yellow, even, smooth, flexuous. **Gills** ascending, distant, very broad and adnate behind with a very broad base, somewhat decurrent, yellow.

Fragile. The smallest of species of the same nature.

In grass field. King's Lynn. Dec.

In 'Hym. Eur.' Fries describes the stem as *somewhat silky*. Name—*muco*, a sharp point. From its acute form. *Fr. Monogr.* ii. p. 140. *Hym. Eur.* p. 418. *Grevillea*, vol. iv. p. 118.

**\*\* Gills adnexed, somewhat separating.**

**46. H. puniceus** Fr.—**Pileus** 5–10 cent. (2–4 in.) broad, glittering blood-scarlet, in dry weather and when old becoming pale especially at the disc, slightly fleshy for its breadth, at first *campanulate, obtuse*, commonly repand or lobed, very irregular, even, smooth, viscid; flesh of the same colour, fragile. **Stem** 7.5 cent. (3 in.) long, 1–2.5 cent. ( $\frac{1}{2}$ –1 in.) thick, solid when young, at length hollow, very stout (not compressed), ventricose (attenuated at both ends), *striate*, and for the most part squamulose at the apex, when dry light yellowish or of the same colour as the pileus, *always white* and often incurved *at the base*. **Gills** ascending, ventricose, 4–8 mm. (2–4 lin.) broad, thick, distant, white-light yellow or yellow and often reddish at the base.

The largest of the group, and very handsome. It certainly differs from *H. coccineus*, for which it is commonly mistaken, in stature, *in the adnexed gills, and in the white base of the striate stem*. The attachment of the gills varies, but from the form of the pileus they ascend to the base of the cone and appear free.

In pastures. Frequent. July–Nov.

Name—*puniceus*, blood-red. *Fr. Monogr.* ii. p. 141. *Hym. Eur.* p. 419. *Sv. atl.* *Sv. t.* 77. *Berk. Out.* p. 201. *B. & Br. n.* 1281. *C. Hbk. n.* 571. *S. Mycol. Scot. n.* 538. *Ag. Fl. Dan. t.* 883. *Bolt. t.* 67. f. 2.

**47. H. obrusseus** Fr.—**Pileus** 5–7.5 cent. (2–3 in.) broad, golden-sulphur-yellow, shining, not becoming pale, very thinly fleshy, rigid-fragile, campanulate then expanded, flexuous, somewhat lobed and often torn, obtuse, dry, even, smooth. **Stem** 7.5 cent. (3 in.) and less in length, 12 mm. ( $\frac{1}{2}$  in.) thick or when compressed still broader, but scarcely ventricose, stout, ascending, *compressed* and commonly unequal, quite *smooth, even, sulphur*

*yellow, becoming tawny at the base. Gills* adnexed, ventricose, but at length separating, free, very broad (12 mm.,  $\frac{1}{2}$  in.), thick, distant, almost white. Hygro-  
phorus.

The stem is hollow and hence flattened, but the walls are slightly firm and somewhat cartilaginous, fragile however and easily splitting into crisped fibres. The gills are much more rigid and firm than in allied species. *A. laceratus* *Bolt. t. 68* which is referred to this species by Secretan is rather a luxuriant form of *H. conicus*.

In woods. Mossburnford. Oct.

Name—*obrussa*, the testing of gold. From the pure golden colour. *Fr. Monogr. ii. p. 141. Hym. Eur. p. 419. Berk. Out. p. 202. C. Hbk. n. 572. S. Mycol. Scot. n. 539. Ag. Batt. t. 19. f. D.*

**48. *H. conicus* Fr.—Pileus** commonly light yellow, becoming black, somewhat membranaceous, *conical, acute*, smooth, often lobed, then expanded and cracked. **Stem** hollow, cylindrical, tense and straight, *fibroso-striate*. **Gills** attenuato-free, ventricose, thin, somewhat crowded, varying white, light yellow, &c.

Very easily distinguished from all neighbouring species by its juicy and fragile substance, &c. Viscid when moist, shining when dry. It is smooth even when dry. The primary colour depends upon the weather. It has the singular character of the flesh becoming black when broken, but in some forms (springing in dry weather) it is unchangeable. The following forms may be noted : a) *coccinea*, scarlet and shining, gills yellow, often reddish at the base : b) *lutea*, pileus and gills yellow—neither becoming black : c) pileus reddish or tawny, gills yellow : d) *sulphurea*, pileus bright light yellow-sulphur-colour, sometimes greenish, gills often whitish—both becoming black when broken : e) *squalida*, pileus livid or fuliginous-light yellow : f) *tristis*, pileus at the first fuliginous, soon shining black—both of their own accord becoming wholly black.

In pastures. Very common. July–Nov.

Pileus 2.5–6 cent. (1–2  $\frac{1}{2}$  in.) Stem varying very much in length and thickness. Spores sphaeroid-ellipsoid, 10–11  $\times$  6–8 mk. *K.* ; 11  $\times$  6 mk. *W.G.S.* Name—from its *conical* shape. *Fr. Monogr. ii. p. 142. Hym. Eur. p. 419. Berk. Out. p. 202. C. Hbk. n. 573. S. Mycol. Scot. n. 540. Ag. Scop.—Schæff. t. 2. Fl. Bat. t. 310. Bull. t. 50 (524 f. 3. var. *tristis*). Batsch f. 28.*

**49. *H. calyptræformis* B. & Br.—Pileus** about 2.5 cent. (1 in.) high, 18 mm. ( $\frac{3}{4}$  in.) broad at the base, *pink, becoming pallid*, thin, acutely conical, lobed below, moist when unexpanded, striate under a lens, minutely *innato-fibrillose*. **Stem** 2.5 cent. (1 in.) or more long, hollow, the walls fibrous within, remarkably smooth, slightly striate, brittle, often splitting longitudinally, *white*. **Gills** acutely attenuated behind, very narrow, distinct, rose-coloured at length pallid.

The gills are often almost evanescent behind. The stem is pure white, except within the pileus where it has often a roseate tinge. Very distinct from *H. conicus*. It does not turn black when bruised.

Hygro-  
phorus.

In pastures. Uncommon. Aug.-Oct.

I have found several specimens larger than described by Berkeley, quite as large as full-sized *H. conicus* which it resembles very much in shape. Spores  $3 \times 4$  mk. *W.G.S.* Name—*calyptra*, a hood; *forma*, form. From its shape. *Berk. Out. p.* 202. *Trans. Woolth. Club, t.* 21. *f.* 4-6. *C. Hbk. n.* 576. *S. Mycol. Scot. n.* 541. *Fr. Hym. Eur. p.* 420.

**50. *H. chlorophanus* Fr.—Pileus** 2.5 cent. (1 in.) broad, commonly bright sulphur-yellow, sometimes however scarlet, not changing colour, somewhat membranaceous, very fragile, at first *convex*, then plane, *obtuse*, orbicular and lobed, and at length cracked, smooth, *viscid*, striate. **Stem** 5-7.5 cent. (2-3 in.) long, 4-6 mm. (2-3 lin.) thick, hollow, equal, round, rarely compressed, wholly *even*, smooth, *viscid* when moist, shining when dry, wholly unicolorous, rich light yellow. **Gills** emarginato-*adnexed*, very ventricose, with a thin decurrent tooth, thin, distant, distinct.

Very much allied to *H. conicus*, but never becoming black, and otherwise certainly distinguished by its convex, obtuse, striate pileus, by its *even* and *viscous* stem, and by its *emarginato-free*, thin, somewhat distant, whiter gills. Like *H. ceraceus* in appearance.

In grassy and mossy places. Common. Aug.-Oct.

Spores pruiniform, 8 mk. *Q.* Name—*χλωρός*, greenish-yellow; of colour generally, pale; *φαίνω*, to appear. From the *pale* yellow colour. *Fr. Monogr. ii. p.* 143. *Hym. Eur. p.* 420. *Icon. t.* 167. *f.* 4. *B. & Br. n.* 933. *C. Hbk. n.* 574. *S. Mycol. Scot. n.* 542. *Hoffm. Ic. t.* 5. *f.* 1.

**51. *H. psittacinus* Fr.—Pileus** about 2.5 cent. (1 in.) broad, for the most part yellow *tinged with green*, slightly fleshy, campanulate then expanded, umbonate, smooth, very *glutinous* when young, slightly viscid when more fully grown, becoming pale but not changing colour and not torn. **Stem** 5 cent. (2 in.) long, 4-6 mm. (2-3 lin.) thick, hollow, equal, but often bent, even, toughish, *greenish*, at least at the apex. **Gills** *adnate*, but very ventricose, rather thick, broad, somewhat distant, most frequently green.

The colour of the pileus and stem is very variable, when full grown for the most part yellowish, but varying brick-colour, reddish, whitish, most frequently with a greenish tinge. In the toughness of the stem it approaches *H. latus*. Easily distinguished by the greenish gluten with which it is smeared, and which persistently tinges the apex of the stem. It varies very much in size.

In pastures. Common. Aug.-Oct.

In some stages of growth it is exceedingly tough. Spores ellipsoid-sphaeroid,  $7-8 \times 5-6$  mk. *K.*;  $6 \times 7$  mk. *W.G.S.* Name—*psittacus*, the ringed green parrot. From the colours. *Fr. Monogr. ii. p.* 143. *Hym. Eur. p.* 420. *Berk. Out. p.* 202. *C. Hbk. n.* 575. *S. Mycol. Scot. n.* 543. *Hussey i. t.* 41. *Ag. Schaff. t.* 301. *Sow. t.* 82. *Grev. t.* 74.—*Bull. t.* 545. *f.* 1. *Ventur. t.* 42. *f.* 1-3. *Batt. t.* 21. *f.* E.

**52. *H. unguinosus* Fr.—Pileus** about 5 cent. (2 in.) broad, smeared with dense fuliginous gluten, slightly fleshy, campanulate then convex, obtuse, even, or at length rimosely incised. **Stem** 5 cent. (2 in.) long, 6 mm. (3 lin.) thick, hollow, commonly attenuated at the base and apex, unequal, somewhat compressed, *glutinous*, of the same colour as the pileus. **Gills** adnate, but very ventricose, distant, thick, broad, connected by veins, *shining white*, becoming glaucous, soft. Hygro-  
phorus.

Inodorous, very fragile, appearing chiefly in later autumn. The fuliginous colour is unchangeable. The tough fuliginous gluten on the pileus distils in drops. It differs sufficiently from *H. irrigatus* in its larger stature, in its watery substance, in the gluten, and in its ventricose separating gills.

In woods and pastures. Frequent. Aug.—Oct.

Name—*unguinus*, oily. From the gluten. *Fr. Monogr.* ii. p. 144. *Hym. Eur.* p. 421. *Icon. t.* 168. *f.* 2. *Berk. Out.* p. 202. *C. Hbk. n.* 577. *S. Mycol. Scot. n.* 544.

**53. *H. nitratus* Fr. *Fuscous-cinereous*.—Pileus** 5 cent. (2 in.) broad, becoming pale, scarcely fleshy, very fragile, convex, obtuse or depressed in the centre, at the first slightly viscid, soon flocculose, then *squamulose* and rimosely incised, irregularly shaped, somewhat repand. **Stem** 5–7.5 cent. (2–3 in.) long, 6–10 mm. (3–5 lin.) thick, hollow, equal, but often twisted, and at length somewhat compressed, fragile, externally polished, smooth, whitish, woven with thick crisped fibres. **Gills** broadly emarginate, very broad, as much as 18 mm. ( $\frac{3}{4}$  in.) broad (obliquely ovate), distant, thick, mucid-soft, connected by veins, whitish then becoming glaucous.

The above is the larger and rarer form. B. The smaller form is the commoner. Pileus 1–2.5 cent. ( $\frac{1}{2}$ –1 in.) broad, viscous when young and moist, even, fuscous, but soon dry and rimoso-squamulose, cinereous. Stem 2.5 cent. (1 in.) and more long, 2–6 mm. (1–3 lin.) thick, unequal, commonly compressed and undulated, sometimes yellowish. Gills at first wholly adnate, even with a decurrent tooth, then ventricose and glaucous, but broad and thick, as if smeared.

Widely removed from all others by its *strong nitrous odour*, and by the colour being by no means bright. As regards substance and structure it agrees well with *H. coccineus*, and affords the best example of the generic character; the gills are thick, juicy, and the hymenium, which is waxy soft as if composed of fat, is as easily rubbed off from the trama as if it were fat. Very changeable in stature according to locality.

In pastures. Uncommon. Aug.—Sept.

Name—*nitrum*, nitre. From its *nitrous odour*. *Fr. Hym. Eur.* p. 421. *S. Mycol. Scot. n.* 545. *H. murinaceus* *Fr. Monogr.* ii. p. 145. *Berk. Out.* p. 203. *C. Hbk. n.* 578.

\* ***H. glauco-nitens* Fr.—Pileus** olivaceous-black or fuliginous,

Hygro-  
phorus.

becoming pale, *streaked with fibrils*. **Stem** equal, shining. **Gills** becoming glaucous.

Stiff. It seems to be a young state of *H. nitratus*, scorched by the sun, but it is reckoned distinct by Persoon and others. Odour the same, but weaker.

In mixed woods. Pass of Killiecrankie. 1877. Aug.

Name—*glaucus*, glaucous; *nitens*, shining. *Fr. Hym. Eur. p. 421. B. & Br. n. 1671. S. Mycol. Scot. n. 546. Ag. Batsch f. 192.*

Lactarius.

GENUS IX.—**Lactarius** (*lac*, milk). *Fr. Epicr. p. 333.*

Hymenophore continuous with the stem. Gills unequal, membranaceous-waxy, slightly rigid, *milky*, acute at the edge. Spores globose, white, rarely yellowish. *Fleshy fungi, growing on the ground (with two exceptions), putrescent, pileus depressed, gills adnato-decurrent and often branched.*



XLIX. *Lactarius subdulcis*.  
One-third natural size.

A very distinct genus and easily recognised. Pileus in some species zoned which is peculiar to this genus. Watery milk is never a normal feature of the genus, but where it does occur it is owing to the dampness of the station where the specimens have grown. *Fr. Hym. Eur. p. 421.*

The nearest allies to the *Lactarii* are the *Russulæ*, which are at once distinguished by their want of milk. The genus includes some species which are edible and delicious, while others are exceedingly poisonous.

The best mark of distinction of species is the nature of the milk.

**Tribe I. PIPERITES** (*piperitis*, pepper-wort). Stem central. Gills unchangeable, naked, not changing colour and not pruinose. Milk at the first white, (commonly) acrid.

\* **TRICHOLOMOIDEI**. *Pileus viscid when moist, margin at first involute, tomentose.*

\*\* **LIMACINI**. *Pileus viscid when moist, pelliculose, margin naked.*

\*\*\* **PIPERATI**. *Pileus without a pellicle, hence absolutely dry, most frequently unpolished.*

**Tribe II. DAPETES** (*daps*, food). Stem central. Gills naked; milk at the first deeply coloured.



*Tribe III. RUSSULARES.* Stem central. Gills pallid then changing colour, Lactarius afterwards darker, glancing when turned to the light, at length white-pruinose. Milk at the first white, mild or from mild turning acrid.

\* *Pileus at the first viscid.*

\*\* *Pileus unpolished, squamulose, villous or pruinose.*

\*\*\* *Pileus polished, smooth.*

*Tribe IV. PLEUROPUS.* Stem excentric or lateral. Growing on trunks.

## TRIBE I.—PIPERITES.

\* TRICHOLOMOIDEI. *Pileus viscid when moist, &c.*

1. **L. scrobiculatus** Fr.—**Pileus** 10 cent. (4 in.) and more broad, yellow, commonly zoneless, fleshy, but not compact, at first convex, umbilicate, at length infundibuliform, obtuse, with agglutinated villous down, very viscid when moist, margin involute, bearded, at length when old unfolded and almost naked; flesh lax, whitish, but commonly becoming yellow when broken. **Stem** 4-6 cent. (1½-2½ in.) long, as much as 2.5 cent. (1 in.) and more thick, stout, often stuffed when young, but soon remarkably hollow, equal, light yellow, pitted with broad roundish spots, smooth, somewhat viscid, pubescent at the base. **Gills** adnato-decurrent, thin, crowded, quaternate, whitish. **Milk** plentiful, at first white, soon sulphur-yellow when exposed to the air.

Very acrid, odour not unpleasant. Pileus sometimes as much as 30 cent. (12 in.) broad. The pileus becomes pale in open places, and is sometimes zoned. In very watery places when broken it occasionally becomes azure-blue-purple, milk white, soon serous.

On the ground. Bristol.

Spores subsphæroid, echinulate, 11-12 × 8-10 mk. *K.*; subglobose, 6-8 mk. *C.B.P.* Name—*scrobis*, a trench. From the pitted stem. *Fr. Monogr.* ii. p. 152. *Hym. Eur.* p. 422. *Grevillea*, vol. xii. p. 42. *Ag. Schæff. t.* 227. *Krombh. t.* 58. f. 1-6. B. milk watery white becoming violet. *Barla t.* 18. f. 3-6.

2. **L. torminosus** Fr.—**Pileus** 7.5 cent. (3 in.) and more broad, pallid flesh-colour, laxly fleshy, somewhat fragile, obtuse, depressed, becoming even, viscid when moist, somewhat zoned, with a white-fibrillose beard at the margin and towards the margin which is at first involute; flesh pallid. **Stem** 6-9 cent. (2½-3½ in.) long, 18 mm.-2.5 cent. (¾-1 in.) thick, stuffed soon hollow, equal or attenuated downwards, delicately adpressedly tomentose or becoming smooth, even or obsoletely pitted, dry. **Gills** adnato-decurrent, 2-3 mm. (1-1½ lin.) broad, very thin and crowded, unequal but rarely connected by branches, paler than the pileus. **Milk** white, unchangeable, acrid.

Inodorous. It frequently varies pale-ochraceous. It also occurs with the

Lactarius.

pileus wholly white and tomentose *Sow. t.* 103. In another form the pileus is smooth, zoneless, white. In a variety the pileus is at length excentric, whitish, cracked in a tessellated manner, margin spreading, naked.

In mixed woods. Common. Aug.—Nov.

Spores subsphæroid, echinulate, 6-8 mk. *K.*; 9×6 mk. *W.G.S.* Name—*tormina*, gripes. Causing gripes. *Fr. Monogr. ii. p.* 153. *Hym. Eur. p.* 422. *Sv. ätl. Sv. t.* 28. *Berk. Out. p.* 203. *C. Hbk. n.* 582. *S. Mycol. Scot. n.* 547. *Ag. Schæff. t.* 12. *Fl. Dan. t.* 1068. *Sv. Bot. t.* 184. *Bull. t.* 529. *f. 2. Krombh. t.* 13. *f.* 15-23. *Ventur. t.* 30. *f. 2. Barla t.* 18. *f.* 7-10. *Harz. t.* 11.

**3. *L. cilicioides* Fr.—Pileus** 5-10 cent. (2-4 in.) broad, *flesh-colour inclining to fuscous*, fleshy, convex then flattened and depressed in the centre, everywhere tomentose, viscous, zoneless, margin involute, fibrilloso-woolly; flesh not very compact, yellowish-white. **Stem** 5-7.5 cent. (2-3 in.) long, 2.5 cent. (1 in.) thick, firm, *stuffed*, somewhat hollow only when old, equal, *even*, pruinato-silky under a lens, *pallid, never pitted or spotted*. **Gills** decurrent, crowded, *branched*, white inclining to yellow. **Milk** acrid, *white or light yellow-white*.

The stem is dingy, not of a pure colour as in *L. torminosus*, &c. Intermediate between *L. torminosus* and *L. turpis*.

In woods. Frequent. Sept.—Nov.

Name—*κιλίκιον*, goat's-hair cloth; *εἶδος*, appearance. *Fr. Monogr. ii. p.* 154. *Hym. Eur. p.* 422. *Berk. Out. p.* 203. *C. Hbk. n.* 583. *S. Mycol. Scot. n.* 548. *Ag. Schæff. t.* 228. *Krombh. t.* 58. *f.* 11-13 var. *B. & Br. n.* 1887.

**4. *L. turpis* Fr.—Pileus** large, as much as 30 cent. (12 in.) broad, *olivaceous inclining to umber*, fleshy, rigid, convex becoming plane, disc-shaped or umbilicate, at length depressed, innato-villous at the circumference or wholly, covered over with tenacious gluten, *zoneless*, sometimes tawny towards the margin, *at length entirely inclining to umber; margin for a long time involute, at the first villous, olivaceous-light-yellow*, then more or less flattened, at length often densely rivuloso-sulcate; flesh compact, white, then slightly rufescent. **Stem** 4-7.5 cent. (1½-3 in.) long, 1-2.5 cent. (½-1 in.) and more thick, *solid*, hard, equal or *attenuated downwards*, even or pitted and uneven, but not spotted, viscid or dry, *pallid or dark olivaceous*, ochraceous-whitish at the apex. **Gills** adnato-decurrent, thin, 2-4 mm. (1-2 lin.) broad, much crowded, forked, *white straw-colour*, spotted fuscous when broken or bruised. **Milk** acrid, white, unchangeable.

Gregarious, *rigidly and compactly fleshy*; habit almost that of *Paxillus involutus*. It varies with the stem hollow, and the pileus somewhat zoned.

In woods, roadsides, &c. Common. Sept.—Nov.

The pileus varies in size from about 7.5-30 cent. (3-12 in.) Spores *Lactarius*. sphaeroid or subsphaeroid, uniguttate, echinulate, 6-8 mk. *K.* Name—*turpis*, base. From its ugly appearance. *Fr. Monogr.* ii. p. 154. *Hym. Eur.* p. 423. *Sv. atl. Sv. t.* 60. *Berk. Out. p.* 203. *C. Hbk. n.* 584. *S. Mycol. Scot. n.* 549. *Ag. Weinm.—Fl. Dan. t.* 1913. *Krombh. t.* 69. *f.* 1-6. *Harz. t.* 60.

**5. *L. controversus* Fr.—Pileus** 7.5 cent. (3 in.) and more broad, fleshy, compact, rigid, at the first convex, broadly umbilicate, when fuller grown *somewhat infundibuliform, oblique*, on emerging from the ground dry, flocculose, *whitish*, then with rain smooth, viscid, *reddish, with blood-coloured spots and zones* (especially towards the margin), margin acute when young, closely involute, more or less villous; *flesh very firm*. **Stem** commonly 2.5 cent. (1 in.) long and thick, sometimes however 5 cent. (2 in.) long and then manifestly attenuated towards the base and often excentric, *solid, obese*, even but pruinose and as if striate at the apex from the obsoletely decurrent tooth of the gills, wholly *white*, never pitted. **Gills** decurrent, thin, very crowded, 2-4 mm. (1-2 lin.) broad, with many shorter ones intermixed, but rarely branched, pallid - white - flesh - colour. **Milk** white, unchangeable, plentiful.

Odour weak but pleasant, taste very acrid. Allied to *L. piperatus*.

In woods. Uncommon. Aug.—Oct.

Spores echinulate, 8×6 mk. *W.G.S.* Name—*contra*, against; *verto*, to turn. From its becoming at length infundibuliform. *Fr. Monogr.* ii. p. 155. *Hym. Eur. p.* 423. *Sv. atl. Sv. t.* 29. *B. & Br. n.* 1282. *C. Hbk. n.* 585. *S. Mycol. Scot. n.* 550. *Trans. Woolh. Club*, 1868, fig. *Ag. Pers.—Vittad. t.* 37. *Ventur. t.* 51. *f.* 1-4. *Barla t.* 18. *f.* 1, 2. *Bull. t.* 538. *Batsch f.* 201. *Krombh. t.* 56. *f.* 5-7 (var.)

**6. *L. pubescens* Fr.—Pileus** about 5 cent. (2 in.) broad, *whitish passing into flesh-colour*, fleshy, *rather plane*, depressed in the centre, *then broadly infundibuliform*, quite *zoneless*, even, dry, becoming smooth at the disc, shining, margin involute, *fibrilloso-pubescent*; *flesh thin but firm*, pliant, white, or flesh-coloured in the cuticle, unchangeable. **Stem** *short*, scarcely 2.5 cent. (1 in.) long, 12 mm. ( $\frac{1}{2}$  lin.) thick at the apex, *attenuated downwards* and sometimes compressed, *stuffed*, soft internally, even (not pitted), commonly pruinose - pubescent when young, but commonly becoming smooth, *flesh-colour then white*. **Gills** adnate or slightly decurrent, crowded, quaternate, narrower than the flesh of the pileus, pallid, slightly flesh-coloured. **Milk** white, unchangeable, not plentiful.

Inodorous, taste very acrid. It is like a poor form of *L. controversus*, but is very distinct.

Lactarius.

Among pebbles. Loch Kenord, Aberdeenshire.

Name—from the *pubescent* margin of the pileus. *Fr. Monogr.* ii. p. 155. *Hym. Eur.* p. 424. *B. & Br. n.* 1282. *S. Mycol. Scot. n.* 551. *Grevillea, t.* 76. f. 2. *Ag. Krombh. t.* 13. f. 1-14.

**Var. *marginetomentosus*** B. & Br.—**Pileus** clothed with fine matted down, margin *tomentose*. **Stem** nearly equal, 3 cent. (1¼ in.) long, about 10 mm. (5 lin.) thick.

Odour pungent. In pastures. Aboyne. Aug. *B. & Br. n.* 1015. *C. Hbk. n.* 586. *S. Mycol. Scot. n.* 551. *Krombh. t.* 13. f. 1, 2.

**\*\* LIMACINI.** *Pileus viscid when moist, pelliculose, &c.*

**7. L. *insulsus*** Fr.—**Pileus** 7.5-10 cent. (3-4 in.) broad, *yellowish-brick-colour*, fleshy, not very compact, deeply umbilicate and at length infundibuliform, rigid, *zoned* (especially round the margin), *smooth*, with a somewhat separable viscid pellicle, margin naked; flesh firm, pallid, somewhat *zoned* under the pellicle. **Stem** *stuffed* when young, but varying hollow when old, most frequently *curt* and attenuated downwards, 4 cent. (1½ in.) long, as much as 2.5 cent. (1 in.) thick, rarely lengthened to 7.5 cent. (3 in.) and then equal, thinner, smooth, even or more rarely pitted-spotted. **Gills** decurrent, very crowded, forked at the base, 2 mm. (1 lin.) or little more broad, whitish, becoming pale. **Milk** *white*, unchangeable, acid.

Its stature and size are those of *L. deliciosus*, but the colours are paler; the milk when the plant is old is often watery (not coloured); otherwise when wounded it does not change colour as *L. deliciosus* does.

In mixed woods. Common. Aug.-Oct.

Spores globose, pallid. *Fr.* Name—*insulsus*, insipid, tasteless. *Fr. Monogr.* ii. p. 157. *Hym. Eur.* p. 424. *Berk. Out. p.* 204. *t.* 13. f. 2. *C. Hbk. n.* 587. *S. Mycol. Scot. n.* 552. *Krombh. t.* 12. f. 1-6. *Hussey, i.* t. 59.

**8. L. *zonarius*** Fr.—**Pileus** 5-10 cent. (2-4 in.) broad, *pallid orange or pale yellowish to deeper yellow*, fleshy, convex becoming plane, then depressed, somewhat umbilicate, margin *thin*, naked, long *involute*, at first even, smooth, with a closely adnate pellicle, viscid, beautifully *zoned* most frequently towards the margin, *at length*—and at the first at the circumference—*minutely ruguloso-flocculose*; flesh *compact*, white. **Stem** 5-7.5 cent. (2-3 in.) long, equal, or 2.5 cent. (1 in.) and attenuated downwards, 1-2.5 cent. (½-1 in.) thick, *firm, solid*, elastic, spongy-soft internally, even, smooth, dry, white then yellowish, unspotted, pale upwards. **Gills** rounded-adnate and adnato-decurrent,

arcuate, thin, narrow, somewhat crowded, branched, scarcely broader than 2 mm. (1 lin.), whitish, at length becoming dingy yellowish, becoming dingy or even somewhat æruginous when bruised. **Milk** white, unchangeable, acrid.

Pileus sometimes deeply umbilicate. Like *L. insulsus*, but well distinguished from it by its compact substance, &c. It varies in having the stem hollow, the walls of the cavity at first white, then ochraceous-tawny, and also in the pileus being flexuous, smooth, zoneless.

In woods. Rare. Aug.—Oct.

Spores minutely echinulate, almost globular, 7 mk. *W.G.S.* Name—*zona*, a zone. Zoned. *Fr. Monogr.* ii. p. 158. *Hym. Eur.* p. 425. *Berk. Out.* p. 204. *C. Hbk. n.* 588. *S. Mycol. Scot. n.* 553. *Fl. Bat. t.* 825 middle fig. *Ag. Bull. t.* 104. *Vaill. Par. t.* 12. f. 7.

**9. L. blennius** Fr.—**Pileus** 5–10 cent. (2–4 in.) broad, *pallid olivaceous or æruginous-grey*, fleshy, plano-depressed, *glutinous*, often *concentrically guttate* or somewhat zoned, the *margin, which is at the first bent inwards*, slightly downy when young, soon naked; flesh rigid, white. **Stem** 4–5 cent. (1½–2 in.) long, 12–18 mm. (½–¾ in.) thick, stuffed then hollow, somewhat equal or attenuated downwards, even, viscid, pallid olive or pallid-grey. **Gills** adnato-decurrent, thin, crowded, about 2 mm. (1 lin.) broad, white, *becoming cinereous when wounded*. **Milk** white, unchangeable.

Firm, acrid. Dry in fine weather. When young it must not be confounded with *L. turpis*. It varies with the disc of the pileus rufescent: *Fl. Dan. t.* 1961. f. 1.

In woods. Common. Aug.—Nov.

Spores sparingly echinulate, 6×8 mk. *W.G.S.* Name—*βλεννός*, mucous matter. Slimy. *Fr. Monogr.* ii. p. 159. *Hym. Eur.* p. 425. *Berk. Out.* p. 204. *C. Hbk. n.* 589. *S. Mycol. Scot. n.* 554. *Krombh. t.* 69. f. 7–9. *Krapf. t.* 4. f. 11–13. *Sterb. t.* 5. E. B. disc rufescent. *Fl. Dan. t.* 1691. f. 2.

**10. L. hyginus** Fr.—**Pileus** 6–10 cent. (2½–4 in.) broad, *red-flesh-colour* or reddish brown, fleshy, not thick, umbilicate, becoming plane, *the thin margin inflexed*, even, viscous, zoneless or somewhat zoned, often shining; flesh white, somewhat rigid. **Stem** 5–10 cent. (2–4 in.) long, 18 mm.–2.5 cent. (¾–1 in.) thick, stuffed then hollow, often inflated, constricted at the apex, attenuated at the base, smooth, here and there *pitted or somewhat spotted*. **Gills** adnato-decurrent, thin, crowded, 2–4 mm. (1–2 lin.) broad, *white then light yellow-ochraceous*. **Milk** white, unchangeable.

Taste acrid; *rigid-fragile*, pileus somewhat repand. The gills are not pruinose as in the Russulares. It varies with the stem curt, firm, even, the

**Lactarius.** pileus at length darker in the centre, becoming pale, tan-flesh-colour, round the margin.

In woods. Rare. Aug.-Oct.

Spores globose, echinulate, 5-6 mk. *C.B.P.* Name—*βουγιών*, a vegetable dye, crimson or scarlet. From the colour. *Fr. Monogr.* ii. p. 160. *Hym. Eur.* p. 426. *Icon.* t. 169. f. 2. *Berk. Out.* p. 204. *C. Hbk.* n. 590. *S. Mycol. Scot.* n. 555. *Ag. Krombh.* t. 14. f. 15, 16.

**11. L. trivialis** Fr.—**Pileus** 10-17.5 cent. (4-7 in.) broad, at first dark lurid, becoming pale when full grown, pallid yellowish, tan-flesh-colour, zoneless, convex then soon depressed, at length infundibuliform, margin at first involute, then flattened, at length the pellicle of the margin only inflexed, even, smooth, viscous; flesh equal, somewhat rigid, white. **Stem** stout, 2.5-15 cent. (1-6 in.) long, 2.5 cent. (1 in.) and more thick, commonly inflated-hollow, even, smooth, slippery, unspotted, almost paler than the pileus. **Gills** somewhat decurrent, tense and straight, rather broad, 4-8 mm. (2-4 lin.), somewhat thin, crowded, whitish, becoming pale. **Milk** white, unchangeable.

Acrid, large, fleshy, rigid-fragile. The stem is shorter and firmer in drier situations.

In woods, chiefly pine. Uncommon. Aug.-Oct.

Name—*trivialis*, common. Occurring everywhere in Northern Europe. *Fr. Monogr.* ii. p. 160. *Hym. Eur.* p. 426. *B. & Br.* n. 934. *C. Hbk.* n. 591. *S. Mycol. Scot.* n. 556. *Ag. Krombh.* t. 14. f. 17, 18.

**12. L. circellatus** Fr.—**Pileus** 5-7.5 cent. (2-3 in.) broad, rufous inclining to fuscous in wet weather, but becoming pale, variegated with darker zones, fleshy, umbilicato-convex then becoming plane, depressed in the centre, for the most part repand when older, even, smooth, very viscous in wet weather; flesh grumous-compact, white. **Stem** 4-5 cent. (1½-2 in.) long, 12 mm. (½ in.) and more thick, solid, equal or attenuated downwards and often ascending, even, smooth, pale. **Gills** horizontal (not arcuate), somewhat decurrent with a tooth, very thin and crowded, narrow, often forked, whitish, at length becoming yellow. **Milk** white, unchangeable.

Stature curt and firm; odour not unpleasant; taste acrid, stinging.

In woods. Rare.

Spores spherical, echinulate, 10 mk. *Q.* Name—*circellus*, a small ring. Ringed. From the zones. *Fr. Monogr.* ii. p. 161. *Hym. Eur.* p. 426. *Berk. Out.* p. 204. *C. Hbk.* n. 592. *Ag. Batt.* t. 13. *D. Sow.* t. 203. *Ventur.* t. 34. f. 4, 5.

**13. L. uvidus** Fr.—**Pileus** 5-10 cent. (2-4 in.) broad, hoary-

*whitish, grey-flesh-colour-livid, becoming fuscous, fleshy, somewhat thin, convex, plane, then depressed, margin at first involute, very soon almost straight, even, smooth, viscous, zoneless; flesh rather lax, white then lilac when broken. Stem* 4-9 cent. ( $1\frac{1}{2}$ - $3\frac{1}{2}$  in.) long, 12-18 mm. ( $\frac{1}{2}$ - $\frac{3}{4}$  in.) thick, soon hollow, even or somewhat pitted, smooth, viscous, whitish becoming light yellow, sometimes with yellowish-ferruginous spots. **Gills** adnate and somewhat decurrent, somewhat arcuate, 2-4 mm. (1-2 lin.) and more broad, thin, crowded, very *unequal*, here and there connected by branches and anastomosing with veins, *shining white to dead white*, soon spotted with lilac when wounded, at length here and there pallid ochraceous-ferruginous, even tinged with pallid brick-colour. **Milk** *white then lilac.*

*Soft-fragile; somewhat insipid, at length slowly acrid; odour weak, somewhat nauseous-pungent, somewhat aromatic.* It varies with the pileus obsoletely zoned, slightly pellucid-striate when old, and with the gills sometimes yellowish. The milk rarely remains white or at length turns dingy tan-colour; commonly it becomes lilac.

In woods. Uncommon. Aug.-Sept.

Spores globose, echinulate, 5-6 mk. C.B.P. Name—*uvidus*, moist. *Fr. Monogr.* ii. p. 162. *Hym. Eur.* p. 426. *Berk. Out.* p. 205. *C. Hbk. n.* 593. *S. Mycol. Scot. n.* 557.

\*\*\* PIPERATI. *Pileus without a pellicle, &c.*

**14. L. flexuosus** Fr.—**Pileus** 5-10 cent. (2-4 in.) broad, *lead-grey or violet-grey*, becoming pale, zoned or zoneless, fleshy, convex, becoming plane, depressed, somewhat repand, margin at first and for a long time bent inwards, at length spreading, erect, *dry*, at the first smooth, somewhat shining, then *rivuloso-scaly, unpolished*, opaque; flesh hard, grumoso-compact, white. **Stem** 5-9 cent. ( $2-3\frac{1}{2}$  in.) long, 2.5 cent. (1 in.) and more thick upwards, *solid*, obese or equally attenuated downwards, not rarely excentric, here and there somewhat lacunose, not however pitted, pallid grey, the apex whitish, the base somewhat yellowish. **Gills** adnate, somewhat horizontal, *thick, distant*, 2-3 mm. ( $1-1\frac{1}{2}$  lin.) broad, connected by branches, *light yellowish*, at length becoming whitish-flesh-colour. **Milk** white, unchangeable.

\*\*\* Remarkable; *hard, firm*, very acrid. Under a lens the stem is clothed with very delicate adpressed down. There is a form *roseo-zonatus* Post, approaching *L. flexuosus* almost as *Gomphidius roseus* approaches *Gomphidius glutinosus*. *Fr. Icon. t.* 169. *f.* 3.

In pine woods, &c. Haywood Forest. Rothiemurchus. Glamis. Oct.

Lactarius.

Edible according to some, but doubtful from its very acrid taste. Spores sphaeroid or subsphaeroid, uniguttate, 6-8 mk. *K.* Name—*flexus*, a bending. Flexuous. From the margin of the pileus. *Fr. Monogr.* ii. p. 163. *Hym. Eur.* p. 427. *Grevillea*, vol. x. p. 45. *S. Mycol. Scot. Supp. Scot. Nat.* 1885, p. 73. *Ag. Schæff. t.* 235. *Harz. t.* 43.

**15. *L. pyrogalus* Fr.—Pileus** 5-7.5 cent. (2-3 in.) broad, *cinereous-grey*, at length becoming dingy yellow, firm, fleshy, convex becoming plane, margin very soon spreading, depressed, under a lens delicately grumoso-uneven, smooth, in rainy weather moist but not viscid, somewhat zoned; (flesh somewhat thin, but grumoso-firm, white). **Stem** 4 cent. (1½ in.) and more long, 6-10 mm. (3-5 lin.) thick, *stuffed then soon hollow*, often attenuated downwards, even or here and there delicately grumoso-tubercular under a lens, smooth, pallid-white, somewhat dingy. **Gills** adnato-decurrent, *thin*, somewhat distant, 2 mm. (1 lin.) and more broad, *light yellow-wax-colour*, the colour rich, inclining to ochraceous. **Milk** plentiful, white, unchangeable, *very acrid*.

Pileus dry in fine weather. Allied to *L. flexuosus*, but *smaller and thinner*. There is a var. like *L. fuliginosus* with the pileus zoneless and tan-colour, not fuliginous, and the gills yellow and distant.

In woods and pastures. Frequent. Aug.—Oct.

Poisonous. Spores sphaeroid or subsphaeroid, echinulate, 6-10 mk. *K.*; sparingly echinulate, white or with a suggestion of ochre, almost globular, 6 mk. *W.G.S.* Name—*πῦρ*, fire; *γάλα*, milk. From the milk being extremely acrid. *Fr. Monogr.* ii. p. 163. *Hym. Eur.* p. 427. *Berk. Out.* p. 205. *C. Hbk. n.* 594. *S. Mycol. Scot. n.* 558. *Ag. Bull. t.* 529. f. 1. *Krombh. t.* 14. f. 1-9.

**16. *L. squalidus* Fr.—Pileus** pallid lurid, compact, convexo-plane, umbilicate, dry, smooth, zoneless; flesh white, sprinkled at the margin with saffron-yellow dots. **Stem** solid, equal, smooth, pallid brown. **Gills** adnate, *narrow, yellowish*. **Milk** whitish, *sweet*.

Its affinity is not clear. Krombholz compares it with *L. fuliginosus*, but its stature is that of *L. pyrogalus*, &c., and the pileus is smooth.

In moist places. Scotland, 1875; locality not recorded.

Name—*squalidus*, dirty. From the dingy colour. *Fr. Hym. Eur.* p. 428. *B. & Br. n.* 1556. *S. Mycol. Scot. n.* 559. *Ag. Krombh. t.* 40. f. 23-25.

**17. *L. capsicum* Schulz.—Pileus** 7.5 cent. (3 in.) broad, chestnut, paler at the disc, compact, pulvinate, dry, margin closely involute; flesh yellowish, becoming fuscous on exposure to the air. **Stem** about 2.5 cent. (1 in.) thick, solid, firm, whitish, *striate with tawny or rufescent fibres*. **Gills** adnato-decurrent, somewhat crowded, tawny, somewhat orange. **Milk** white, acrid.



Very robust. The gills are described as anastomosing.

Lactarius.

In wood. Dumfries, 1884. Sept.

Name—probably from the taste being hot like that of *Capsicum*. Schulz.—*Fr. Hym. Eur. p. 428. S. Mycol. Scot. Supp. Scot. Nat. 1885, p. 73.*

**18. L. chrysorheus** Fr.—**Pileus** 5–7.5 cent. (2–3 in.) broad, somewhat flesh-colour, with darker zones or spots, fleshy, firm, umbilicate then infundibuliform; flesh white, bright sulphur-yellow when broken. **Stem** 5–7.5 cent. (2–3 in.) long, 1–2.5 cent. ( $\frac{1}{2}$ –1 in.) thick, rigid, stuffed then soon hollow, equal, even, delicately pruinose under a lens, dry, *white*, changing colour only when old. **Gills** decurrent, very thin and crowded, narrow, 4 mm. (2 lin.) broad, pallid-yellowish, not truly changing colour and not pruinose. **Milk** *white then bright sulphur-yellow (golden)*, very acrid.

It is easily distinguished from all neighbouring species by the change of colour in the flesh and milk. Pileus when dry sprinkled with thin, whitish lustre, at length darker.

In mixed woods. Frequent. Sept.–Oct.

Margin of pileus wavy, involute, and minutely downy when young; flesh firm, crisp; gills slightly forked; stem downy at the base. *M. J. B.* Name—*χρῖσος*, gold; *ρέω*, to flow. From the yellow milk. *Fr. Monogr. ii. p. 165. Hym. Eur. p. 428. Berk. Out. p. 206. C. Hbk. n. 597. S. Mycol. Scot. n. 560. Ag. Bolt. t. 144. Krombh. t. 12. f. 7–14.*

**19. L. acris** Fr.—**Pileus** 7.5 cent. (3 in.) broad, cinereous-fuliginous, sometimes darker sometimes paler, fleshy, firm, rigid, irregular, often excentric or emarginate on one side, at first convex then plane, at length obliquely infundibuliform, moist rather than viscous, scarcely zoned, here and there spotted; flesh white. **Stem** about 4–5 cent. ( $1\frac{1}{2}$ –2 in.) long, 12 mm. ( $\frac{1}{2}$  in.) or a little more thick, stuffed then hollow, *attenuated downwards*, often *oblique*, ascending or curved, *pallid*, at length fragile. **Gills** somewhat decurrent, thin, somewhat crowded, somewhat divided behind, at first pallid, then yellow-flesh-colour. **Milk** *white, soon reddish*, plentiful.

Strong-smelling, acrid. Very different from *L. fuliginosus*, which is of the same colour, but approaching *L. pyrogalus*.

In woods. Rare. Aug.–Nov.

Spores echinulate, yellow,  $6 \times 8$  mk. *W.G.S.* Name—*acer*, sharp. From the acrid taste. *Fr. Monogr. ii. p. 164. Hym. Eur. p. 428. Berk. Out. p. 207. C. Hbk. n. 596. S. Mycol. Scot. n. 561. Ag. Bolt. t. 60. Batsch f. 68. Batt. t. 13. E.*

**20. L. pergamenus** Fr. *White*.—**Pileus** fleshy, pliant, convex

**Lactarius.** then plano-depressed, repand, zoneless, slightly wrinkled, smooth. **Stem** stuffed, smooth, changing colour. **Gills** *adnate*, very narrow, *horizontal*, very crowded, branched, white then straw-colour. **Milk** white, acrid.

Very much allied to *L. piperatus*, but differing in the *stem* being stuffed, at length softer internally, elongated (7.5 cent., 3 in.), unequal, attenuated downwards and here and there ascending, *quite smooth*; in the *pileus* being *thinner*, *pliant*, elastic, most frequently irregular and excentric, for the most part flexuous, at first convex (not umbilicate), then *rather plane*, the surface *very smooth*, but unpolished and *wrinkled* in a peculiar manner; and in the *gills* being *adnate*, not decurrent, *very crowded*, *very narrow* (scarcely 2 mm., 1 lin. broad), always *straight and horizontal*, not arcuate or extended upwards, *soon straw-colour*. The flesh is very milky, but the gills are sparingly so.

In woods. Haywood Forest, &c. Oct.

Spores subglobose, rather irregular, 6-8 mk. *C.B.P.* Name—*pergamenus*, *pergamena*, parchment. From its toughness and parchment-like appearance. *Fr. Monogr.* ii. p. 166. *Hym. Eur.* p. 430. *Grevillea*, vol. xi. p. 71. *Ag. Swartz.*—*Krombh.* t. 57. f. 1-3. *Batsch* f. 59.

**21. L. piperatus** Fr.—**Pileus** 10-22.5 cent. (4-9 in.) broad, *white*, fleshy, rigid, umbilicate when young, reflexed (margin at first involute) at the circumference, when full grown wholly *infundibuliform*, for the most part regular, even, smooth, zoneless; flesh white. **Stem** 2.5-5 cent. (1-2 in.) long, 2.5-5 cent. (1-2 in.) thick, solid, obese, equal or obconical, even, obsoletely pruinose, white. **Gills** *decurrent*, *crowded*, *narrow*, scarcely broader than 2 mm. (1 lin.), obtuse at the edge, *dichotomous*, arcuate then all *extended upwards* in a straight line, white, here and there with yellow spots. **Milk** white, unchangeable, plentiful and very acrid.

*Compact, firm, dry*, inodorous. The pileus becomes obsoletely yellow when old. Although the gills are spotted with yellow, they do not change to straw-colour like those of *L. pergamenus*.

In mixed wood. Common. Aug.-Oct.

When bruised changing to umber. *M. F. B.* Said to be edible, but very acrid. Spores not echinulate, generally with an apiculus, 5 × 6 mk. *W. G. S.* Name—*piper*, pepper. From the acrid taste. *Fr. Monogr.* ii. p. 166. *Hym. Eur.* p. 430. *Sv. atl. Sv. t.* 27. *Berk. Out.* p. 205. *C. Hbk. n.* 598. *S. Mycol. Scot. n.* 563. *Ag. Scop.*—*Fl. Dan. t.* 1132. *Barla t.* 22. f. 1-5. *Harz. t.* 39. *Krombh. t.* 56. f. 1-4. *Bull. t.* 200. *Paul. t.* 68. f. 3-4.

**22. L. vellereus** Fr.—**Pileus** 12.5-17.5 cent. (5-7 in.) broad, white, fleshy, compact, *convexo-saucer-shaped*, the margin for a long time sloping downwards, *innato-pubescent*, dry, zoneless. **Stem** 5-7.5 cent. (2-3 in.) long, 2.5-4 cent. (1-1½ in.) thick, *stout*, solid, *equal*, covered over with innate, very thin pubescence. **Gills** *arcuate*, *adnato-decurrent*, rather thick, acute at the edge, *some-*

*what distant, rather broad* (4 mm., 2 lin. and more), *connected by* Lactarius. *branches* (but not dichotomous), pallid, watery white. **Milk** white, scanty, very bitter.

*Large, hard, rigid.* The pileus tends to become pallid reddish-tan. Very much allied to *L. piperatus*. On being broken it sometimes changes to pallid sulphur-yellow, but after a time becomes white again. Easily confounded with *Russula delicata*.

In woods. Common. Sept.-Dec.

Spores hardly echinulate,  $4 \times 8$  mk. *W.G.S.* Name—*vellus*, fleece. From the pubescence. *Fr. Monogr.* ii. p. 167. *Hym. Eur.* p. 430. *Berk. Out.* p. 206. *C. Hbk. n.* 599. *S. Mycol. Scot. n.* 565. *Hussey i. t.* 63. *Ag. Klotzsch Fl. Bor. t.* 469. *Barla t.* 22. f. 6-8. *Krombh. t.* 57. f. 10-13. *Sow. t.* 104.

**23. L. exsuccus** Smith.—**Pileus** clothed with adpressed down, fleshy, depressed, with an involute margin. **Stem** very short, white, clothed with adpressed down. **Gills** decurrent, connected by veins, and forked, white, shaded with verdigris.

Whole plant rigid and brittle, *milkless*.

In woods. Uncommon.

Considered by some to be a var. of *L. vellereus*. It must be regarded as a distinct species, but I retain it among *Lactarii* only in deference to the opinion of others. Spores covered with spines, almost globular, 9 mk. *W.G.S.* Name—*ex, succus*, juice; juiceless. *Sm. Journ. Bot.* 1873, p. 336. *S. Mycol. Scot. n.* 566. *L. vellereus* var. *exsuccus*. *Fr. Hym. Eur.* p. 430. *Berk. Out.* p. 206. *C. Hbk. n.* 599. *Ag. Berk. Eng. Fl. v. p.* 31.

**24. L. scoticus** B. & Br.—**Pileus** depressed-tomentose then smooth, the involute margin tomentose; flesh firm. **Stem** somewhat unequal, smooth, somewhat flesh-colour. **Gills** thin, scarcely branched. **Milk** persistently white, acrid.

Odour pungent.

Among moss. Aboyne. 1862.

Name—*Scottish*. *B. & Br. n.* 1783. *S. Mycol. Scot. n.* 564.

## TRIBE II.—DAPETES.

*Milk at first deeply coloured, &c.*

**25. L. deliciosus** Fr.—**Pileus** 5-15 cent. (2-6 in.) broad, *orange-brick-colour*, becoming pale, fleshy, when quite young *depressed in the centre*, margin naked, involute, then plano-depressed or broadly infundibuliform with the margin unfolded, smooth, slightly viscid, *zoned* (zones sometimes obsolete); flesh soft, not compact, pallid, coloured at the circumference only by the juice. **Stem** 2.5-5 cent. (1-2 in.) and more long, about 2.5

Lactarius.

cent. (1 in.) thick, stuffed then hollow, at length fragile, equal or attenuated at the base, spotted in a pitted manner, of the same colour as the pileus or paler. **Gills** somewhat decurrent, crowded, narrow, arcuate, often branched, typically *saffron-yellow*, but *becoming pale and always becoming green when wounded*. **Milk** aromatic, from the first red brick-saffron.

The pileus becomes pale in various degrees, often greenish; it occurs also cinereous, fuscous-zoned, &c.

In woods, chiefly pine. Common in some localities. July–Nov.

Edible; when young and fresh one of the most delicious of fungi, and from its orange-coloured milk it cannot be mistaken for any other. Spores white. *Fr.*; sphaeroid, echinulate, 7–8 mk. *K.*; 6 mk. *W.G.S.* Name—from its delicious flavour. *Fr. Monogr.* ii. p. 168. *Hym. Eur.* p. 431. *Sv. atl. Sv. t. 6.* *Berk. Out. p. 206.* *C. Hbk. n. 600.* *S. Mycol. Scot. n. 567.* *Husseyi. t. 67.* *Ag. Schæff. t. 11.* *Fl. Dan. t. 1751.* *Sv. Bot. t. 173.* *Letell. t. 632.* *Viv. t. 13.* *Barla t. 19.* *Vittad. t. 42.* *Krombh. t. 11.* *Harz. t. 10.*

### TRIBE III.—RUSSULARES.

#### \* *Pileus at the first viscous.*

**26. L. pallidus** Fr.—**Pileus** 7.5–15 cent. (3–6 in.) broad, flesh-colour or clay-colour to *pallid*, somewhat *tan*, fleshy, umbilicato-convex, depressed, obtuse, margin broadly and for a long time involute, smooth, viscous, *zoneless*; flesh pallid. **Stem** 5 cent. (2 in.) and more long, about 18 mm. ( $\frac{3}{4}$  in.) thick, somewhat equal, stuffed then *hollow*, even, smooth, of the same colour as the pileus. **Gills** somewhat decurrent, arcuate, rather broad (3–4 mm.,  $1\frac{1}{2}$ –2 lin. and more), somewhat thin, crowded, somewhat branched, whitish at length of the same colour as the pileus. **Milk** white, unchangeable.

Taste somewhat mild. Stature that of *L. deliciosus*, but more lax in texture and always *pallid*. There is a var. with the pileus inclining to fuscous.

In mixed woods. Uncommon. Sept.–Oct.

Spores echinulate, almost round, 8 mk. *W.G.S.* Name—from the *pallid* colour. *Fr. Monogr.* ii. p. 169. *Hym. Eur.* p. 431. *Sv. atl. Sv. t. 61.* *Berk. Out. p. 207.* *C. Hbk. n. 601.* *S. Mycol. Scot. n. 568.* *Ag. Pers.—Krombh. t. 56. f. 10–12?*

**27. L. quietus** Fr.—**Pileus** 7.5 cent. (3 in.) broad, fleshy, depressed, obtuse, margin deflexed, smooth, at the first viscid, somewhat *cinnamon*, flesh-colour, disc darker, somewhat *zoned*, soon dry, somewhat *silky*, opaque, *becoming pale*; flesh white then rufescent. **Stem** 5–7.5 cent. (2–3 in.) long, 12 mm. ( $\frac{1}{2}$  in.) and more thick, stuffed, *spongy*, smooth, rufescent, at length

beautifully *rubiginous*. **Gills** adnato-decurrent, somewhat forked at the base, 3-4 mm. ( $1\frac{1}{2}$ -2 lin.) broad, *white then soon brick-rufescent*. **Milk** white, unchangeable, *sweet*. Lactarius.

Abounding in milk. *Soft*, not so obese and fleshy as *L. pallidus*. The pileus is sometimes of one colour, rufescent.

In woods. Common. Aug.-Nov.

Margin of pileus delicately downy. Stem bearing strong pressure without breaking, when old less firm. Odour oily, and somewhat like that of bugs. *M. J. B.* Spores papillose,  $8 \times 5$  mk. *W. G. S.*; spherical, echinulate, 11 mk. *Q.* Name—*quietus*, mild. From the taste. *Fr. Monogr.* ii. p. 169. *Hym. Eur.* p. 431. *Berk. Out.* p. 207. *C. Hbk.* n. 602. *S. Mycol. Scot.* n. 569. *L. pallidus Saund. & Sm.* t. 16. *Ag. Krombh.* t. 40. f. 5-7.

**28. L. theiogalus Fr.**—Pileus 4-6 cent. ( $1\frac{1}{2}$ -2½ in.) broad, *rufous-tawny*, fleshy, thin, convex when young, then depressed, at length infundibuliform in the centre, margin more or less thin, rather broadly deflected, even, smooth, *viscous*, shining when dry, *zoneless*; flesh whitish. **Stem** 2.5-5 cent. (1-2 in.) long, 4-8 mm. (2-4 lin.) thick, stuffed then hollow, equal, even, *of the same colour as the pileus*. **Gills** adnato-decurrent, about 2 mm. (1 lin.) broad, thin, crowded, pallid then rufescent. **Milk** *white then sulphur-yellow*.

*Slowly acrid*, inodorous. Var. B. pileus zoneless, dry, gills becoming instantly yellow. *Bolt.* t. 9.

In mixed woods. Frequent. Sept.-Oct.

Name—*θειον*, brimstone; *γαλα*, milk. From the sulphur-yellow milk. *Fr. Monogr.* ii. p. 170. *Hym. Eur.* p. 432. *Berk. Out.* p. 206. *C. Hbk.* n. 603. *S. Mycol. Scot.* n. 570. *Ag. Bull.* t. 567. f. 2. *Krombh.* t. 1. f. 23, 24. *Barla* t. 20. f. 14-16. *Paul.* t. 71.

**29. L. vietus Fr.**—Pileus 4-6 cent. ( $1\frac{1}{2}$ -2½ in.) broad, *flesh-colour or livid-grey, becoming pale*, somewhat papillate becoming plane, then *umbilicate*, at length *somewhat infundibuliform*, margin somewhat deflexed, even, smooth, at the first viscid, opaque, *slightly silky when dry*; flesh whitish. **Stem** 4-6 mm. (2-3 lin.) thick, attenuated upwards or somewhat equal, stuffed then hollow, even, smooth, dry, somewhat of the same colour as the pileus. **Gills** adnato-decurrent, thin, somewhat crowded, somewhat flaccid, scarcely broader than 2 mm. (1 lin.), *whitish*. **Milk** *white then somewhat grey*.

*Small, thin, soft, fragile*, changeable in form. *Slowly acrid*, odour weak, somewhat pungent. The gills become at length yellowish.

In grassy woods. Stoke Poges. Glamis. Oct.

Name—*vietus*, shrunk, withered. From its appearance. *Fr. Monogr.*

Lactarius.

ii. p. 171. *Hym. Eur.* p. 432. *Icon. t.* 170. *B. & Br. n.* 1672. *S. Mycol. Scot. Supp. Scot. Nat.* 1882, p. 217.

**30. L. cyathula** Fr.—**Pileus** 2.5–5 cent. (1–2 in.) broad, slightly fleshy, convexo-plane, *umbonate*, at length *plano-depressed*, umbo often vanishing, viscid in wet weather, soon dry, always very *opaque*, when in full vigour even, rufescent-brick or flesh colour, *somewhat zoned*, when dry becoming *pale*, livid or flesh-clay-colour, hoary-tan, *rimoso-rivulose*; flesh white-flesh-colour. **Stem** about 5 cent. (2 in.) long, 2–10 mm. (1–5 lin.) thick, stuffed, internally spongy-soft, equal, round, even, smooth, pale, at length whitish. **Gills** decurrent, very *crowded*, thin, scarcely reaching 2 mm. (1 lin.) in breadth, linear, white-flesh-colour then yellowish. **Milk** white, *unchangeable*, acrid.

Growing in troops, *for the most part very thin and very small*, but larger, firmer, and darker specimens occur. Allied to *L. vietus*, but remarkable for its stature, thinness, and mode of growth; easily distinguished by the *white milk being unchangeable when the gills are wounded*, by its having at the first scarcely any odour, but a *strong smell of bugs when withering or half-dried*. It varies with the pileus zoneless, not umbonate.

In woods. Rare. Aboyne, &c. Aug.–Nov.

Name—*cyathula*, a little cup. From its shape. *Fr. Monogr.* ii. p. 172. *Hym. Eur.* p. 433. *B. & Br. n.* 1016. *C. Hbk. n.* 604. *S. Mycol. Scot. n.* 571.

**\*\* Pileus unpolished, squamulose, &c.**

**31. L. rufus** Fr.—**Pileus** 7.5–10 cent. (3–4 in.) broad, *bay-brown-rufous*, fleshy, *umbonate when young*, soon depressed with an umbo, and at length *infundibuliform*, wholly *zoneless*, dry, at the first *flocculoso-silky*, but soon *polished*, smooth, somewhat *shining*, margin *involute when young*, somewhat *whitish-tomentose*; flesh not compact, pallid. **Stem** 5–7.5 cent. (2–3 in.) long, about 12 mm. ( $\frac{1}{2}$  in.) thick, stuffed, somewhat fragile, equal, *obsoletely pruinose or wholly smooth*, rufescent, but paler than the pileus, white-pubescent at the base. **Gills** adnato-decurrent, crowded, about 3 mm. ( $1\frac{1}{2}$  lin.) broad, quaternate, scarcely branched, ochraceous or pallid then *rufescent*. **Milk** white, unchangeable.

Odour none. Taste *very acrid, stinging*. The colour of the pileus when older or dried passes into cinnamon-rufous. There is a smaller var. which must be carefully distinguished from *L. subdulcis*; very bitter.

In pine woods. Common. June–Nov.

The umbo is remarkably persistent and distinct even when most deeply *infundibuliform*. Very poisonous. Spores globose, white. *Fr.*; subsphaeroid, 8 mk. *K.*; scarcely echinulate, almost round, 5 mk. *W.G.S.* Name—*rufus*,

red. *Fr. Monogr.* ii. p. 173. *Hym. Eur.* p. 433. *Sv. ätl. Sv. t.* 11. *Berk. Lactarius.* *Out.* p. 208. *C. Hbk. n.* 605. *S. Mycol. Scot. n.* 572. *Hussey i. t.* 15. *Ag. Scop.—Krombh. t.* 39. f. 12-15.—*Paul. t.* 22 bis.

**32. L. helvus** Fr.—**Pileus** 5-10 cent. (2-4 in.) broad, *pale yellowish-brick-colour, becoming pale*, fleshy, *fragile*, at first convex, then *flattened, somewhat umbonate, the surface wholly broken up into granuloso-squamulose flocci*; flesh soft, of the same colour as the pileus but paler. **Stem** 5-7.5 cent. (2-3 in.) long, 12 mm. ( $\frac{1}{2}$  in.) and more thick, stuffed, or stuffed then hollow, moderately firm but soft, equal, *brick-colour inclining to pale*, everywhere slightly *pruinat*e, and somewhat tomentose at the base. **Gills** decurrent, crowded, in groups of 8, often dichotomous, 2-3 mm. (1-1 $\frac{1}{2}$  lin.) broad, fragile, at first whitish, then somewhat flesh-colour, at length yellowish. **Milk** scanty, somewhat acrid, white.

Odour weak, sweetish; taste commonly mild. *It occurs most frequently in marshes with watery juice, not coloured.* Very much allied to *L. rufus*.

In fir wood. Grantown, Morayshire. Aug.

Spores sphaeroid, echinulate, 6-7 mk. *K.* Name—*helveus*, pale yellowish; dun-colour. *Fr. Monogr.* ii. p. 174. *Hym. Eur.* p. 433. *S. Mycol. Scot. Supp. Scot. Nat.* 1881, p. 36. *Ag. Krombh. t.* 40. f. 17, 18.

**33. L. mammosus** Fr.—**Pileus** *grey-fuscous*, truly fleshy, at the first convex, *acutely umbonate*, depressed on the umbo at length *vanishing*, soft, everywhere *clothed with depressed down*, dry, zoneless, the margin involute, white-pubescent. **Stem** moderately short and thick, stuffed then hollow, firm, pubescent, *white inclining to pale*. **Gills** adnate, crowded, whitish, then pallid ferruginous. **Milk** white, slowly acrid.

Small, firm, *inodorous*.

In woods. Holme Lacy, 1879.

Name—*mamma*, a breast. Breast-shaped. *Fr. Monogr.* ii. p. 174. *Hym. Eur.* p. 434. *Icon. t.* 170. f. 2. *Grevillea*, vol. viii. p. 98. *Quélet t.* 11. f. 6.

**34. L. glyciosmus** Fr.—**Pileus** 2.5-5 cent. (1-2 in.) broad, grey, brick-colour, fuscous, most frequently passing into violet, zoneless, convex becoming rather plane, acutely papillate, then depressed, the papilla vanishing, *delicately innato-squamulose or unpolished*, opaque; flesh white, acquiring at last a slight tinge of the external colour. **Stem** 2.5-5 cent. (1-2 in.) long, 4-8 mm. (2-4 lin.) thick, stuffed, somewhat equal, even, *pubescent, light yellowish, becoming tawny* when bruised, sometimes somewhat silvery-whitish, especially at the apex. **Gills** arcuate becoming

**Lactarius.** rather plane, adnato-decurrent, 1-2 mm. ( $\frac{1}{2}$ -1 lin.) broad, thin, often connected by branches, *crowded*, straw-colour, becoming in the end tawny-flesh-colour or somewhat ochraceous. **Milk** white, somewhat mild, at length acrid.

*Strong-smelling, with a sweet-spirituos pungent odour*, which distinguishes it from all others; rigid, fragile, variable in stature, resembling *L. vietus*, but sufficiently distinguished from it by the *pileus being absolutely dry, and by the gills being very crowded, straw-colour-ochraceous, and not spotted when wounded*. The milk is sometimes unchangeable, sometimes becoming bright straw-colour-green when dried on the broken flesh, but it does not change colour on the gills. Var. stem elongated, hollow; pileus at first silky, even; also umbilicate, repand, zoned.

In woods, chiefly pine. Uncommon. Sept.-Nov.

Spores sphaeroid, echinulate, 6-8 mk. *K.* Name—*γλύκυσ*, sweet; *δσμή*, smell. Sweet-scented. *Fr. Monogr.* ii. p. 175. *Hym. Eur.* p. 434. *Icon. t.* 170. f. 3. *Berk. Out.* p. 209. *C. Hbk.* n. 606. *S. Mycol. Scot.* n. 573. *Ag. Krombh. t.* 39. f. 16-18?

**35. L. fuliginosus Fr.**—**Pileus** 2.5-10 cent. (1-4 in.) broad, fleshy, spongy-compact, at length softer, becoming plane, *somewhat repand*, not rarely uneven, margin at first inflexed, but very soon spreading, flexuous, somewhat erect; depressed, *very dry*, at first even, *tan-whitish, zoneless, sprinkled with innate, fuliginous pruina*, somewhat velvety to the touch, rather hard, somewhat slippery when wetted, *at length naked*, pelliculose, now and then somewhat rugulose, *soapy-livid*, fawn-brick-colour, the disc sometimes at last brown; flesh almost equally attenuated towards the margin of the pileus, moderately thick in the centre. **Stem** 4-7.5 cent. ( $1\frac{1}{2}$ -3 in.) long, 6-10 mm. (3-5 lin.) thick, spongy-stuffed, somewhat equal, even or at length obsoletely rugulose, smooth, dead white to shining white then dingy, tan, somewhat rufescent-brick-colour, fuliginous. **Gills** at the first rounded-adfixed, soon decurrent, 2-6 mm. (1-3 lin.) broad, somewhat thin, somewhat distant, *white at length light yellow-ochraceous*, connected by branches and veins, the intermediate ones at length crisped. **Milk and flesh when broken white, soon rose-colour, at length saffron-yellow.**

*Taste mild, soon acrid*, again after a time, and *when old, mild*, pleasant. Odour weak, somewhat nauseous, somewhat pungent. *Slightly firm, somewhat elastic*; when full grown and seen from above somewhat resembling *A. (Clitocybe) clavipes*. The milk sometimes remains white; it occurs also thin and watery.

In woods. Frequent. Aug.-Oct.

The colour of the pileus is exactly that of coffee and milk. *M. J. B.* The change of colour in milk and flesh is remarkable. Spores yellowish. *Fr.*; sphaeroid, echinulate, uniguttate, 6-10 mk. *K.*; almost globular, 7 mk.



*W.G.S.* Name—*fuligo*, soot. From the smoky colour. *Fr. Monogr.* ii. p. Lactarius. 178. *Hym. Eur.* p. 434. *Berk. Out.* p. 209. *C. Hbk. n.* 608. *S. Mycol. Scot.* n. 574. *Ag. Krombh. t.* 14. f. 10-12. *Barla t.* 21. f. 6, 7. *Harz. t.* 19. *Bull. t.* 567. f. 3.

**36. L. picinus** Fr.—**Pileus** 7.5 cent. (3 in.) broad, *umber or blackish-umber*, zoneless, fleshy, *convex becoming plane*, umbonate, orbicular, dry, even, at first everywhere *villous, somewhat velvety* (the down innate, not depressed or silky) then when the disc has become smooth wholly continuous on the surface (not *rivuloso-flocculose* or *squamulose*); flesh firm, but not very thick, pallid. **Stem** 5-7.5 cent. (2-3 in.) long, comparatively thin, scarcely more than 12 mm. ( $\frac{1}{2}$  in.) thick, stuffed, internally spongy, equal, *even, smooth*, paler than the pileus. **Gills** adnate, thin, *very crowded*, straight, plane, *ochraceous*. **Milk** *white, unchangeable, acrid*.

Very distinguished, rigid, regular.

In pine woods. Foxley, Hereford. Rothiemurchus. Sept.-Oct.

Spores round, *ochraceous*. *Fr.*; sphaeroid, echinulate, 5-6 mk. *C.B.P.* Name—*picinus*, pitch-black. *Fr. Monogr.* ii. p. 176. *Hym. Eur.* p. 435. *Grevillea*, vol. viii. p. 110. *S. Mycol. Scot. Supp. Scot. Nat.* 1885, p. 74. *Ag. Krombh. t.* 40. f. 20-22 (f. 19 rather *L. acris*).

**37. L. lilacinus** Lasch.—**Pileus** 5 cent. (2 in.) broad, *rosy-lilac*, becoming pale, zoneless, fleshy, thin, convex then depressed, papillate, *floccose when dry*, granular. **Stem** stuffed then hollow, *white-mealy*, pallid. **Gills** adnate, somewhat distant, pallid flesh-colour. **Milk** white, acrid.

Fragile. Milk plentiful.

In woods. Hawthornden. Sept.

Spores sphaeroid or sphaeroid-ellipsoid, echinulate, uniguttate, 7-10 mk. *K.* Name—from the *lilac* colour. *Lasch.—Fr. Hym. Eur.* p. 435. *Grevillea*, vol. viii. p. 98. *S. Mycol. Scot. Supp. Scot. Nat.* 1885, p. 74.

\*\*\* *Pileus polished, smooth.*

**38. L. volemus** Fr.—**Pileus** 7.5 cent. (3 in.) and more broad, *rufous-tawny, golden*, becoming pale, zoneless, *compact, rigid, obtuse*, plano-convex, with the margin at the first bent inwards, at length depressed, even, *at length rimoso-rivulose*, dry. **Stem** 5-6 cent. (2-2½ in.) long, 18 mm.-3 cent. ( $\frac{3}{4}$ -1¼ in.) thick, *obese, hard, solid*, somewhat equal or attenuated upwards, even, *pruinose*, of the same colour as the pileus. **Gills** adnato-decurrent, 2-4 mm. (1-2 lin.) broad, thin, crowded, white to yellowish. **Milk** *white, sweet, plentiful*.

Lactarius.

*The largest and most robust of the Russulariæ. The milk sometimes becomes yellow. Wounds on the gills become dingy.*

In mixed woods. Frequent. Sept.-Oct.

Stem sculptured longitudinally. *M. J. B.* Edible, delicious; savoury even when raw. Name—*volēma pira*, a kind of large pear. From the shape. *Fr. Monogr.* ii. p. 178. *Hym. Eur.* p. 435. *Sv. atl. Sv. t.* 10. *Berk. Out.* p. 207. *C. Hbk. n.* 609. *S. Mycol. Scot. n.* 575. *Hussey i. t.* 87. *Ag. Ventur. t.* 34. f. 1-3. *Letell. t.* 624. *Fl. Bat. t.* 874. *Krombh. t.* 39. f. 1-4.

**39. L. ichoratus** Fr.—**Pileus** 7.5-10 cent. (3-4 in.) broad, *tawny-brick-colour*, disc often brown, brick-colour and zoned, fleshy, *thin*, at first rigid, then softer, obtuse, plano-depressed, often unequal, excentric, occasionally repand, *even* (not pruinete), very dry, opaque; flesh pallid. **Stem** 4-7.5 cent. (1½-3 in.) long, 6-10 mm. (3-5 lin.) thick, solid, at first hard, then softer, internally spongy, *equal or fusiform downwards*, even, *smooth*, at first tawny, then rufescent. **Gills** adnate, decurrent with a tooth, scarcely crowded, white then ochraceous, never spotted. **Milk** white, unchangeable, *sweet*.

Allied to *L. volemus*, but thinner, *very compact when young, then softer, strong-smelling. The stem is not so obese as that of L. volemus.*

In woods. Hereford. Oct.

Name—*ιχώρ*, lymph. From the colour of the milk. *Fr. Monogr.* ii. p. 179. *Hym. Eur.* p. 436. *Grevillea*, vol. viii. p. 98. *Ag. Batsch f.* 60.

**40. L. serifluus** Fr.—**Pileus** brown-tawny, zoneless, fleshy, plane then depressed, somewhat flexuous, dry, smooth, margin inflexed. **Stem** solid, yellowish, paler than the pileus, somewhat incurved. **Gills** crowded, of the same colour as the stem. **Milk** scanty, *somewhat insipid, of the colour of serum.*

In mixed woods. Frequent. Aug.-Nov.

**Pileus** 2.5-5 cent. (1-2 in.) broad. It must be carefully distinguished from *L. camphoratus* to which it is similar in appearance and in the nature of the milk. The scent of *L. camphoratus* distinguishes it. Name—*serum*, whey; *fluo*, to flow. From the nature of the milk. *Fr. Hym. Eur.* p. 436. *Berk. Out.* p. 207. t. 13. f. 4. *C. Hbk. n.* 607. *S. Mycol. Scot. n.* 576. *Ag. Krombh. t.* 40. f. 15, 16.

**41. L. mitissimus** Fr.—**Pileus** 2.5-7.5 cent. (1-3 in.) broad, *golden-tawny*, zoneless, fleshy, thin, somewhat rigid, convex, *papillate*, depressed, papilla vanishing, even, smooth, somewhat slippery when moist; flesh pallid. **Stem** elongated, 2.5-7.5 cent. (1-3 in.) long, 8-12 mm. (½-½ in.) thick, stuffed then hollow, even, smooth, of the same colour as the pileus. **Gills** adnato-decurrent, somewhat arcuate, then tense and straight, 2-3 mm.

(1-1½ lin.) and more broad, thin, crowded, a little paler than the *Lactarius* pileus, most frequently stained with minute rufous spots. **Milk** white, *mild, plentiful*.

Thin; very much allied to *L. subdulcis*, but distinguished by the *taste* being *mild then somewhat bitterish*, and especially by the *bright, golden-tawny, resplendent* colour of the pileus and stem.

In mixed and pine woods. Common. Aug.-Nov.

Spores sphaeroid, echinulate, 6-7 mk. *C.B.P.* Name—*mitis*, mild. *Very mild* in taste. *Fr. Monogr.* ii. p. 180. *Hym. Eur.* p. 437. *Sv. atl. Sv.* t. 78. *Berk. Out.* p. 208. *C. Hbk. n.* 610. *S. Mycol. Scot. n.* 577. *Hoffm. Ic. t.* 2. *Ag. Krombh. t.* 39. f. 19-20.

**42. L. subdulcis** Fr.—**Pileus** 5-7.5 cent. (2-3 in.) broad, *rufescent*, zoneless, fleshy, thin, papillate, at length depressed, even, smooth, dry. **Stem** 2.5-5 cent. (1-2 in.) long, 4-8 mm. (2-4 lin.) thick, stuffed then hollow, equal, somewhat pruinose, of the same colour as the pileus. **Gills** adnate, crowded, paler. **Milk** *somewhat mild*, white.

Inodorous. Slightly rigid. The colour is variable, rufous-cinnamon, bay-brown-rufous, &c., but not becoming pale. Flesh rufescent, compact although thin. Gills sometimes dark ferruginous-rufous, sometimes paler. All the forms agree in the *slender growth*, in the *pileus* being *even, smooth, dry, zoneless*, and, as well as the equal *stem, rufescent*, in the *gills* being somewhat rigid, crowded, *somewhat rufescent*, in the *taste* being *somewhat mild*, and in the *milk* being *white, unchangeable*.

In woods, mixed and pine. Common. July-Nov.

The substance of the stem is looser than in *L. quietus*. *M. F.B.* Spores between papillose and echinulate, 7 mk. *W.G.S.* Name—*sub*, and *dulcis*, sweet. Somewhat sweet in taste. *Fr. Monogr.* ii. p. 180. *Hym. Eur.* p. 437. *Berk. Out.* p. 208. *C. Hbk. n.* 611. *S. Mycol. Scot. n.* 578. *Ag. Bull. t.* 227. *Sow. t.* 204. *Bolt. t.* 3. *Harz. t.* 53. *Barla t.* 20. f. 4-10. *Quel. t.* 11. f. 3. *Lenz. f.* 11.

**43. L. Terrei** B. & Br.—**Pileus** 12 mm. (½ in.) broad, bay-brown, corrugated, depressed. **Stem** 18 mm.—2.5 cent. (¾-1 in.) long, 4 mm. (2 lin.) thick, hollow, thickened at the base, of the same colour as the pileus, with orange down. **Gills** decurrent, pallid.

Cæspitose. Odour sweet. Allied to *L. subdulcis*.

On the ground. Stoke Poges, 1876. Nov.

Name—after Michael Terry. *B. & Br. n.* 1673.

**44. L. camphoratus** Fr.—**Pileus** brown-brick-red, *somewhat zoned*, fleshy, thin, depressed, dry, smooth. **Stem** stuffed, somewhat undulated, of the same colour as the pileus. **Gills** adnate, crowded, *yellowish-brick-colour*. **Milk** *mild*, white.

Lactarius.

*Strong-smelling.* So like *L. subdulcis*, that it can be distinguished safely only by its odour of *Melilot* when dried.

In woods, chiefly pine. Frequent. Aug.-Nov.

Pileus about 2.5-5 cent. (1-2 in.) broad. Stem short. The pileus is sometimes quite zoneless. The milk is scarcely white as described by Fries, rather pale watery like that of *L. seriffuus*. The unmistakable odour is slight when the plant is fresh, but very strong when dried, nor does it lose its scent with years. The milk, as well as the scent, distinguishes it from *L. subdulcis*. Spores spherical, echinulate, 6-7 mk. *Q.* Name—*camphorated*, strong-scented. *Fr. Hym. Eur. p.* 437. *Berk. Out. p.* 208. *C. Hbk. n.* 612. *S. Mycol. Scot. n.* 579. *Quel. t.* 11. f. 5. *Ag. Bull. t.* 567. f. 1. *Ventur. t.* 55. f. 3. *Barla t.* 20. f. 11-13. *Krombh. t.* 39. f. 21-24.

**45. *L. obnubilus* Lasch.**—Pileus scarcely 2.5 cent. (1 in.) broad, brown-fuliginous, zoneless, fleshy, thin, convex then umbilicate, smooth, somewhat striate. Stem stuffed then hollow, thin, paler than the pileus. Gills somewhat crowded, yellowish. Milk white, somewhat sweet.

Small, fragile; pileus at first papillate; varying in colour like *L. subdulcis*, of which it was commonly taken to be a subspecies.

In damp places below alder and in pastures. Greeshop, Morayshire. Glamis. Sept.-Nov.

Name—*obnubilo*, to overcloud, obscure. Dark. *Lasch Linn. n.* 71. under *Agaricus*. *Fr. Hym. Eur. p.* 438. *B. & Br. n.* 1784. *S. Mycol. Scot. n.* 580. *A. obscuratus Lasch. Fr. Monogr. ii. p.* 181. *Ag. Fl. Dan. t.* 1674. Compare *Ag. tristis Krombh. t.* 40. f. 26-29.

**46. *L. minimus* Smith.**—Pileus  $\frac{1}{4}$ - $\frac{5}{8}$  in. broad, pallid clay-colour, fleshy, pulvinate, rounded or slightly umbonate, margin incurved. Stem very short, generally excentric, of the same colour as the gills. Gills somewhat decurrent, arcuate, branched, moderately distant. Milk abundant, white, mild.

*Very small.*

In pastures. Abergavenny, 1871. Forres. Oct.

Spores echinulate, 8 mk. Name—*smallest* in size. *Smith Journ. Bot.* 1873, p. 205. *B. & Br. n.* 1557. *S. Mycol. Scot. n.* 581. *Fr. Hym. Eur. p.* 438.

#### TRIBE IV.—PLEUROPUS.

No British species.


GENUS X.—**Russula** (*russulus*, reddish). Pers.

Russula.

Fr. Epicr. i. p. 349.

Veil none. Hymenophore descending unchanged and forming a *vesicular trama*. Gills rigid, fragile and not milky, acute at the edge. Spores rounded, often echinulate, white or yellowish. *Growing on the ground, fleshy, putrescent, with polished stem, and pileus at the first or at length depressed.*

The Russulæ constitute a very natural group, allied to the Lactarii, but very easily distinguished from them by their want of milk. Various Russulæ however exude watery drops, especially in rainy weather. The gills of most of them are equal in length or furcate, but in the tribe of the *Compactæ* they are quite unequal. The species are for the most part striking in appearance: the *Fragiles*, the most common tribe, are difficult to determine on account of the variations of their colour. *Fr. Hym. Eur. p. 439.*



*L. Russula aurata.* One-fourth natural size.

The most protean species is *R. integra*. Fries warns beginners to beware of making out new species in the neighbourhood of this, and of ascribing forms of it to truly distinct species which are of comparatively rare occurrence.

On account of the danger of mistaking noxious species for the few which are edible, their use as food is not to be commended.

I. *COMPACTÆ* (*compingo*, to put together: compact). Pileus fleshy throughout, hence the margin is at first bent inwards and always without striæ, without a distinct viscid pellicle (in consequence of which the colour is not variable, but only changes with age and the state of the atmosphere). Flesh compact, firm. Stem solid, fleshy. Gills unequal.

II. *FURCATÆ* (*furca*, a fork. With *forked* gills). Pileus compact, firm, covered with a thin, closely adnate pellicle, which at length disappears, margin abruptly thin, at first inflexed, then spreading, *acute, even*. Stem at first compact, at length spongy-soft within. Gills *somewhat forked*, with a few shorter ones intermixed, commonly attenuated at both ends, thin and normally narrow.

III. *RIGIDÆ* (*rigidus*, rigid). Pileus without a viscid pellicle, *absolutely dry, rigid, the cuticle commonly breaking up into flocci or granules*. Flesh thick, compact, firm, vanishing away short of the *margin which is straight* (never involute), soon spreading, and always *without striæ*. Stem solid, at

Russula.

first hard, then softer and spongy. Gills, a few dimidiate, others divided, rigid, *dilated in front, and running out with a very broad rounded apex*, whence the margin of the pileus becomes obtuse and is not inflexed. *Exceedingly handsome*, but rather rare.

IV. HETEROPHYLLÆ (*R. heterophylla*, the typical species of the section). Pileus fleshy, firm, with a thin margin which is at first inflexed, then expanded and striate, covered with a thin adnate pellicle. The gills consist of many shorter ones mixed with longer ones, along with others which are forked. Stem solid, stout, spongy within.

V. FRAGILES (*fragilis*, fragile or brittle). Pileus more or less fleshy, rigid-fragile, covered with a pellicle which is always continuous, and in wet weather viscid and somewhat separable; margin membranaceous, at first convergent and not involute, in full-grown plants commonly sulcate and tubercular. Flesh commonly floccose, lax, friable. Stem spongy, at length wholly soft and hollow. Gills almost all equal, simple, broadening in front, free in the pileus when closed. Several doubtful forms occur. *R. integra* is specially fallacious from the variety of its colours.

\* Gills and spores white.

\*\* Gills and spores white, then light yellowish or bright lemon-yellow.

\*\*\* Gills and spores ochraceous.

#### I.—COMPACTÆ.

1. **R. nigricans** Fr.—Pileus 5-10 cent. (2-4 in.) and more broad, olivaceous-fuliginous, *at length black*, fleshy to the margin which is at first bent inwards, convex then flattened, umbilicato-depressed, when young and moist slightly viscid and even (without a separable pellicle), at length rimoso-squamulose; flesh firm, white, when broken becoming red on exposure to the air. **Stem** 2.5 cent. (1 in.) thick, persistently solid, equal, pallid when young, *at length black*. **Gills** rounded behind, slightly adnexed, *thick, distant*, unequal, paler, reddening when touched.

Compact, obese, inodorous, within and without *at length wholly black*, in which it differs from all others. The flesh becomes red when broken because it is saturated with red juice, although it does not exude milk. Sometimes a very few of the gills are dimidiate.

In woods. Common. June.—Nov.

Spores papillose, 8 mk. *W.G.S.* Coarse in habit. Name—*nigricans*, to be blackish. *Fr. Monogr.* ii. p. 184. *Hym. Eur.* p. 439. *Berk. Out.* p. 209. *C. Hbk.* n. 613. *S. Mycol. Scot.* n. 582. *Hussey i.* t. 73. *Ag. Bull.* t. 579. *f.* 2, t. 212. *Krombh.* t. 70. *f.* 14, 15. *Barla* t. 17. *Sow.* t. 36.

2. **R. adusta** Fr.—Pileus pallid or whitish, *cinereous-fuliginous*, equally fleshy, compact, depressed then somewhat infundibuliform, margin at first inflexed, smooth, then erect, without striæ; flesh unchangeable. **Stem** solid, obese, of the same colour as the pileus. **Gills** adnate then decurrent, *thin, crowded*, unequal, white then dingy, not reddening when touched.

It can only be compared with *R. nigricans*, but is sufficiently distinct;

stature commonly smaller, flesh juiceless, not reddening, &c. The pileus does *Russula*. not become black, but only of a scorched appearance.

In woods. Frequent. Aug.—Oct.

Well distinguished by its comparatively thin, crowded gills, &c. *M. J. B.* Spores sphaeroid, echinulate, 7–9 mk. *K.*; globose, rough, 8 mk. *C. B. P.* Name—*aduro*, to scorch. From its scorched appearance. *Fr. Monogr.* ii. p. 184. *Hym. Eur.* p. 439. *Berk. Out.* p. 209. *C. Hbk.* n. 614. *S. Mycol. Scot.* n. 583. *Ag. Pers.*—*Krombh.* t. 70. f. 7–11. *Batt.* t. 13.

**3. *R. semicrema* Fr.**—**Pileus** *persistently white*, compactly fleshy, convex then plane, even, smooth, dry, disc umbilicate, margin at first involute, always even; flesh juiceless, white, unchangeable. **Stem** sometimes curt, sometimes 5–7.5 cent. (2–3 in.) long, almost 5 cent. (2 in.) thick, fleshy, solid, firm, white, but when broken becoming cinereous internally, and *at length becoming black*. **Gills** decurrent, crowded, thin, persistently white.

Odour none, taste mild. Intermediate between *R. adusta* and *R. delicata*, but easily distinguished from both.

In mixed woods. Glamis. Aug.

Name—*semi*, half; *crêmo*, to burn. From the blackening stem. *Fr. Monogr.* ii. p. 185. *Hym. Eur.* p. 440. *Icon.* t. 172. f. 1. *B. & Br.* n. 1674. *S. Mycol. Scot.* n. 584.

**4. *R. delicata* Fr.** White.—**Pileus** 7.5–12.5 cent. (3–5 in.) broad, fleshy throughout, firm, umbilicate then infundibuliform, regular, everywhere even, smooth with a *whitish lustre*, the involute margin without striæ; flesh firm, juiceless, not very thick, white. **Stem** curt, 2.5–5 cent. (1–2 in.) long, 12 mm. ( $\frac{1}{2}$  in.) and more thick, solid, even, smooth, white. **Gills** *decurrent, thin, distant*, very unequal, white, exuding small watery drops in wet weather.

The stature and *unchangeable colours* are wholly those of *L. vellereus* and *L. piperatus*, but it is readily distinguished by the *gills being juiceless*, though they exude watery drops when young.

In mixed woods. Uncommon. Sept.—Oct.

Name—*delicus*, weaned; without juice or milk in the gills, as distinguished from *L. vellereus*, &c. *Fr. Monogr.* ii. p. 185. *Hym. Eur.* p. 440. *Berk. Out.* p. 210. *C. Hbk.* n. 615. *S. Mycol. Scot.* n. 585. *Vent.* t. 48. f. 3, 4. *Batt.* t. 17. *A. Paul.* t. 73. f. 1.

**5. *R. elephantina* Fr.**—**Pileus** 7.5 cent. (3 in.) broad, fuscous-tan, paler round the margin, but not changeable, equally fleshy, firm, at first convexo-umbilicate; margin bent inwards, smooth, undulated, and almost exceeding the gills, but never striate, covered with a pellicle which is not separable; flesh equal,

**Russula.** compact, but comparatively thin. **Stem** 5-7.5 cent. (2-3 in.) long, 2.5 cent. (1 in.) thick, solid, obese, very hard, even, smooth, externally and internally shining white. **Gills** obtuse or somewhat *sinuato-adnate*, *arcuate*, *somewhat crowded*, *thin*, divided behind, unequal in front, white, spotted pale yellowish when touched.

Habit that of *R. fetens*, but inodorous.

In woods. Stoke Poges. Sept.

Name—*elephantinus*, of ivory. From the white ivory-like stem. *Fr. Monogr.* ii. p. 186. *Hym. Eur.* p. 440. *B. & Br. n.* 1785—not Bolt.

## II.—FURCATÆ.

**6. *R. furcata* Fr.**—**Pileus** about 7.5 cent. (3 in.) broad, sometimes æruginous-greenish, sometimes umber-greenish, fleshy, compact, gibbous then plano-depressed or infundibuliform, *even*, smooth, but often *sprinkled with slightly silky lustre*, pellicle here and there separable, margin thin, at first inflexed, then spreading, always *even*; flesh firm, somewhat cheesy, white. **Stem** 5 cent. (2 in.) or a little more long, solid, firm, equal or attenuated downwards, *even*, white. **Gills** *adnato-decurrent*, *rather thick*, somewhat distant but broad, attenuated at both ends, frequently forked, shining white.

Taste bitterish. The structure is that of *R. sanguinea* or *rubra*, not of *R. virescens*, with which it agrees in colour, but from which it differs in the form of the pileus, in the acute and at first inflexed margin, in the pellicle being separable and not dividing into warts, and in the entire type of the gills.

In woods. Common. Sept.

Poisonous. Spores globose, echinulate, 6-7 mk. *C.B.P.* Name—*furca*, a fork. With *forked gills*. *Fr. Monogr.* ii. p. 187. *Hym. Eur.* p. 441. *Berk. Out.* p. 210. *C. Hbk. n.* 616. *S. Mycol. Scot. n.* 586. *Ag. Pers.*—*Krombh. t.* 62. *f.* 1, 2, *t.* 69. *f.* 18-22. *Bull. t.* 26. *Schæff. t.* 94. *f.* 1. *Barla t.* 16. *f.* 1-9. *Harz. t.* 54, *t.* 63. *f.* 5. *Paul. t.* 74. *f.* 1. *Buxb. C. v. t.* 47. *f.* 2.

**7. *R. sanguinea* Fr.**—**Pileus** 5-7.5 cent. (2-3 in.) broad, blood-red or becoming pale round the *even*, spreading, *acute margin*, fleshy, firm, at first convex, obtuse, then depressed and infundibuliform and commonly gibbous in the centre, polished, *even*, *moist* in damp weather; flesh firm, cheesy, white. **Stem** stout, spongy-stuffed, at first contracted at the apex, then equal, slightly striate, white or reddish. **Gills** at first adnate, then truly decurrent, very crowded, very narrow, connected by veins, fragile, somewhat forked, shining white.

Taste *acrid*, peppery. Often confounded with *R. rubra*, which is of the



same colour, but entirely different from it in the firm solid flesh, in the gills *Russula*. being adnate, then deeply decurrent, and acuminate in front.

In woods, chiefly fir. Uncommon. Aug.-Sept.

Poisonous. Name—*sanguis*, blood. Blood-coloured. *Fr. Monogr.* ii. p. 188. *Hym. Eur.* p. 442. *Berk. Out.* p. 210. *C. Hbk.* n. 617. *S. Mycol. Scot.* n. 587. *Ag. Bull.* t. 42.

**8. *R. rosacea* Fr.**—**Pileus** 5-10 cent. (2-4 in.) broad, somewhat flesh-coloured varying in intensity, becoming whitish when the pellicle disappears, often *variegated with darker spots when dry*, compactly fleshy, at first convex, then expanded, obtuse, commonly *unequal*, repand, even incised, covered with a pellicle which is *viscid* and separable in wet weather, *margin acute, even*; flesh firm, cheesy, white. **Stem** about 5 cent. (2 in.) long, solid, firm, at length spongy internally, even, smooth, occasionally ventricose, white or reddish. **Gills** in every stage of growth adnate, thin, crowded, fragile, forked behind, with dimidiate ones intermixed, always persistently white.

Taste *slowly acrid*. Allied to *R. sanguinea* but irregular, often excentric with the pileus somewhat repand, *scarcely depressed*, and the gills less crowded, broader, less divided, scarcely connected.

In mixed woods. Frequent. Sept.-Oct.

Spores papillose, almost globose, 7 mk. *W.G.S.* Name—*rosa*, a rose. Rose-coloured. *Fr. Monogr.* ii. p. 188. *Hym. Eur.* p. 442. *Berk. Out.* p. 210. *C. Hbk.* n. 618. *S. Mycol. Scot.* n. 588. *Ag. Bull.* t. 509. f. Z.

**9. *R. sardonia* Fr.**—**Pileus** 5-7.5 cent. (2-3 in.) broad, reddish, &c., fleshy, compact, convex then plane, rarely depressed, but here and there repand, with an adnate pellicle which is viscid in wet weather and soon changes colour—and then often spotted, *margin even*. **Stem** 4-5 cent. (1½-2 in.) long, almost 2.5 cent. (1 in.) thick, solid, firm, but at length spongy within, even, white or reddish. **Gills** adnate, *crowded*, broad, somewhat forked, *white*, exuding watery drops in wet weather, whence arise *yellowish* spots when dry.

Robust, firm. The colour is very changeable, sometimes reddish, sometimes pallid with yellow spots, sometimes dingy yellow, opaque. Flesh same as in *R. rosacea*, &c. Intermediate between *R. rosacea* and *R. expallens*, but distinct from both in the colour becoming yellow.

In woods, chiefly fir. Uncommon. Sept.

Name—from its acrid taste. *Herba Sardonia* (probably *Ranunculus sceleratus*), screwing the mouth with its bitterness. *Fr. Monogr.* ii. p. 189. *Hym. Eur.* p. 442. *Berk. Out.* p. 211. *C. Hbk.* n. 619. *S. Mycol. Scot.* n. 589. *Ag. Krombh.* t. 68. f. 1-4.—*Schæff.* t. 16. f. 5, 6.

**10. *R. depallens* Fr.**—**Pileus** pallid reddish or inclining to

Russula.

fuscous, &c., fleshy, firm, convex then plane, more rarely depressed, but commonly *irregularly shaped and undulated*, even, the thin adnate pellicle presently changing colour especially at the disc, the spreading margin even, but slightly striate when old; flesh white. **Stem** about 4 cent. ( $1\frac{1}{2}$  in.) long, solid, firm, commonly attenuated downwards, *white, becoming cinereous* when old. **Gills** adnexed, broad, crowded, distinct, but commonly forked at the base, often with shorter ones intermixed.

Inodorous, taste mild. The colour of the pileus is at first pallid reddish or inclining to fuscous, then whitish or yellowish, opaque in every stage of growth. It approaches nearest to the *Heterophyllæ*.

In beech woods, pastures, &c. Uncommon. Aug.-Sept.

Name—*de*, and *paleo*, to be pale. Becoming pale. *Fr. Monogr.* ii. p. 189. *Hym. Eur.* p. 442. *Berk. Out.* p. 211. *C. Hbk.* n. 620. *S. Mycol. Scot.* n. 590. *Krombh.* t. 66. f. 12, 13.

**11. R. drimeia** Cke.—**Pileus** 5–10 cent. (2–4 in.) broad, bright purple, compact, firm, convex then depressed, scarcely viscid when moist, opaque when dry, margin somewhat incurved, even. **Stem** 5–7.5 cent. (2–3 in.) long, 12–18 mm. ( $\frac{1}{2}$ – $\frac{3}{4}$  in.) thick, solid, firm, cylindrical, equal, tinged with purple. **Gills** adnexed, scarcely crowded, narrow and furcate at the base, at first pale sulphur-yellow, then deeper yellow, never white.

Acrid, peppery. Spores pale ochre. The colour and habit similar to *R. Queletii*, but distinguished by the yellow gills, ochraceous spores, and intensely peppery taste.

On the ground among larch. Black Park, 1881, &c. Sept.-Nov.

Name—*δριμύς*, pungent. *Cke. in Grevillea*, vol. x. p. 46.

### III.—RIGIDÆ.

**12. R. lactea** Fr.—**Pileus** 5 cent. (2 in.) broad, at the first *milk-white, then tan-white, throughout compactly fleshy*, campanulate then convex, often excentric, without a pellicle, always dry, at the first even, then slightly cracked when dry, margin straight, thin, obtuse, even; flesh compact, white. **Stem** 4–5 cent. ( $1\frac{1}{2}$ –2 in.) long, 4 cent. ( $1\frac{1}{2}$  in.) thick, solid, very compact, but at length spongy-soft within, equal, even, always white. **Gills** *free*, very broad, *thick, distant*, rigid, forked, white.

Mild. The gills are at length adnate, forked at the base and apex. Care must be taken not to confound it with other *Russulæ* which have changed colour and become white.

In mixed woods. Uncommon. Aug.

Edible, but not commended for use. Name—*lac*, milk. Milk-white. *Fr.*

*Monogr.* ii. p. 190. *Hym. Eur.* p. 443. *B. & Br.* 1133. *C. Hbk.* n. 621. *Russula.*  
*S. Mycol. Scot.* n. 591. *Ag. Pers.*—*Krombh.* t. 61. f. 1-2. *Barla* t. 15. f. 1-13.  
*Paul.* t. 74. f. 2.

**13. R. virescens** Fr.—**Pileus** green, compactly fleshy, globose then expanded, at length depressed, often unequal, always dry, not furnished with a pellicle, wherefore the *flocculose cuticle is broken up into patches or warts*, margin straight, obtuse, *even*; flesh white, not very compact. **Stem** solid, internally spongy, firm, *somewhat rivulose*, white. **Gills** free, somewhat crowded, sometimes equal, sometimes forked, with a few shorter ones intermixed, *white*.

Taste mild. It varies in size and colour of pileus, which is sometimes deep sometimes pallid green, sometimes yellowish then green. The gills are not so broad in front as those of neighbouring species. It is very easily distinguished from all others by the *green* pileus being without a pellicle and *innato-flocculose*, then rivulose, and scaly in the form of patches.

In woods. Frequent. July–Sept.

Stem variable in form, slightly reticulated with raised lines. *M. F. B.* Pileus sometimes as much as 10 cent. (4 in.) broad. Edible. Spores scarcely echinulate, almost globular, 6 mk. *W. G. S.* Name—*vireo*, to be green. *Fr. Monogr.* ii. p. 190. *Hym. Eur.* p. 443. *Berk. Out.* p. 212. t. 13. f. 6. *C. Hbk.* n. 622. *S. Mycol. Scot.* n. 592. *Hussey* ii. t. 11. *Ag. Schæff.* t. 94 excluding f. 1. *Vittad.* t. 31. *Sturm Deutschl. Fl.* iii. 3. t. 31. *Barla* t. 16. f. 10-12. *Ventur.* t. 17. f. 1, 2. *Krombh.* t. 67. f. 1-10.

**14. R. Du Portii** Phill.—**Pileus** 4–6.5 cent. ( $1\frac{1}{2}$ – $2\frac{3}{4}$  in.) broad, centre rufous or flesh-red, margin bluish, compact, fleshy, firm, convexo-plane, depressed, smooth, dry, margin even, obtuse. **Stem** 2.5 cent. (1 in.) or more high, 10–16 mm. (5–8 lin.) broad, spongy-stuffed, minutely striate, glabrous, white. **Gills** rounded behind, broad, distant, white.

The flesh turns reddish-brown when cut, and the odour is that of the common crab.

In wood. Mattishall, Norfolk.

Spores globose, asperate, 9 mk. *W. P.* Name—after Rev. J. M. Du Port. *Phill. Grevillea*, vol. xiii. p. 49.

**15. R. lepida** Fr.—**Pileus** 7.5 cent. (3 in.) broad, *blood-red-rose*, becoming pale, whitish especially at the disc, somewhat equally fleshy, convex then expanded, scarcely depressed, obtuse, opaque, unpolished, *with a silky appearance, at length often rimoso-squamulose*, margin spreading, obtuse, without striæ. **Stem** as much as 7.5 cent. (3 in.) long, often 2.5 cent. (1 in.) thick, *even, white or rose-colour*. **Gills** rounded behind, rather thick, somewhat crowded, often forked, connected by veins, white, often red at the edge.

Russula.

Taste mild; wholly compact and firm, but the flesh is cheesy not somewhat grumous. The gills are often red at the edge chiefly towards the margin, on account of the margin of the pileus being continuous with the gills.

In mixed woods. Frequent. Sept.-Oct.

Edible. Name—*lepidus*, neat, elegant. *Fr. Monogr.* ii. p. 191. *Hym. Eur.* p. 444. *Sv. ätl. Sv. t.* 59. *Berk. Out.* p. 212. *C. Hbk. n.* 623. *S. Mycol. Scot. n.* 593. *Hussey ii. t.* 32. *Hogg & Johnst. t.* 4. *Ag. Krombh. t.* 64. f. 19, 20. *Batsch t.* 13 (*very small*).

**16. R. rubra** Fr.—**Pileus** unicolorous, cinnabar-vermilion, but becoming pale (tan) when old, disc commonly darker, compact, hard but fragile, convex then flattened, here and there depressed, absolutely dry, *without a pellicle, but becoming polished-even*, often rivuloso-rimose when old, margin spreading, obtuse, even, always persistent; flesh white, *reddish under the cuticle*. **Stem** 5-7.5 cent. (2-3 in.) long, about 2.5 cent. (1 in.) thick, solid, even, varying white and red. **Gills** obtusely adnate, somewhat crowded, whitish, then yellowish, with dimidiate and forked ones intermixed.

Very *acid*, very hard and rigid, most distinct from all the others of this group in the *pileus becoming polished-even*, although without a pellicle, in the *flesh being somewhat grumous*, and in the *very acid taste*. Gills often red at the edge.

In mixed woods. Frequent. Aug.-Nov.

Poisonous. Spores whitish. *Fr.*; sphaeroid, 8-10 mk. *K.* Name—*ruber*, red. *Fr. Monogr.* ii. p. 191. *Hym. Eur.* p. 444. *Sv. ätl. Sv. t.* 49. *Berk. Out.* p. 212. *C. Hbk. n.* 624. *S. Mycol. Scot. n.* 594. *Ag. Decand.*—*Barla t.* 15. f. 1-10. *Krombh. t.* 65. *Vitt. Mang. t.* 38. f. 2, not Bull. *Schæff. t.* 15. f. 4-6.

**17. R. Linnæi** Fr.—**Pileus** 7.5-10 cent. (3-4 in.) broad, unicolorous, dark purple, blood-red or bright rose, opaque, not becoming pale, everywhere fleshy, rigid, plano-depressed, sometimes repand, even, smooth, *dry, without a separable pellicle*, margin spreading, obtuse, without striæ; flesh thick, *spongy-compact, white*. **Stem** 4 cent. (1½ in.) and more long, 2.5 cent. (1 in.) and more thick, stout, firm, but spongy-soft within, somewhat ventricose, *obsoletely fibrilloso-reticulate*, intensely blood-red. **Gills** adnate, *somewhat decurrent*, rather thick, not crowded, *broad* (more than 12 mm., ½ in.), fragile, sparingly connected by veins, white, becoming yellow when dry, with a few dimidiate ones intermixed, somewhat anastomosing behind.

*Taste mild.* The habit is exactly that of *R. emetica*; the substance also is floccose, but very compact, firm, and thick. The stem according to v. Post is hollow.

In woods. Stoke Poges.

Spores wholly white. *Fr.*; ellipsoid-sphæroid, echinulate, 11 mk. *Q.* Russula. Name—after Linnæus. *Fr. Monogr.* ii. p. 192. *Hym. Eur.* p. 444. *Icon. t.* 172. f. 3. *B. & Br. n.* 1786.

**18. *R. xerampelina* Fr.**—**Pileus** 7.5 cent. (3 in.) and more broad, *rosy-purple*, &c., disc becoming pale, fleshy, compact, convex then flattened, at length depressed, *without a distinct pellicle*, wherefore always dry, opaque, even, but when older commonly very thinly *rimulose*, so that the cuticle under a lens is very thinly *granulated or punctate*, margin spreading, without striæ; flesh *compact, white inclining to yellowish*. **Stem** 5–7.5 cent. (2–3 in.) long, 2.5 cent. (1 in.) thick, stout, firm, internally at length soft, spongy, even hollow, commonly thickened at the base, even, white or reddish. **Gills** adnexed, somewhat crowded, broader in front, forked behind, *whitish*, then yellowish.

*Taste mild.* The colour of the pileus is changeable, commonly rosy-purple, with the disc becoming pale, yellowish-white, sometimes inclining to olivaceous. The stem is redder when moist. It approaches *R. integra*, but differs in the pileus not having a distinct pellicle, &c.

In Scotch fir wood. Glamis, 1876. Oct.

Spores dingy tan. *Fr.* A very distinct species. Name—*ξεραπέλινα*, of the colour of dry vine-leaves. Dark rosy red. *Fr. Monogr.* ii. p. 192. *Hym. Eur.* p. 445. *B. & Br. n.* 1675. *S. Mycol. Scot. n.* 595. *Ag. Schæff. t.* 214, 215.

**19. *R. olivacea* Fr.**—**Pileus** dingy purple then olivaceous or wholly fuscous-olivaceous, fleshy, convexo-flattened and depressed, *slightly silky and squamulose*, margin spreading, even; *flesh white, becoming somewhat yellow*. **Stem** firm, ventricose, rose-colour to pallid, spongy-stuffed within. **Gills** adnexed, wide, *yellow*, with shorter and forked ones intermixed.

*Mild.* Near to *R. rubra*, but certainly distinct in the stem being definitely spongy, in the pileus being unpolished, and in the gills being soft and brightly coloured; corresponding with *R. alutacea*.

In woods. Slough.

Spores light yellow. *Fr.*; sphæroid, punctate, 10 mk. *Q.* Name—*oliva*, an olive. Olivaceous in colour. *Fr. Hym. Eur.* p. 445. *B. & Br. n.* 1562. *Ag. Schæff. t.* 204. *Krombh. t.* 68. f. 13.

**20. *R. cutefracta* Cke.**—**Pileus** 7.5–10 cent. (3–4 in.) or more broad, green, purple, dull red, &c., fleshy, firm, dry, opaque, convex then a little depressed in the centre, cuticle cracking from the margin inwards into minute firm adnate areolæ, otherwise even; flesh beneath the cuticle tinged with purple. **Stem** 7.5 cent. (3 in.) long, often 2.5 cent. (1 in.) thick, firm, solid, nearly equal, or a little attenuated above, smooth, slightly tinged

*Russula*. with purple. **Gills** somewhat crowded, narrowed behind, furcate, adnexed or nearly free, white.

Mild. Allied to *R. virescens* which it resembles in the cracking of the cuticle, but differs in the purple tint beneath even in green specimens, and in the tinted stem, as well as in the colour of the pileus, which is of a darker and different shade of green, and sometimes of a deep bluish-purple, as well as of a madder-red.

On the ground in woods. Epping, &c.

Name—*cutis*, skin; *frango*, to break. From the cracked cuticle. *Cke. in Grevillea*, vol. x. p. 46.

#### IV.—HETEROPHYLLÆ.

**21. *R. vesca*** Fr.—**Pileus** *red-flesh-colour, disc darker*, fleshy, slightly firm, plano-depressed, *slightly wrinkled with veins*, with a viscid pellicle, margin at length spreading; flesh cheesy, firm, shining white. **Stem** *solid*, compact, externally rigid, *reticulated and wrinkled* in a peculiar manner, often attenuated at the base, shining white. **Gills** adnate, crowded, thin, shining white, with many unequal and forked ones intermixed, but scarcely connected by veins.

Of middle stature. *Taste mild*, pleasant.

In mixed woods. Frequent. Sept.—Oct.

Reckoned edible. Name—*vesco*, to feed. From its edible qualities. *Fr. Monogr.* ii. p. 193. *Hym. Eur.* p. 446. *Sv. atl. Sv. t.* 63. *Berk. Out.* p. 211. *C. Hbk. n.* 625. *S. Mycol. Scot. n.* 596. (*Hussey i. t.* 89.)

**22. *R. cyanoxantha*** Fr.—**Pileus** 5–7.5 cent. (2–3 in.) and more broad, *lilac or purplish then olivaceous-green*, disc commonly becoming pale often yellowish, *margin* commonly becoming *azure-blue or livid-purple*, compact, convex then plane, then depressed or infundibuliform, sometimes even, sometimes wrinkled or streaked, viscous, margin deflexed then expanded, remotely and slightly striate; flesh firm, cheesy, white, commonly reddish beneath the separable pellicle. **Stem** 5–7.5 cent. (2–3 in.) long, as much as 2.5 cent. (1 in.) thick, *spongy-stuffed*, but firm, often cavernous within when old, equal, smooth, *even*, shining white. **Gills** rounded behind, connected by veins, not much crowded, broad, forked with shorter ones intermixed, shining white.

Allied to *R. vesca* in its *mild*, pleasant *taste* and in other respects, but constantly different in the colour of the pileus, which is very variable, whereas in *R. vesca* it is unchangeable. The peculiar combination of colours in the pileus, though very variable, always readily distinguishes it.

In mixed woods, &c. Common. Aug.—Oct.

Sometimes considerably larger than Fries describes. Name—*κύανος*, blue;

ξανθός, yellow. From the colours. *Fr. Monogr.* ii. p. 194. *Hym. Eur.* p. Russula. 446. *B. & Br.* n. 1131. *C. Hbk.* n. 626. *S. Mycol. Scot.* n. 597. *Ag. Schæff.* t. 93. *Krombh.* t. 67. f. 16-19. *Paul.* t. 76. f. 1-3.

**23. *R. heterophylla* Fr.**—**Pileus** very variable in colour, but *never becoming reddish or purple*, fleshy, firm, convexo-plane then depressed, *even, polished*, the very thin pellicle disappearing, margin thin, even or densely but slightly striate; flesh white. **Stem** solid, firm, somewhat equal, *even*, shining white. **Gills** *reaching the stem in an attenuated form, very narrow, very crowded*, forked and dimidiate, shining white.

Taste *always mild*, as in *R. cyanoxantha*, from which it differs in its smaller stature, in the pileus being thinner, even, *never reddish* or purplish, with a thin closely adnate pellicle, in the stem being *firm and solid*, and in the gills being *thin, very narrow, very crowded*, &c. The apex of the stem is occasionally dilated in the form of a cup, so that the gills appear remote.

In woods. Common. July–Oct.

Edible, of a sweet nutty flavour. Spores echinulate, 5×7 mk. *W.G.S.* Name—ἔτερος, φύλλον, a leaf. With gills of different lengths. *Fr. Monogr.* ii. p. 194. *Hym. Eur.* p. 446. *Berk. Out.* p. 211. t. 13. f. 5. *C. Hbk.* n. 627. *S. Mycol. Scot.* n. 598. *Hogg & Johnst.* t. 9. *Hussey* i. t. 84. *Badh.* i. t. 10. f. 3; ii. t. 3. f. 3, 4. *Ag. Fl. Dan.* t. 1909. f. 1. *Paul.* t. 75. f. 1-5.

\* ***R. galochroa* Fr.** Smaller.—**Pileus** *at the first milk-white*, then greenish, rather plane, viscid in wet weather, commonly dry, even, sometimes sprinkled with *white floccose spots*, slightly striate at the margin. **Stem** 2.5–5 cent. (1–2 in.) long, 12 mm. (½ in.) thick, solid, firm, never becoming red. **Gills** very thin, more or less forked and unequal.

The flocci at the disc in the form of spots or warts are not like those of *R. virescens*. In woods. Slough. Name—γάλα, milk; χρώς, colour. Milk-white. *Fr. Monogr.* ii. p. 195. *Hym. Eur.* p. 447. *B. & Br.* n. 1563. *Ag. Bull.* t. 509. f. L.M. *Batt.* t. xii. f. E.

**24. *R. consobrina* Fr.**—**Pileus** 7.5 cent. (3 in.) broad, *dark cinereous* or fuscous olivaceous, fleshy, fragile, campanulate then expanded, at length depressed, margin spreading, even, though membranaceous; *flesh white, cinereous under the thick, viscous, separable pellicle*. **Stem** 5–7.5 cent. (2–3 in.) long, almost 2.5 cent. (1 in.) thick, solid, but *soft*, equal, even, smooth, shining white, *at length becoming cinereous*. **Gills** at the first free, then appearing adnate when the pileus is flattened, broad, crowded, shining white, *very many of them dimidiate and forked*.

Taste *very acrid*. Not foetid. Stature in general that of *R. emetica*, but differing in the colour of the pileus, and in the very unequal gills.

In mixed woods. Glamis, 1876. Oct.

Russula.

Spores granular, 10 mk. *Q.* Name—*consobrinus*, cousin. *Distantly related* to neighbouring species. *Fr. Monogr.* ii. p. 195. *Hym. Eur.* p. 447. *B. & Br.* n. 1676. *S. Mycol. Scot.* n. 599.

\* **R. sororia** Larbr.—**Pileus** convex then plane or depressed, *striate at the margin*. **Gills** somewhat distant, connected by veins.

The habit and colour are the same, but it differs by many marks. Stem white. Many of the gills dimidiate, but scarcely forked. Name—*soror*, a sister. From its relation to *R. consobrina*. In fir woods. North Wooton. *Aug. Larbr. t.* 19. f. 7. *Fr. Hym. Eur.* p. 447. *Icon. t.* 173. f. 1. *Grevillea*, vol. x. p. 67.

**25. R. fœtens** Fr.—**Pileus** 10–12.5 cent (4–5 in.) and more broad, dingy yellow, often becoming pale, thinly fleshy, at first bullate, then expanded and depressed, covered with a pellicle which is adnate, not separable, and viscid in wet weather, margin broadly membranaceous, at the first bent inwards *with ribs which are at length tubercular*; flesh thin, *rigid-fragile*, pallid. **Stem** 5 cent. (2 in.) and more long, 1–2.5 cent. ( $\frac{1}{2}$ –1 in.) thick, stout, stuffed then hollow, whitish. **Gills** adnexed, crowded, connected by veins, with very many *dimidiate and forked* ones intermixed, whitish, at the first *exuding watery drops*.

*Fœtid. Taste acrid.* Very *rigid*, most distinct from all others in its *very heavy empyreumatic odour*. In very dry weather the odour is often obsolete. The margin is more broadly membranaceous and hence marked with *longer furrows* than in any other species. It differs from all the preceding ones in the gills at the first exuding watery drops. The gills become obsoletely light yellow, and dingy when bruised.

In woods, &c. Very common. July–Sept.

Stem ruggedly hollow within as if eaten by snails. *M. J. B.* A very coarse and easily recognised species. Reckoned poisonous, though eaten by slugs. Spores minutely echinulate, almost globular, 8 mk. *W. G. S.* Name—*fœtens*, stinking. *Fr. Monogr.* ii. p. 195. *Hym. Eur.* p. 447. *Sv. atl. Sv. t.* 40. *Berk. Out. p.* 213. *C. Hbk. n.* 628. *S. Mycol. Scot. n.* 600. *Ag. Pers.—Krombh. t.* 70. f. 1–6. *Viv. t.* 41. *Bull. t.* 292. *Ventur. t.* 33. f. 1–3.

**26. R. subfœtens** Smith.—**Pileus** bullate, somewhat viscid, disc fleshy, margin somewhat membranaceous. **Stem** not stout. **Gills** thick, distant, branched.

Odour somewhat disagreeable, taste slightly acrid. Smaller than *R. fœtens*, stem not so stout, margin different. This is the plant referred to by Fries *Syst. Myc. i. p.* 58. as a var. of *R. fragilis*.

On the ground. Various localities.

Name—*sub*, and *fœtens*, somewhat stinking. *Worth. Smith Journ. Bot.* 1873, p. 337.

**27. R. fellea** Fr.—**Pileus** 2.5–5 cent. (1–2 in.) broad, *straw-*



*colour often with a deeper yellowish tinge*, disc darker, thinly fleshy, *Russula*. convex then plane, polished, smooth, margin even, slightly striate when old; pellicle closely adnate, thin; *flesh firm, of the same colour as the gills*. **Stem** 5 cent. (2 in.) long, 12 mm. ( $\frac{1}{2}$  in.) thick, spongy-stuffed then hollow, fragile, equal, even, white then *straw-colour*. **Gills** adnate, crowded, thin, narrow, obsoletely connected by veins, intermixed with a few which are dimidiate or bifid behind, *straw-colour*.

Distinguished, *very acrid*, odour none. The gills exude watery drops in damp weather.

In beech woods. Common. Aug.—Nov.

Easily distinguished from all others by its unchangeable persistent yellowish straw-colour. The gills vary more distant. Name—*felleus*, full of gall. Bitter. *Fr. Monogr.* ii. p. 196. *Hym. Eur.* p. 447. *Icon. t.* 173. *f.* 2. *B. & Br. n.* 1787. *S. Mycol. Scot. Supp. Scot. Nat.* 1881, p. 36. *Paul. t.* 76. *f.* 4.

**28. R. Queletii Fr.**—**Pileus** *violaceous-black* or fuscous, *purple-lilac* at the margin which is slightly striate, compact, campanulate-convex then plane, *even*, viscous; flesh firm, white, red-purple under the pellicle. **Stem** spongy, *mealy*, violaceous-purple. **Gills** attenuated, unequal or forked, *exuding drops*, white.

*Acrid*. When the drops on the gills are dried, they leave azure-blue-cinereous or pallid olivaceous spots.

In woods. Common. Aug.—Nov.

Pileus commonly about 7.5 cent. (3 in.) broad; stem about same in length and 2.5 cent. (1 in.) thick. Easily distinguished from *R. integra* by the white gills. Name—after L. Quélet. *Fr. in Quél. Jur.* p. 185. *t.* 24. *f.* 6. *Hym. Eur.* p. 448. *B. & Br. n.* 1888. *S. Mycol. Scot. n.* 601.

#### V.—FRAGILES.

\* *Gills and spores white.*

**29. R. emetica Fr.**—**Pileus** 7.5–10 cent. (3–4 in.) broad, at first rosy then *blood-colour*, tawny when old, sometimes becoming yellow and at length (in moist places) white, at first campanulate then flattened or depressed, polished, *margin* at length *sulcate and tubercular*; *flesh white, reddish under the separable pellicle*. **Stem** spongy-stuffed, stout, elastic when young, fragile when older, even, white or reddish. **Gills** somewhat *free*, broad, somewhat distant, shining white.

Handsome, regular, moderately firm, but fragile when full grown, *taste very acrid*. The gills are never dusty. There is one form taller, with the pileus campanulate then expanded, and the gills persistently free, another with the pileus convex then expanded, and gills appearing adfixed.

Russula.

In woods and open ground. Common. July-Dec.

Spores shining white, *Fr.*; sphaeroid, echinulate, 8-10 mk. *K.*; 7 mk. *W.G.S.* Stem longitudinally rugulose. *M.F.B.* Easily distinguished by the flesh remaining red under the cuticle when peeled off. Poisonous. Said to act as its name implies as an emetic; certainly dangerous. Name—*emetica*, a vomit. *Fr. Monogr.* ii. p. 197. *Hym. Eur.* p. 448. *Sv. ätl. Sv. t.* 21. *Berk. Out.* p. 212. *C. Hbk. n.* 629. *S. Mycol. Scot. n.* 602. *Ag. Harz. t.* 63. *Barl. t.* 14. f. 4. *Lenz f.* 15.

**30. *R. pectinata* Fr.—Pileus** 7.5 cent. (3 in.) broad, at first viscous, *toast-brown*, then dry, becoming pale, tan, with the *disc* always *darker*, fleshy, *rigid*, convex then flattened and depressed or concavo-infundibuliform; margin thin, *pectinato-sulcate*, here and there irregularly shaped; *flesh white, light yellowish under the pellicle*, which is not easily separable. **Stem** curt, 5 cent. (2 in.) long, 18 mm.—2.5 cent. ( $\frac{3}{4}$ —1 in.) thick, *rigid*, spongy-stuffed, longitudinally *slightly striate, shining white*, often attenuated at the base. **Gills** *attenuato-free* behind, broader towards the margin, somewhat crowded, *equal*, simple, white.

Odour weak, but nauseous, approaching that of *R. fetens*. The gills are narrower than those of neighbouring species, but dilated from the stem to the margin, not exuding drops, and not changeable when touched.

In mixed woods. Glamis, 1874. Aug.

Name—*pecten*, a comb. From the furrows at the margin being like the teeth of a comb. *Fr. Monogr.* ii. p. 197. *Hym. Eur.* p. 449. *B. & Br. n.* 1564. *S. Mycol. Scot. n.* 603. *Ag. Bull. t.* 509. f. N.

**31. *R. ochroleuca* Fr.—Pileus** *yellow, becoming pale*, fleshy, flattened or depressed, polished, with an adnate pellicle, the spreading margin *becoming even*. **Stem** spongy, stuffed, firm, *slightly reticulato-wrinkled, white becoming cinereous*. **Gills** *rounded behind*, united, broad, *somewhat equal*, white becoming pale.

Odour obsolete, but pleasant. The pileus is never reddish. It agrees wholly with *R. emetica* in structure and stature, as well as in the *acid taste*; it differs however in the stem being slightly reticulato-wrinkled, white becoming cinereous, in the adnate pellicle of the pileus, in the margin remaining for a long time *even* (remotely striate, but not tubercular, only when old), and in the gills being rounded behind and becoming pale. The colour of the pileus is constant. The gills remain free and do not exude drops.

In woods, chiefly fir. Frequent. Aug.—Nov.

Spores papillose, 7 mk. *W.G.S.* Name—*ὀχρός*, pale yellow; *λευκός*, white. From the yellow pileus and white stem and gills. *Fr. Hym. Eur.* p. 449. *Monogr.* ii. p. 197. *Berk. Out.* p. 212. *C. Hbk. n.* 630. *S. Mycol. Scot. n.* 604. *Ag. Pers.—Krombh. t.* 64. f. 7-9. *Buxb. C. v. t.* 45. f. 2.

**32. *R. fragilis* Fr.—Pileus** 2.5-4 cent. (1-1½ in.) broad, rarely

more, flesh-colour, changing colour, very thin, fleshy only at the disc, at the first convex and often umbonate, then plane and depressed, pellicle thin, becoming pale, slightly viscid in wet weather; *margin* very thin, *tuberculo-striate*. **Stem** 4-5 cent. ( $1\frac{1}{2}$ -2 in.) long, spongy within, soon hollow, often slightly striate, white. **Gills** slightly adnexed, very *thin*, *crowded*, broad, *ventricose*, all equal, shining white. Russula.

Very acrid. Smaller and more fragile than the rest of the group, directly changing colour. The colour is variable, often opaque, typically flesh-colour, when changed in colour white externally and internally, often with reddish spots. Among varieties of colour is to be noted a livid-flesh-coloured form, with the disc becoming fuscous.

It is not easy to define it from fragile forms of *R. emetica*, but the gills are much more crowded, thinner, and often slightly eroded at the edge, ventricose; the pileus thinner and more lax, &c.

In woods. Common. Aug.-Oct.

Name—*fragilis*, fragile or brittle. *Fr. Monogr.* ii. p. 198. *Hym. Eur.* p. 450. *Berk. Out.* p. 213. *C. Hbk.* n. 631. *S. Mycol. Scot.* n. 605. *Ag. Pers.* —*Krombh.* t. 64. f. 12-18. *Bull.* t. 509. f. T, U. *Barla* t. 14. f. 10-12. *Corda ap. Sturm* xi. t. 53. *Vent.* t. 33. f. 4, 5.

**\*\* Gills and spores white then light yellowish, &c.**

**33. R. veternosa** Fr.—**Pileus** 5 cent. (2 in.) broad, rose or flesh-colour, soon becoming pale, commonly whitish or yellowish at the disc, slightly fleshy, plane when full grown, depressed in the middle, covered with a thin, adnate, somewhat viscid pellicle, *margin even*, scarcely membranaceous; flesh soft, white. **Stem** 5-7.5 cent. (2-3 in.) long, 12 mm. ( $\frac{1}{2}$  in.) thick, *soft, spongy then hollow, fragile*, equal, even, white. **Gills** adnate, crowded, narrow, becoming broader in front, with shorter ones intermixed, *white then straw-colour*.

Taste acrid.

On the ground. Chatteris.

Name—*veternosus*, drowsy, languid. *Fr. Monogr.* ii. p. 201. *Hym. Eur.* p. 450. *B. & Br.* n. 1132. *C. Hbk.* n. 635. *Ag. Krombh.* t. 66. f. 18, 19. *Paul.* t. 74. f. 3.

**34. R. integra** Fr.—**Pileus** typically red, changing colour, fleshy, campanulato-convex then expanded and depressed, fragile when full grown, with a viscous pellicle, at length *sulcate and somewhat tubercular* at the margin; *flesh white*, sometimes yellowish above. **Stem** at first short, conical, then clavate or *ventricose*, as much as 7.5 cent. (3 in.) long, spongy-stuffed, commonly stout, *even*, shining *white*. **Gills** somewhat free, very broad, equal or bifid at the stem, somewhat distant, connected

Russula.

by veins, pallid-white, at length light yellow, *somewhat ochraceous-pulverulent with the spores*.

*Taste mild*, often astringent. The most changeable of all species, especially in the colour of the pileus which is typically red, but at the same time inclining to azure-blue, bay-brown, olivaceous, &c. Sometimes the gills are sterile and remain white.

In woods, chiefly pine. Common. Aug.-Nov.

Pileus 7.5-10 cent. (3-4 in.) broad. The white stem and yellow-pulverulent gills distinguish it. Spores ellipsoid-sphaeroid or sphaeroid, echinulate, 10 × 8 or 7 × 9 mk. *K.*; globose, rough, 8-9 mk. *C.B.P.* Name—*integer*, whole, entire. Perfect in form. *Fr. Monogr.* ii. p. 199. *Hym. Eur.* p. 450. *Berk. Out.* p. 213. *C. Hbk.* n. 632. *S. Mycol. Scot.* n. 606. *Ag. Schaff.* t. 92. *Vittad.* t. 21. *Vent.* t. 63. f. 5, 6. *Harz.* t. 59. *Krombh.* t. 66. f. 14, 15, and 16, 17? *Batt.* t. 16. *C.*

**35. *R. decolorans* Fr.**—Pileus 7.5 cent. (3 in.) broad, unicolorous, at first orange-red, then light yellow and becoming pale, fleshy, spherical then expanded and depressed, remarkably regular, viscid when moist, thin and at length striate at the margin; flesh *white, but becoming somewhat cinereous* when broken, and more or less *variegated with black spots* when old. **Stem** elongated (10 cent., 4 in.) cylindrical, solid, but spongy within, often *rugoso-striate, white then becoming cinereous* especially within. **Gills** adnexed, often in pairs, thin, crowded, fragile, white then yellowish.

*Taste mild.* Colours changeable according to a fixed rule, but not variable. The gills are not ochraceous-pulverulent as in *R. integra*, nor shining and pure yellow as in *R. aurata*, &c.

In woods. Uncommon. Aug.-Sept.

Spores pallid ochraceous Fr. Name—*de*, and *coloro*, to colour. Changing colour. *Fr. Monogr.* ii. p. 199. *Hym. Eur.* p. 451. *Berk. Out.* p. 214. *C. Hbk.* n. 633. *S. Mycol. Scot.* n. 607.

**36. *R. aurata* Fr.**—Pileus 5-7.5 cent. (2-3 in.) broad, varying *lemon-yellow, orange and red*, disc darker, fleshy, *rigid*, brittle however, hemispherical then plane, disc not depressed, pellicle thin, adnate, viscid in wet weather, *margin even*, and slightly striate only when old, but sometimes wrinkled; *flesh lemon-yellow* under the pellicle, white below. **Stem** 5-7.5 cent. (2-3 in.) long, solid, *firm*, but spongy within, cylindrical, obsoletely striate, white or lemon-yellow. **Gills** rounded free, connected by veins, broad, equal, shining, never pulverulent, whitish inclining to light yellow, but vivid *lemon-yellow at the edge*.

Taste at first mild, then slightly acrid, odour pleasant. It is among the most beautiful and most easily recognised species by the colours alike of the pileus and of the flesh and gills.

In woods. Uncommon. Aug.-Sept.

Russula.

The bright yellow edge of the gills is a very distinctive mark. Name—*aurum*, gold. From its colour. *Fr. Monogr.* ii. p. 200. *Hym. Eur.* p. 452. *Berk. Out.* p. 213. *C. Hbk.* n. 634. *S. Mycol. Scot.* n. 608. *Ag. With.*—*Krampf.* t. 5. *Schæff.* t. 15. f. 1-3. *Krombh.* t. 66. f. 8-11.

**37. *R. nitida* Fr.**—**Pileus** about 5 cent. (2 in.) broad, commonly bay-brown-purplish, disc darker, shining, fleshy, stiff, convex then plane or slightly depressed, covered with a pellicle which is viscid in wet weather, margin striate, somewhat tubercular; flesh white. **Stem** 5-7.5 cent. (2-3 in.) long, 12 mm. ( $\frac{1}{2}$  in.) thick, spongy-stuffed, soft, somewhat equal or attenuated downwards, even, white inclining to pale. **Gills** adnexed, separating, crowded, pallid then *bright sulphur-yellow, not pulverulent*.

Taste mild, at length nauseous; said to be *somewhat fœtid*. Distinguished at first sight from all others by the *gills being bright lemon-yellow, shining, and never ochraceous-pulverulent*.

In woods. Uncommon. Aug.-Sept.

Name—*nitidus*, shining. *Fr. Monogr.* ii. p. 201. *Hym. Eur.* p. 452. *Berk. Out.* p. 214. t. 13. f. 7. *C. Hbk.* n. 636. *S. Mycol. Scot.* n. 609. *Ag. Pers.*—*Krombh.* t. 66. f. 1-3.—*Schæff.* t. 254.

\*\*\* *Gills and spores ochraceous.*

**38. *R. alutacea* Fr.**—**Pileus** 5-10 cent. (2-4 in.) broad, commonly bright blood-colour or *red*, even black-purple, but becoming pale, especially at the disc, fleshy, campanulate then convex, flattened and somewhat umbilicate, even, *with a remarkably viscous pellicle, margin thin, at length striate, tubercular; flesh snow-white*. **Stem** 5 cent. (2 in.) long, solid, stout, equal, even, white, most frequently *variegated-reddish*, even purple. **Gills** at first free, *thick, very broad*, connected by veins, all equal, somewhat distant, at first pallid light yellow, then bright ochraceous, not pulverulent.

Handsome. Taste mild, pleasant; at length soft and very fragile. *Changeable in colour*; in shady beech woods a form occurs with the *pileus æruginous*. In one form the stem is wholly dark blood-colour, and the gills are rounded behind and remote from the stem. It is distinguished from *R. integra* by its gills not being pulverulent.

In mixed woods. Common.

Edible and very good. Name—*aluta*, tanned leather. From the yellowish gills. *Fr. Monogr.* ii. p. 202. *Hym. Eur.* p. 453. *Berk. Out.* p. 214. t. 13. f. 8. *C. Hbk.* n. 637. *S. Mycol. Scot.* n. 610. *Price f.* 36. *Ag. Vittad.* t. 34. *Krombh.* t. 64. f. 1-3. *Barla t.* 14. f. 1-3.

**39. *R. lutea* Fr.**—**Pileus** 2.5-5 cent. (1-2 in.) broad, *yellow*,

Russula.

at length becoming pale, and occasionally wholly white, thinly fleshy, soon convexo-plane or plano-depressed, viscous when moist, *even* or when old obsoletely striate *at the margin*; *flesh white*. **Stem** 4 cent. (1½ in.) long, 6–8 mm. (3–4 lin.) thick, stuffed then *hollow*, soft, fragile, equal, even, white, never reddish. **Gills** somewhat free, connected by veins, *crowded*, *narrow*, all equal, ochraceous-egg-yellow.

Always small, very regular; taste mild. *When young the pileus is always of a beautiful yellow.*

In woods. Uncommon. July–Oct.

Spores yellow, echinulate, 8 mk. *W.G.S.*; globose, rough, 6–7 mk. *C.B.P.* Name—*luteus*, yellow. *Fr. Monogr.* ii. p. 203. *Hym. Eur.* p. 454. *Berk. Out.* p. 214. *C. Hbk.* n. 638. *S. Mycol. Scot.* n. 611. *Ag. Huds.*

**40. R. nauseosa** Fr.—**Pileus** variable in colour, typically *purplish at the disc*, then livid, but becoming pale and often whitish, laxly fleshy, thin, at first plano-gibbous, then depressed, viscid in wet weather, *sulcate* and somewhat tubercular at the somewhat membranaceous *margin*; flesh soft, white. **Stem** short, about 2.5 cent. (1 in.) long, 8 mm. (4 lin.) thick, spongy-stuffed, slightly striate, white. **Gills** adnexed, ventricose, *somewhat distant*, here and there with a few shorter ones intermixed, light yellow then dingy ochraceous.

The taste is mild, but also nauseous, as the odour often is. The habit is that of *R. nitida*, of the same colour of pileus, but differing in the colour of the gills.

In woods, chiefly pine. Coed Coch, &c. Oct.

Name—*nauseosus*, nauseous. *Fr. Monogr.* ii. p. 204. *Hym. Eur.* p. 454. *B. & Br.* n. 1283. *Ag. Schæff. t.* 16. f. 4.

**41. R. vitellina** Fr.—**Pileus** 2.5 cent. (1 in.) broad, *unicolorous*, light yellow then wholly pallid, somewhat membranaceous, at length *tuberculoso-striate*, somewhat dry, disc very small, slightly fleshy. **Stem** thin, scarcely exceeding 2.5 cent. (1 in.) long, 4 mm. (2 lin.) thick, equal. **Gills** separating-free, equal, *distant*, rather thick, connected by veins, saffron-yellow.

Pretty, very fragile, strong-smelling, mild.

In fir and mixed woods. Uncommon. Aug.–Sept.

Name—*vitellus*, yolk of egg. Egg-yellow. *Fr. Hym. Eur.* p. 454. *Berk. Out.* p. 215. *C. Hbk.* n. 639. *S. Mycol. Scot.* n. 612. *Ag. Pers.*—*Batsch f.* 72.

**42. R. chamæleontina** Fr.—**Pileus** 2.5–5 cent. (1–2 in.) broad, thinly fleshy, soon flattened, sometimes oblique with a thin,

separable, viscid pellicle, which is at first flesh-colour, then *Russula*. presently changing colour, becoming yellow at the disc and at length wholly yellow, margin even, then slightly striate. **Stem** as much as 7.5 cent. (3 in.) long, but thin, somewhat hollow, slightly striate, white. **Gills** more or less adnexed, *thin, crowded*, equal, narrow, somewhat forked, light-yellow-ochraceous.

Mild, inodorous, very fragile. Pileus *rosy-blood-red, purplish lilac*, &c. Sometimes even at the first yellowish at the disc.

In woods. Rare. Sept.

Name—from its changing colours like the *chamæleon*. *Fr. Monogr.* ii. p. 204. *Hym. Eur.* p. 455. *B. & Br. n.* 1014. *C. Hbk. n.* 640. *S. Mycol. Scot. n.* 613.

GENUS XI.—**Cantharellus** Adans.—(κάνθαρος, a vase or cup.) *Cantharellus*.  
Fr. Syst. Myc. i. p. 316.

Hymenophore continuous with the stem, descending unchanged into the trama. Gills thick, fleshy-waxy, *fold-like*, somewhat branched, *obtuse at the edge*. Spores white. *Fleshy, membranaceous, putrescent fungi, without a veil*. *Fr. Hym. Eur.* p. 455.

The distinguishing character of the genus consists in the fold-like gills. The *Mesopodes* commonly grow on the ground; the rest commonly on mosses, very rarely on wood. Intermediate between *Agaricus* and *Craterellus*. Some are edible, others reckoned poisonous.



L1. *Cantharellus cibarius*. One-third natural size.

I. MESOPUS (μέσος, middle; πούς, a foot). Pileus entire. Stem central.

\* *Pileus and solid stem fleshy.*

\*\* *Pileus somewhat membranaceous, stem tubular and polished.*

II. MERISMA (μερίζω, to divide). Stems very numerous, united into an elongated column or branched.

III. PLEUROPUS (πλευρόν, the side; πούς, a foot). Dimidiate, stem exactly lateral.

IV. RESUPINATUS (*resupinatus*, lying on the back). Pileus entire, at first cup-shaped, adfixed by the vertex, then somewhat reflexed.

\* *Bryophilus* (moss-loving).

\*\* *Lignatiles* (growing on wood), doubtful species.

Canthar-  
ellus.

## I.—MESOPUS.

\* *Pileus and solid stem fleshy.*

1. **C. cibarius** Fr. *Egg-yellow*.—**Pileus** *truly fleshy*, firm, at first convex, then plane or somewhat depressed, repand, *always smooth*. **Stem** solid, thickened upwards. **Gills** decurrent, *thick, distant*, branched, narrow, fold-like.

Taste pleasant. Growing in troops. The stem passes into the pileus which is hence turbinate. The gills are *more swollen than those of the other species*. It varies wholly white.

In woods, especially beech. Common. July–Dec.

Berkeley records a var. white, here and there tinged with pink. The smell, especially when the plant is old, is that of ripe apricot. Edible, delicious. It must have four hours' slow cooking. Spores sphaeroid-ellipsoid,  $8-9 \times 5-6$  mk. *K.*;  $6 \times 8$  mk. *W.G.S.*; pruiniform, nucleate, 11 mk. *Q.* Name—*cibaria*, food. *Fr. Monogr.* ii. p. 206. *Hym. Eur.* p. 455. *Sv. atl. Sv. t.* 7. *Berk. Out.* p. 215. *C. Hbk. n.* 641. *S. Mycol. Scot. n.* 614. *Harz. t.* 18. *Grev. t.* 258. *Badh. i. t.* 9. f. 2; ii. t. 8. f. 1. *Price f.* 94. *Krombh. t.* 45. f. 1-11. *Vittad. t.* 25. f. 1. *Barla t.* 28. *Ag. Fl. Dan. t.* 264. *Sow. t.* 46.

2. **C. Friesii** Quel.—**Pileus** somewhat orange, fleshy, thin, convex then depressed, *villous*. **Stem** solid, slender, *villous*, *at the base white, attenuated*. **Gills** narrow, *fold-like*, branched, yellow.

Exactly intermediate between *C. cibarius* and *C. aurantiacus*, but smaller than either.

In woods. Sydenham Hill.

Name—after Elias Fries. *Quell. Jur. p.* 191. t. 23. f. 2. *Fr. Hym. Eur.* p. 455. *B. & Br. n.* 1782. Apparently *Krombh. t.* 46. f. 3-6 and perhaps *Paul. t.* 51. f. 3-4.

3. **C. aurantiacus** Fr.—**Pileus** 5-7.5 cent. (2-3 in.) broad, orange-ochraceous, fleshy, *soft*, depressed, often excentric and undulated, *somewhat tomentose*, involute at the margin. **Stem** 5 cent. (2 in.) long, stuffed at length *hollow*, somewhat incurved and unequal, ochraceous. **Gills** decurrent, tense and straight, repeatedly dichotomous and *crowded*, often crisped at the base, deep orange.

Taste unpleasant. The pileus is not firm, and the stem is not so much diffused into the pileus as in *C. cibarius*. The stem, especially in marshy places, becomes at length black. *Bull. t.* 505. It varies with light yellow pileus and white gills, and is sometimes wholly white.

In woods, especially pine. Common. Aug.–Nov.

Reckoned poisonous. Spores ellipsoid or sphaeroid-ellipsoid,  $5-6 \times 3$  mk. *K.* Name—*aurantiacus*, orange-yellow. *Fr. Monogr.* ii. p. 206. *Hym. Eur. p.* 455. *Sv. atl. Sv. t.* 79. *Berk. Out. p.* 215. t. 14. f. 1. *C. Hbk. n.* 642. *S. Mycol. Scot. n.* 615. *Ag. Wulf. Jacqu. Coll. ii. t.* 14. f. 3. *Batsch f.* 37. *Sow. t.* 413.



**4. C. Brownii** B. & Br. Ochraceous-white or cream-coloured. *Cantharellus*.  
—**Pileus** about 12 mm. ( $\frac{1}{2}$  in.) broad, thin, convex, somewhat umbonate, obscurely silky. **Stem** 4-5 cent. ( $1\frac{1}{2}$ -2 in.) long, stuffed, slender, tough, nearly equal, somewhat furfuraceous. **Gills** (folds) obtusely decurrent, rather distant, *linear*, extremely narrow, sometimes forked, interstices smooth.

Hymenium nearly white. Stem furnished with a little white fibrillose mycelium at the base, which sometimes forms a small earthy ball. A very distinct species.

Among grass. Hitchin.

Name—after J. Brown. *Berk. Out. p. 215. C. Hbk. n. 643. Fr. Hym. Eur. p. 456.*

**5. C. carbonarius** A. & S.—**Pileus** *date-brown then black*, somewhat fleshy, *striato-squamulose*, umbilicate. **Stem** paler. **Gills** tense and straight, white.

Rooted, fasciculate. The following seem to be the same: B. *C. anthracophilus*, solitary, *black, tough*, pileus fleshy-membranaceous, naked, depressed; stem stuffed, slender, smooth, *slightly rooted*; gills distant, dichotomous. *Leveill. Ann. Sc. Nat. 1841, p. 236. t. 14. f. 2.*—C. *C. radicosus*, slender, pileus 18 mm.-2.5 cent. ( $\frac{3}{4}$ -1 in.), deeply umbilicate, *floccose, black*; stem *rooting*, pallid; hymenium white, gills narrow. Two or three pilei often grow from the same obconical root, which is white and spongy.

On charcoal heaps, &c. Rare. Sept.-Nov.

Spores 9x6 mk. *W.G.S.* Name—*carbo*, charcoal. From its habitat. *A. & S. Fr. Hym. Eur. p. 456. C. radicosus B. & Br. n. 1134. C. Hbk. n. 647. Saund. & Sm. t. 1.*

**6. C. umbonatus** Fr.—**Pileus** 2.5 cent. (1 in.) and more broad, *cinereous-blackish*, slightly fleshy, convex when young, *umbonate, at length depressed*, even, dry, *flocculoso-silky* on the surface, shining brightly especially under a lens; flesh soft, white, often becoming red when wounded. **Stem** 7.5 cent. (3 in.) long, about 8 mm. (4 lin.) thick, *stuffed*, equal, elastic, villous at the base, *cinereous*, but paler than the pileus. **Gills** decurrent, thin, tense and straight, *crowded*, repeatedly dichotomous, *shining white*.

Odour and taste scarcely notable. Gregarious. Among the taller mosses the stem is longer. Often overlooked from its habit being that of an Agaric. It varies with the pileus squamulose and blackish.

In woods. Rare. April-Aug.

Name—*umbo*, the boss of a shield. Umbonate. *Fr. Monogr. ii. p. 207. Hym. Eur. p. 457. Berk. Out. p. 216. C. Hbk. n. 644. S. Mycol. Scot. n. 616. Hoffm. Ic. t. 22. f. 2. Ag. Wulf. Jacqu. Coll. 2. t. 16. f. 1.*

**7. C. albidus** Fr.—**Pileus** 1-2.5 cent. ( $\frac{1}{2}$ -1 in.) broad, *whitish*,

Canthar-  
ellus.

inclining to yellowish or rufescent, fleshy, *very thin*, somewhat infundibuliform, irregular, somewhat repand and torn, *smooth*, somewhat zoned. **Stem** 2.5–4 cent. (1–1½ in.) long, 4 mm. (2 lin.) thick, *solid*, tough, equal or at length compressed, smooth, white or rarely yellowish. **Gills** decurrent, thin, somewhat crowded, repeatedly dichotomous, white.

Tough but not coriaceous. It approaches near to the *Clitocybæ* among Agarics, clearly showing the affinity of these with the *Cantharelli* (*C. cibarius*, *aurantiacus*, *umbonatus*, and *albidus*).

In mossy places. Coed Coch. Sept.–Oct.

Name—*albus*, white. Whitish. *Fr. Monogr.* ii. p. 207. *Hym. Eur.* p. 457. *B. & Br. n.* 1421. *Fl. Dan. t.* 1293. f. 1.

**8. C. Houghtoni** Phill.—**Pileus** 2.5 cent. (1 in.) or more broad, dirty white with a tinge of flesh-colour, thin, convex, umbilicate, smooth. **Stem** 5 cent. (2 in.) long, 2 mm. (1 lin.) thick, stuffed, slender, thickened at the apex, at first delicately fibrillose, rooting at the *base* which is more or less *cottony*. **Gills** somewhat decurrent, narrow, scarcely forked, pallid flesh-colour.

Sometimes 5 cent. (2 in.) broad. Allied to *C. albidus*.

On the ground. Hereford.

Name—after Rev. William Houghton. *Phillips*.—*B. & Br. n.* 1565.

**\*\* *Pileus somewhat membranaceous, stem tubular and polished.***

**9. C. tubæformis** Fr.—**Pileus** fuscous when moist, becoming pale when dry, fleshy-membranaceous, infundibuliform, repand and lobed, flocculose. **Stem** hollow, at length compressed and lacunose, smooth, *orange-tawny*. **Gills** thick, distant, branched with many clefts, yellow or fuliginous, *naked*.

It differs widely from all the preceding species in the somewhat membranaceous, infundibuliform, flocculose pileus, and in the tubular stem. Intermediate between *C. infundibuliformis*, to which it is like in the hymenium, and *Craterellus lutescens*, which is more like it in the entire habit.

**Var. lutescens**, pileus convexo-umbilicate, *rather even*, somewhat regular; stem more equal, attenuated upwards; gills less divided. *Bull. t.* 473. f. 3.

In woods. Common. Aug.–Oct.

Spores 8 × 4 mk. *W.G.S.* Name—*tuba*, a trumpet; *forma*, form. Trumpet-shaped. *Fr. Monogr.* ii. p. 208. *Hym. Eur.* p. 457. *Berk. Out.* p. 216. *C. Hbk. n.* 645. *S. Mycol. Scot. n.* 617. *Fl. Dan. t.* 2080. f. 1. *Merulius Pers. Ic. descr. t.* 6. f. 1. *Dittm. Sturm. t.* 30. *Batt. t.* 23. f. 1.

**10. C. infundibuliformis** Fr.—**Pileus** 2.5–5 cent. (1–2 in.) broad, rarely more, yellowish-cinereous or fuliginous when

moist, becoming pale when dry, somewhat membranaceous, umbilicate then infundibuliform, here and there pervious to the base, *floccoso-wrinkled* on the surface, at length undulated at the margin. **Stem** 5-7.5 cent. (2-3 in.) long, about 4 mm. (2 lin.) thick, *fistulose*, somewhat thickened at the base, even, smooth, always *light yellow*. **Gills** decurrent, thick, distant, dichotomous, tense and straight, sometimes *light yellowish*, sometimes cinereous, especially when old and *then pruinete*.

Gregarious, somewhat caespitose. Often not easily distinguished from *C. tubaeformis*. It does not approach *Crat. lutescens*. The colours are less intense, *moderately persistent when dried*.

In woods. Common. July-Oct.

Spores sphaeroid-ovoid,  $9-11 \times 7-8$  mk. *K.* Name—*infundibulum*, a funnel; *forma*, form. Funnel-shaped. *Fr. Monogr.* ii. p. 208. *Hym. Eur.* p. 458. *Berk. Out.* p. 216. *C. Hbk.* n. 646. *S. Mycol. Scot.* n. 618. *Merulius Scop.*—*Sow. t.* 47. *Krombh. t.* 46. f. 7-9. *Fl. Dan. t.* 1617.

**11. C. cinereus** Fr.—**Pileus** 2.5-5 cent. (1-2 in.) and more broad, hoary-fuliginous, somewhat membranaceous, infundibuliform, *pervious as far as the base of the stem, villososquamous*. **Stem** 5-7.5 cent. (2-3 in.) long, 4-8 mm. (2-4 lin.) thick, hollow, manifestly attenuated downwards, smooth, *fuliginous-blackish*. **Gills** decurrent, thick, very distant, connected by veins but not much branched, cinereous.

As *C. tubaeformis* resembles *Crat. lutescens* this entirely resembles *Crat. cornucopioides*, but is very easily distinguished from it by the hymenium being in the form of gills.

In woods. Rare. Halifax, &c.

Spores  $15 \times 8$  mk. *B. & Br.* Name—*cinis*, ashes. From the cinereous gills. *Fr. Monogr.* ii. p. 209. *Hym. Eur.* p. 458. *Berk. Out.* p. 216. *B. & Br. n.* 1016\*, 1139\*. *C. Hbk.* n. 648. *Krombh. t.* 45. f. 12. *Merulius Pers.*—*Bull. t.* 465. f. 2.

**12. C. cupulatus** Fr.—**Pileus** 12 mm. ( $\frac{1}{2}$  in.) broad, pallid fuscous when damp, becoming pale, somewhat rufescent, when dry, membranaceous, *plano-infundibuliform* (exactly cup-shaped), repand, not zoned, when moist smooth, *striate at the margin*, when dry even, flocculose. **Stem** scarcely 2.5 cent. (1 in.) long, 1 m. ( $\frac{1}{2}$  lin.) thick, *stuffed*, equal, tense and straight, tough, smooth, paler than the pileus. **Gills** decurrent, very distant, *branched with many clefts*, with some *simple ones intermixed*, broad, but obtuse at the edge, grey.

The branches on the gills ramify in a fasciculate manner. It must not be confounded with *Xerotus degener*.

In waste places. Rare. Oct.

Canthar-  
ellus.

Name—*cupa*, a cup. Shaped like a *little cup*. *Fr. Monogr.* ii. p. 209. *Hym. Eur.* p. 458. *Merulius Pers. Syn. t. 5. f. 2.* *Ag. helvelloides Bull. t. 601. f. 3.* *Berk. Out. p. 132.* *C. Hbk. n. 226.*

**13. C. Stevensoni** B. & Br.—**Pileus** about 4 mm. (2 lin.) broad, pallid, orbicular, *umbilicate*, smooth, margin inflexed. **Stem** 6 mm. ( $\frac{1}{4}$  in.) long, 1 m. ( $\frac{1}{2}$  lin.) thick, cylindrical, delicately pulverulent, white then darker. **Gills** decurrent, pallid, becoming fuscous in front.

Stem with a little white mycelium at the base. Very near to *C. cupulatus*; but that is very strongly umbonate when young, and the *umbo* is always visible at the bottom of the umbilicus; the habit, moreover, is different.

On rotten stump among moss. Glamis, 1874. March, April.

Name—after Rev. John Stevenson. *B. & Br. n. 1422.* *S. Mycol. Scot. n. 619.*

## II.—MERISMA.

No British species.

## III.—PLEUROPUS.

**14. C. muscigenus** Fr.—**Pileus** fuscous when moist, cinereous-whitish when dry, somewhat zoned, membranaceous, tough, spathulate, smooth, rather plane, slightly undulated when full grown. **Stem** short, 2–4 mm. (1–2 lin.), round, exactly lateral, *villous at the base*, horizontal as well as the pileus with which it is continuous. **Gills** slightly swollen, diverging from the apex of the stem, *distant, branched*, not anastomosing, of the same colour as the pileus.

The whole plant from the base of the stem to the apex of the pileus scarcely reaching 2.5 cent. (1 in.), commonly smaller.

On mosses. Uncommon. June.

Name—*muscus*, moss; *gigno*, to bear. Growing on mosses. *Fr. Monogr.* ii. p. 210. *Hym. Eur.* p. 460. *Berk. Out. p. 217.* *C. Hbk. n. 649.* *S. Mycol. Scot. n. 620.* *Ag. Bull. t. 288, 498. f. 2.* *Merulius Nees Syst. f. 236.*

## IV.—RESUPINATI.

\* *Bryophili* (moss-loving).

**15. C. retirugus** Fr.—**Pileus** 6–10 mm. (3–5 lin.) broad, whitish-cinereous above, darker, cinereous-fuliginous, beneath, membranaceous, expanded, repando-lobed, very tender, somewhat round, at first entire at the margin then variously split. **Gills** radiating from the centre, *very tender, reticulated*,

Easily distinguished from *C. lobatus* by the *thinner, reticulated gills*; *Cantharellus*. without the rudiment of a stem, *adfixed* to mosses *by fibrils at the vertex*. It approaches the *Cyphellæ* in many respects.

On mosses. Uncommon. May.

Name—*rete*, a net; *ruga*, a wrinkle. From the reticulated gills. *Fr. Monogr. ii. p. 212. Hym. Eur. p. 460. Berk. Out. p. 217. t. 14. f. 2. Helvella Bull. i. 498. f. 1. Sow. t. 348.*

**16. C. lobatus** Fr.—**Pileus** scarcely 2.5 cent. (1 in.), *dark fuscous* when in full vigour, becoming pale when dry, *sessile*, horizontal, membranaceous, somewhat round, at first adfixed, entire, then somewhat lateral and lobed. **Gills** *fold-like*, somewhat distant, *distinct, branched*, of the same colour as the pileus.

On mosses in bogs. Uncommon. Autumn.

When old often very much lobed and crisped, and then the wrinkles, especially at the base, are reticulated, though towards the margin they continue distinct. *M. J. B.* Spores ovoid-pruniform, 10 mk. *Q.* Name—*lobatus*, lobed. *Fr. Monogr. ii. p. 211. Hym. Eur. p. 461. Berk. Out. p. 217. C. Hbk. n. 650. S. Mycol. Scot. n. 622. Fl. Dan. t. 1077. Bolt. t. 177.*

**\*\* Lignatiles** (growing on wood).

No British species.

GENUS XII.—**Arrhenia** Fr. *Summ. Veg. Scand. p. 312.*

*Arrhenia.*

Membranaceous, very tender, hymenium inferior and striate with few, delicate, slightly raised, simple, straight veins, which take the place of gills. Spores pallid. *Minute, moderately persistent, variable in form. Striæ not decurrent.* Compare *Cantharellus Brownii*. *Fr. Hym. Eur. p. 461.*

No British species.

GENUS XIII.—**Nyctälis** (*νύξ*, night. From inhabiting dark places.) *Fr. Syst. Orb. Veg. p. 203.*

*Nyctalis.*

Hymenophore continuous with the stem. Gills fleshy, thick, juicy, obtuse at the edge, not decurrent on the stem nor fold-like. Veil (in species which have been fully observed) floccoso-pruinose. *Fleshy fungi, not reviving, of uncertain and irregular occurrence, differing in many respects from one another and from the rest of the Agaricini.* *Fr. Hym. Eur. p. 462.*

I. **SPELEÆ** (*spelæum*, a cave). Gills crowded, somewhat coalescent. In


Nyctalis.

hollow places. There is scarcely any doubt that these species are forms of others, changed in appearance and disguised owing to their place of growth.

II. PARASITÆ (*parasita*, a parasite). Gills distinct, distant. On putrid fungi.

## I.—SPELEÆ.

1. *N. caliginosa* Smith.—**Pileus** white when dry, marked with colours (as in *Ag. butyraceus*) when wet, very fleshy, floccoso-pruinose when dry, margin involute, slightly exceeding the gills. **Stem** solid, floccoso-pruinose, base naked. **Gills** decurrent, thick, branched.



Odour and taste rank and disagreeable like *Polyporus squamosus*. Closely allied to *N. parasitica*, but at once distinguished by its truly decurrent gills and other characters.

Amongst earth and dead leaves in dense dark wood. Bishop's Wood, Highgate.

LII. *Nyctalis parasitica*. One-half natural size.

Name—*caligo*, darkness. From its habitat. *Worth. Smith Journ. Bot.* 1873, p. 337.

## II.—PARASITÆ.

2. *N. asterophora* Fr.—**Pileus** white when young, then *fawn-colour*, fleshy, conical then hemispherical, often papillate when young, then splitting open, *floccoso-pulverulent*. **Stem** scarcely exceeding 12 mm. ( $\frac{1}{2}$  in.) long, 1–2 mm. ( $\frac{1}{2}$ –1 lin.) thick, stuffed, equal, but often twisted, at first whitish and pruinose, then becoming fuscous. **Gills** adnate, distant, thick, tense and straight, somewhat forked, dingy.

Very abnormal. Gregarious. The dust covering the pileus represents stellate spores under the microscope. B. even underneath and without gills.

On dead *Russula nigricans*, &c. Frequent. Sept.–Nov.

Spores  $2 \times 3$  mk. *W.G.S.* Name—*aster*, a star; *fero*, to bear. From the stellate bodies in the dust on the pileus. *Fr. Monogr.* ii. p. 214. *Hym. Eur.* p. 463. *Berk. Out.* p. 217. *C. Hbk. n.* 652. *S. Mycol. Scot. n.* 623. *Ag. Bull. t.* 516. f. 1. *Asterophora Lycop. Dittm. Sturm t.* 26.

3. *N. parasitica* Fr.—**Pileus** 1–2.5 cent. ( $\frac{1}{2}$ –1 in.) broad, whitish-fuscous, then becoming pale, whitish, slightly fleshy, when young conico-campanulate then convex, soon plane and obtuse

or obsoletely umbonate, unequal, with a *persistent, pruinose, grey* Nyctalis. *pellicle*; flesh dingy. **Stem** 2.5–5 cent. (1–2 in.) long, 3 mm. (1½ lin.) thick, delicately fistulose, equal or slightly attenuated upwards, straight or curved from oblique position, pubescent, white. Veil manifest, fibrillose. **Gills** adnate, thick, distant, with alternate shorter ones intermixed, becoming fuscous, at length contorted and anastomosing.

Gregarious, even cæspitose. The pileus is often irregular (angular, repand and rimose).

On dead *Russulæ*, *adusta* and *fætens*. Common. Aug.–Nov.

Odour like *Polyporus squamosus*. *M. f. B.* Name—from its *parasitic* growth. *Fr. Monogr.* ii. p. 214. *Hym. Eur.* p. 464. *Berk. Out.* p. 218. *t.* 19. *f.* 2. *C. Hbk. n.* 653. *S. Mycol. Scot. n.* 624. *Ag. Bull. t.* 574. *f.* 2. *Sow. t.* 343.


#### GENUS XIV.—*Marasmius* (*μαρasmus*, to wither or shrivel).

Marasmius.

Fr. Gen. Hymen. Epicr. p. 372.

Tough, arid fungi; shrivelling (and not putrescent), reviving when wet. *Hymenophore continuous with the stem, but heterogeneous*, descending into the trama; veil none with one exception. Stem cartilaginous or horny. Gills pliant, somewhat distant, edge acute and quite entire.

This begins a new series of pliant and arid Agaricini. They are commonly epiphytal. Like other plants they may be dried and preserved in herbaria. *Marasmius* is a very natural genus, although it be principally distinguished by biological marks. Many of the species have a strong smell and peculiar taste (all the Agaricini with the smell of garlic seem to belong to this genus, but no one of the species has the smell of new meal). *Fr. Hym. Eur.* p. 464.



LIII. *Marasmius oreades*. One-half natural size.

*Marasmius* is closely allied to *Collybia*. Their biological character of shrivelling, and reviving with moisture is of the utmost importance in the determination of the true *Marasmii*. Fries refers (*Monogr.* ii. p. 215) to two species which hold a doubtful

Marasmius. place between the genuine *Agarics* and the *Marasmii*—viz., *Ag. (Collybia) stipitarius* and *Marasmius oreades*. One species (not British) *M. subannulatus* has a veil.

The species are mostly small, thin, and commonly epiphytal; those which are found on the ground are generally on putrid leaves or roots of grasses. They are too thin to be of much use for food, but some are esteemed for flavouring.

I. COLLYBIA. Fr. Syst. Orb. Veg. i. p. 73. Pileus fleshy-pliant, at length somewhat coriaceous, sulcate or wrinkled, margin at first involute. Stem somewhat cartilaginous, mycelium floccose in a very few not manifest.

A. Scortei (*scorteus*, coriaceous). Stem solid, or stuffed with a pith then hollow, internally fibrous, externally with villous down (pruina, 'Monogr.') that is easily wiped off covering the cartilaginous cuticle. Gills separating-free.

\* Stem woolly or strigose at the base.

\*\* Stem naked at the base, often composed of twisted fibres.

B. Tergini (*tergum*, hide, skin, leather). Stem rooting, definitely tubular, not fibrous, manifestly cartilaginous. Gills separating-free. Pileus thinner than in the former section, hygrophanous, sometimes even, sometimes striate at the margin.

\* Stem woolly downwards, smooth upwards.

\*\* Stem (at least when dry) everywhere pruinato-velvety.

C. Calopodes (*καλός*, beautiful; *πούς*, a foot). Stem curt, rootless, inserted, often furnished with a floccose tubercle at the base. Pileus convexo-involute then plane and depressed; in this state the gills, which are typically adnate, are somewhat decurrent. On stipules, branches, &c., gregarious, arid.

\* Stem very smooth and shining upwards, with a simple base.

\*\* Stem velvety or pruinose, with a somewhat tubercular base.

II. MYCENA. Stem horny, fistulose, but here and there with a pith, tough, arid, mycelium rhizomorphoid corticate and not floccose. Pileus somewhat membranaceous, campanulate then expanded, margin at the first straight and adpressed to the stem.

A. Chordales (*chorda*, a gut. *M. chordalis*). Stem rigid, rooting, or adnate by a dilated base. Pileus campanulate or convex. Type manifestly that of *Mycena*.

B. Rotulæ (*rotula*, a little wheel). Stem filiform, flaccid, inserted at the base. Pileus soon rather plane or umbilicate. Growing on leaves, declining from the common type of *Mycena*.

\* Stem very smooth, shining.

\*\* Stem velvety or pilose.

III. APUS (α, and *πούς*, a foot. Stemless). Pileus sessile, resupinate.

## I. — COLLYBIA.

### A. SCORTEI.

\* Stem woolly or strigose at the base.

1. *M. urens* Fr.—Pileus 5-7.5 cent. (2-3 in.) broad, unicolorous, pale yellowish becoming pale, slightly fleshy, moderately compact at the disc, even, but here and there squamulose or rimoso-



rivulose when dry, smooth, the thin margin involute. **Stem** 5-7.5 cent. (2-3 in.) long, 6 mm. (3 lin.) thick, *solid*, composed of crisp tough fibres, rigid, equal, sometimes however ventricose, 12 mm. ( $\frac{1}{2}$  in.) thick, *everywhere clothed with white flocci, pale*, white-villous at the base. **Gills** free, united behind, *at length remote* from the stem, *distant*, tough, at first pale-wood colour, *then brown*.

Gregarious, somewhat caespitose. *Taste very stinging*. The stem is not strigosely sheathed at the base.

In mixed woods. Frequent. June-Sept.

A curious form has occurred with the pileus turning very dark when full grown. *B. & Br.* Poisonous. Worthington Smith has tested it by accident. It produced headache, swimming of brain, burning in throat and stomach, followed by severe purging and vomiting. Spores  $3 \times 4$  mk. *W.G.S.* Name—*uro*, to burn. From its *burning* taste. *Fr. Monogr.* ii. p. 216. *Hym. Eur.* p. 465. *Berk. Out.* p. 218. t. 14. f. 3. *B. & Br.* n. 1889. *C. Hbk.* n. 654. *S. Mycol. Scot.* n. 625. *Gonn. & Rab.* t. 8. f. 1. *Ag. Bull.* t. 528. f. 1. *Fl. Dan.* t. 2018. f. 1.

**2. M. peronatus** Fr.—**Pileus** 2.5-5 cent. (1-2 in.) and more broad, light yellowish or pallid brick-rufescent, then becoming pale, *wood-colour* or tan, at first fleshy-pliant, then *coriaceo-membranaceous*, convex then plane, obtuse, flaccid, slightly wrinkled, even at the disc, *at length lacunose, striate at the margin*; flesh white. **Stem** 5-7.5 cent. (2-3 in.) long, 2-4 mm. (1-2 lin.) thick, stuffed, fibrous, tough, attenuated upwards, *at length hollow* and compressed, *furnished with a bark*, light yellow then pallid, *cuticle villous* but separating and rufescent when rubbed, somewhat incurved at the base, where it is *clothed with dense, somewhat strigose*, yellowish or white *villous down*. **Gills** *adnexed then separating*, free, moderately *thin*, and *crowded*, when young whitish, *pallid wood-colour*, at length somewhat remote, rufescent.

B. Woolly-sheathed at the base. *Taste acrid* like that of *M. urens*, odour none.

In woods. Common. July-Dec.

Spores pip-shaped,  $7 \times 4$  mk. *W.G.S.* Name—*pero*, a kind of boot; *peronatus*, sheathed. From the strigose sheathing at the base of the stem. *Fr. Monogr.* ii. p. 216. *Hym. Eur.* p. 465. *Berk. Out.* p. 218. t. 14. f. 4. *C. Hbk.* n. 655. *S. Mycol. Scot.* n. 626. *Ag. Bolt.* t. 58. *Sow.* t. 37. *Fl. Dan.* t. 2018. f. 2.

**3. M. porreus** Fr.—**Pileus** 2.5-5 cent. (1-2 in.) broad, *dingy yellowish*, pallid when dry, *coriaceo-membranaceous*, flaccid, convex then flattened, obtuse, even at the disc, *striate* at the margin, opaque; flesh pallid, almost of the same colour as the pileus. **Stem** 7.5 cent. (3 in.) long, stuffed then hollow, tough, juiceless, *pubescent*, somewhat thickened at both ends, *red-fuscous*, paler at the apex, *slightly villous at the base*. **Gills** separating-free,

Marasmius. *distant*, rather thick, tough, broadly linear, at length coriaceous, *light yellowish then becoming pale*.

*Odour of garlic*, but fugacious when the plant is dried.

In woods, especially among oak-leaves. Frequent. Sept.-Nov.

Spores pip-shaped,  $4 \times 6$  mk. *W.G.S.* Name—*porrum*, the leek. From its smell of garlic. *Fr. Monogr.* ii. p. 217. *Hym. Eur.* p. 466. *Berk. Out.* p. 219. *C. Hbk. n.* 656. *S. Mycol. Scot. n.* 627. *Ag. Pers.*—*Sow. t.* 81.

**\*\* Stem naked at the base, &c.**

**4. M. oreades** Fr.—**Pileus** 2.5–5 cent. (1–2 in.) broad, *rufescent then becoming pale, hygrophaneous, whitish when dry, fleshy*, pliant, convex then plane, somewhat umbonate, even, smooth, slightly striate at the margin when moist. **Stem** 5–7.5 cent. (2–3 in.) long, 3 mm. ( $1\frac{1}{2}$  lin.) thick, *solid*, very tough, *equal*, tense and straight, *everywhere clothed with a villous-woven cuticle* which can be rubbed off, pallid; bluntly rooted at the base, naked, not villous or tomentose. **Gills** free, broad, *distant*, the alternate ones shorter, *at first soft*, then firmer, pallid-white.

*Odour* weak, but *pleasant*, stronger when dried, *taste mild*. Commonly growing in circles or rows.

In lawns and pastures. Common. May–Oct.

Edible, delicious when broiled with butter. It may be pickled or dried for future use. Carefully distinguish *M. urens* (poisonous) from it. Spores ellipsoid-sphaeroid,  $6-7 \times 5-6$  mk. *K.* Name—*Ορειάς*, a mountain-nymph. Forming fairy rings. *Fr. Monogr.* ii. p. 218. *Hym. Eur.* p. 467. *Sv. atl. Sv. t.* 31. *Berk. Out.* p. 219. *t.* 14. *f.* 5. *C. Hbk. n.* 657. *S. Mycol. Scot. n.* 628. *Ag. Bolt. t.* 151. *Grev. t.* 323. *Sow. t.* 247. *Price f.* 11. *Badh. i. t.* 8. *f.* 3, ii. *t.* 7. *f.* 4. *Vittad. t.* 10. *f.* 1. *Krombh. t.* 43. *f.* 11–16. *Fl. Bat. t.* 830. *Gonn. & Rab. t.* 8. *f.* 3. *Schaff. t.* 77. *Bull. t.* 144, 528. *f.* 2.

**5. M. scorteus** Fr. *Wholly whitish*.—**Pileus** 6–10 mm. (3–5 lin.) broad, becoming pale when dried, slightly fleshy, convex, scarcely umbonate, even (not striate at the margin), but slightly wrinkled when dried. **Stem** 2.5 cent. (1 in.) long, *obsoletely fistulose*, thin, filiform, *equal*, not rooted, smooth, *delicately pruinose towards the apex*, contorted and becoming fuscous when dry. **Gills** quite free, remarkably broad, rounded behind, *ventricose, white*.

Inodorous, *mild*. Habit that of very small *M. oreades*.

In grass among trees. Moncreiffe. Aug.

Name—*scortum*, hide. Leathery, *wrinkled*. *Fr. Monogr.* ii. p. 219. *Hym. Eur.* p. 468. *B. & Br. n.* 1677. *S. Mycol. Scot. n.* 629. *Batsch t.* 109?

## B. TEGINI.

Marasmius.

\* *Stem woolly downwards, smooth upwards.*

**6. M. varicosus** Fr.—**Pileus** 12 mm. ( $\frac{1}{2}$  in.) or a little more broad, *fuscous-purple, becoming darker when dry*, somewhat membranaceous, pliant, campanulate then convex, plane, umbonate, even. **Stem** 5–7.5 cent. (2–3 in.) long, thin, *fistulose, rubiginous*, among mosses villosa-tomentose beyond the middle, the down rubiginous, blackish when dry, smooth above, *filled with dark blood-coloured juice, which drops when it is broken*. **Gills** separating-free, *very crowded, linear, very narrow*, of the same colour as the pileus, but becoming dark, *umber-fuscous when dry*.

Solitary, inodorous. In places which are not mossy the stem is only 2.5 cent. (1 in.) long, wholly smooth, becoming black. It is very singular in the juicy stem, and quite exceptional in the gills being darker when dry. It cannot be compared with *A. hæmatopus*.

In damp mossy places. Apethorpe. Sept.

Name—*varix*, dilated vein. Full of veins, which yield a *bloody* juice. *Fr. Monogr. ii. p. 220. Hym. Eur. p. 469. Icon. t. 174. f. 1. B. & Br. n. 1788.*

**7. M. fusco-purpureus** Fr.—**Pileus** 1–2.5 cent. ( $\frac{1}{2}$ –1 in.) broad, dark purple when in vigour, becoming pale (tan) when dry, slightly fleshy, at first hemispherical, then plane, obsoletely umbilicate, slightly wrinkled. **Stem** 2.5–7.5 cent. (1–3 in.) long, 2–4 mm. (1–2 lin.) thick, fistulose, smooth, at first pallid, then rufous or *dark purple, sheathed towards the base with strigose, rubiginous down*. **Gills** adnexed in the form of a ring, then free, *distant*, narrow, rufescent.

*Inodorous, juiceless.* There are two forms, one larger, solitary, with the stem (longer) decumbent, another smaller, caespitose, with the stem (shorter) tense and straight.

In woods, on sticks, &c. Frequent. Aug.

Name—*fuscus*, fuscous; *purpureus*, purple. *Fr. Monogr. ii. p. 220. Hym. Eur. p. 469. Berk. Out. p. 219. C. Hbk. n. 658. S. Mycol. Scot. n. 630. Ag. Pers. Ic. descr. t. 4. f. 1–3.*

**8. M. terginus** Fr.—**Pileus** about 2.5 cent. (1 in.) broad, *flesh-colour* when moist, whitish when dry, *shining*, tough, slightly fleshy, convex then plane, obtuse, at length somewhat depressed, margin striate when moist. **Stem** 5–7.5 cent. (2–3 in.) long, 2–4 mm. (1–2 lin.) thick, fistulose, slightly attenuated upwards, longitudinally fissile, smooth, shining, and pallid upwards, and

Marasmius. not somewhat pruinose when dried, *reddish downwards, white-villous at the rooting base*. **Gills** separating-free, *somewhat crowded, narrow, pallid*.

Inodorous and juiceless, always solitary.

Among leaves, beech, &c. Dursley. Nov.

Name—*tergum*, hide, leather. *Fr. Monogr.* ii. p. 221. *Hym. Eur.* p. 469. *Icon. t.* 174. f. 4. *B. & Br. n.* 1361. *C. Hbk. n.* 661. *M. Stephensii Berk. Out. p.* 220.

**\*\* Stem (at least when dry) everywhere pruinato-velvety.**

**9. M. Wynnei B. & Br.**—**Pileus** 2.5–4 cent. (1–1½ in.) broad, *lilac-brown*, tardily changing colour, fleshy, convexo-plane, somewhat umbonate. **Stem** 5 cent. (2 in.) long, 3 mm. (1½ lin.) thick, fistulose, *furfuraceous*, somewhat of the same colour as the pileus. **Gills** adnexed, thick, distant, bright-coloured, beautifully tinged with *lilac*; interstices even.

Inodorous. Gregarious or caespitose. The stem springs from a white mycelium, but is by no means strigose or tawny at the base. Quite distinct from *M. fusco-purpureus*.

Among leaves, twigs, &c. Coed Coch.

Name—after Mrs Lloyd Wynne. *B. & Br. n.* 802. *Berk. Out. p.* 219. *t.* 19. f. 3. *C. Hbk. n.* 659. *S. Mycol. Scot. n.* 631. *Fr. Hym. Eur. p.* 470. *Gonn. & Rab. viii.-ix. t.* 8. f. 2.

**10. M. erythropus Fr.**—**Pileus** about 2.5 cent. (1 in.) broad, hygrophanous, pale, almost whitish when dry, slightly fleshy, convex then plane, obtuse, slightly wrinkled when dry. **Stem** 5–7.5 cent. (2–3 in.) and more long, 4 mm. (2 lin.) thick, fistulose, firm, tough, round or at length compressed, dark red, smooth above, at the first paler, somewhat pruinose when dry, white-strigose below, internally villous at the sides. **Gills** separating-free, *broad, lax*, connected by veins, not crowded, quite entire, whitish.

Inodorous, juiceless, single or somewhat caespitose. The pileus varies in colour with the moisture of the atmosphere. It varies with the stem decumbent, even contorted. In habit it approaches nearest to *Ag. (Collybia) confluens* and *acervatus*; in these, however, the gills are very narrow and very crowded.

Among leaves near stumps. Frequent.

Pileus pallid, somewhat shaded with pink. *M. J. B.* Name—*ερυθρός*, red; *πούς*, a foot. Red-stemmed. *Fr. Monogr.* ii. p. 221. *Hym. Eur. p.* 470. *Icon. t.* 174. f. 2. *Berk. Out. p.* 220. *C. Hbk. n.* 660. *S. Mycol. Scot. Supp. Scot. Nat.* 1885, p. 74.

**11. M. archyropus** Fr.—**Pileus** 2.5 cent. (1 in.) broad, tan- Marasmius. colour, becoming pale, slightly fleshy, convex then plane or depressed, smooth. **Stem** very long, but scarcely 2 mm. (1 lin.) thick, stuffed then hollow, *rigid, tense and straight, pallid rufescent under the white tomentose pruina which forms an outer covering, similar at the base.* **Gills** adnexed, separating, crowded, linear, pallid.

Inodorous. Fasciculato-caespitose. Analogous with *M. terginus*. The covering of the stem is almost that of *M. oreades*.

Among leaves. Rare. Bristol.

Name—From the very tall stem. *Fr. Hym. Eur. p. 471. Berk. Out. p. 220. C. Hbk. n. 663. Gonn. & Rab. t. 8. f. 6. Ag. Pers. Myc. Eur. 3. p. 135. t. 25. f. 4.*

**12. M. torquescens** Quel.—**Pileus** pallid, disc tawny, membranaceous, thin, convexo-plane, *rugoso-striate*. **Stem** somewhat filiform, *delicately velvety*, brown, *smooth*, and whitish at the apex, twisted and sulcate when dry. **Gills** free, *thin*, ventricose, distant, white or reddish.

Among oak-leaves. Glamis, 1877.

The gills are finely serrulated. In the very young plant, when the pileus is conical, there is a slight indication of a veil. *B. & Br.* Name—*torqueo*, to twist. From the twisted stem. *Quell. p. 198. t. 22. f. 3. Fr. Hym. Eur. p. 471. B. & Br. n. 1678. S. Mycol. Scot. n. 632.*

**13. M. impudicus** Fr.—**Pileus** 1-2.5 cent. ( $\frac{1}{2}$ -1 in.) broad, bay-brown-rufous, becoming pale when dry, slightly fleshy at the disc, convex then plane, often depressed in the centre, soft; membranaceous, striato-plicate, and paler from the circumference to the middle. **Stem** 5 cent. (2 in.) long, 2 mm. (1 lin.) thick, fistulose, equal, but broader in becoming compressed, *rooting* at the attenuated *base*, tough, flexile; when moist rufous or rufous-fuscous, sometimes violaceous-purple, naked, but *wholly covered over with white villous down* (pruinato-velvety) *when dry.* **Gills** at first reaching the stem, then free, truncate behind, connected by veins, ventricose, at first crowded, then distant, whitish-flesh-colour.

Gregarious, odour strong, disgusting, of the same kind as that of *M. fæstens* or *perforans*, but very heavy.

On and about pine-trunks. Hanham. Sept.

Name—*impudicus*, disgusting. Of the odour. *Fr. Monogr. ii. p. 222. Hym. Eur. p. 471. Br. Bath. Trans. 1870, p. 77. B. & Br. n. 1789. C. Hbk. n. 662.*

Marasmius.

## C. CALOPODES.

\* *Stem very smooth and shining upwards, with a simple base.*

**14. M. scorodonius** Fr.—**Pileus** 12 mm. ( $\frac{1}{2}$  in.) and more broad, rufous when young, but soon becoming pale, whitish (not hygrophanous), slightly fleshy, pliant, convex then soon plane, obtuse, always arid; even when young, at length wrinkled and crisped. **Stem** 2.5 cent. (1 in.) long, scarcely 2 mm. (1 lin.) thick, *horny, tough, fistulose, equal, very smooth throughout, shining, rufous, inserted and naked at the base.* **Gills** adnate, often separating, connected by veins, at length crisped in drying, whitish.

Commonly gregarious. *Readily distinguished from neighbouring species by its strong odour of garlic.*

Heaths and dry pastures on twigs, &c. Rare.

Edible; esteemed for flavouring. Name—*σκόρδιον*, a plant that smells like garlic. *Fr. Monogr.* ii. p. 223. *Hym. Eur.* p. 472. *Sv. atl. Sv. t.* 32. *Berk. Out.* p. 220. *C. Hbk. n.* 644. *Ag. Schæff. t.* 99. *Paul. t.* 122 bis f. 2, 3. *Lenz f.* 17. *Sv. Bot. t.* 173?

**15. M. calopus** Fr.—**Pileus** about 8 mm. (4 lin.) broad, *whitish*, slightly fleshy, tough, convex then flattened, obtuse, rarely depressed, even, smooth, slightly wrinkled when dried. **Stem** 2.5 cent. (1 in.) long, 2 mm. (1 lin.) thick, fistulose, slightly attenuated upwards, even, *smooth, tough, rufous or bay-brown-rufous, shining, somewhat rooted.* **Gills** slightly emarginate, in groups of 2-4, thin, white.

*Inodorous.* Almost smaller than *M. scorodonius*, but the stem is longer, otherwise very like it.

On twigs, &c. Bristol.

In 'Hym. Eur.' the stem is described as *not rooted*. Name—*καλός*, beautiful; *πούς*, a foot. With beautiful stem. *Fr. Monogr.* ii. p. 223. *Hym. Eur.* p. 472. *B. & Br. n.* 1790. *Quel. t.* 13. f. 5. *Ag. Pers.—Bull. t.* 550. f. 1.

**16. M. Vaillantii** Fr.—**Pileus** 12 mm. ( $\frac{1}{2}$  in.) and more broad, whitish, somewhat membranaceous, pliant, soon flattened and depressed in the middle, *plicato-rugose.* **Stem** 2.5 cent. (1 in.) long, *stuffed, thickened and paler upwards*, here and there obsoletely pruinose; otherwise smooth, *date-brown*, bright shining, *inserted at the base*, where it is naked and blackish. **Gills** adnate, *somewhat decurrent on account of their triangular form*, broad, distant, distinct, simple, white.

Very remarkable. *Inodorous*, tough, arid. The pileus is plicate as in the *Rotulæ*.

On dead wood. Royal Botanic Garden, Regent's Park. Sept. Marasmius.

Perhaps imported. *B. & Br.* Name—after Vaillant. *Fr. Monogr.* ii. p. 223. *Hym. Eur.* p. 472. *Berk. Out.* p. 221. *C. Hbk. n.* 665. *Ag. Fr. Vaill. t.* 11. f. 21-23. *Buxb. iv. t.* 36. f. 2.

**17. *M. angulatus* B. & Br.—***Pileus* fuscous-whitish, somewhat membranaceous, hemispherical then flattened, *at length plicato-angular*. *Stem* 2.5 cent. (1 in.) and more long, thin, rufescent-grey, *thickened at both ends*, hairy at the base. *Gills* distant, whitish.

Very different from *M. Vaillantii*.

On grass. Cefn, Denbighshire, above Bone-cave.

Name—*angulus*, an angle. From the *angular* pileus. *B. & Br. n.* 1018. *Fr. Hym. Eur.* p. 473. *Ag. Pers. Myc. Eur.* 3. p. 155. t. 26. f. 3, 4. *Michel. t.* 74. f. 4 (*whitish var.*)

**18. *M. languidus* Fr.—***Pileus* 12 mm. ( $\frac{1}{2}$  in.) broad, white, inclining to flesh-colour or light yellow, slightly fleshy, at first convex, margin involute, then more expanded and umbilicate, *flocculose, rugoso-sulcate*. *Stem* scarcely 2.5 cent. (1 in.) long, 1-2 mm. ( $\frac{1}{2}$ -1 lin.) thick, stuffed, thickened upwards, naked, pallid, becoming fuscous and commonly white villous at the base. *Gills* adnate then decurrent, distant, connected by veins, *narrow, white*.

Gregarious, tough, pliant, inodorous, especially remarkable for its decurrent gills. *Its appearance is that of Agaricus.*

On dead leaves of grass. Coed Coch.

Name—*languo*, to be weak, languid. Limp and tough as opposed to brittle. *Fr. Monogr.* ii. p. 224. *Hym. Eur.* p. 473. *B. & Br. n.* 1017. *C. Hbk. n.* 667. *Ag. Lasch—Batt. t.* 27. f. O.

**\*\* *Stem velvety or pruinata, with a somewhat tubercular base.***

**19. *M. foetidus* Fr.—***Pileus* 12 mm. ( $\frac{1}{2}$  in.) broad, *bay-brown-tawny, membranaceous*, somewhat pellucid, pliant, convex then expanded, at length umbilicate, *striato-plicate*. *Stem* scarcely reaching 2.5 cent. (1 in.) long, 1 m. ( $\frac{1}{2}$  lin.) thick, horny, *fistulose*, equal, *date-brown, and everywhere velvety*, inserted with a very small floccose tubercle at the base. *Gills* adnexed in the form of a ring, somewhat decurrent when the disc is depressed, distant, connected by veins, *yellowish-rufescent*.

*Odour foetid.* There is another form differing in the colour being wholly umber, and in the absence of the villous tubercle at the base. This and the other plicate species differ from the *Rotulae* in the mycelium not being rhizomorphoid, and in the margin of the pileus being at first somewhat involute.

Marasmius. On decayed twigs, &c. Rare. Aug–Nov.

Name—*fœtidus*, stinking. *Fr. Monogr.* ii. p. 224. *Hym. Eur.* p. 473. *Berk. Out.* p. 221. *C. Hbk.* n. 668. *S. Mycol. Scot.* n. 633. *Ag. Sow.* t. 21.

**20. M. amadelphus** Fr.—**Pileus** 6–8 mm. (3–4 lin.) broad, pale yellowish, *becoming pale*, sometimes whitish, *fleshy-membranaceous*, convex then plane, always obtuse, at length depressed at the darker disc, *somewhat pruinose, even, at length striate only at the margin.* **Stem** curt, 12 mm. ( $\frac{1}{2}$  in.) long, scarcely reaching 2 mm. (1 lin.) thick, *stuffed*, equal, *somewhat mealy, pallid*, commonly inserted with an obsolete tubercle. **Gills** adnate, *somewhat decurrent, broad, distant, pallid.*

Gregarious, arid, *inodorous*. In a var. in fir woods the gills are still more distant, swollen in the form of veins, at length separating, rufescent-wood-colour (*pallid umber*); pileus thinner, whitish.

On dead branches, &c. Rare. Sept.

Name—*ἀμ, ἀδελφός*, brother. From its relation to *M. ramealis* next to which Fries placed it, though differing in some respects from the group. *Fr. Monogr.* ii. p. 225. *Hym. Eur.* p. 474. *Berk. Out.* p. 221. *C. Hbk.* 669. *S. Mycol. Scot.* n. 634. *Ag. Bull.* t. 550. f. 3.

**21. M. ramealis** Fr.—**Pileus** 6–8 mm. (3–4 lin.) broad, *white*, rufescent at the disc, *somewhat fleshy*, firm, convex then plane, obtuse or depressed in the centre, *slightly wrinkled*, but *without striæ.* **Stem** 6–10 mm. (3–5 lin.) long, scarcely 2 mm. (1 lin.) thick, *stuffed*, but commonly with a white line in the centre indicating that it is typically fistulose, often incurved, *mealy* (or with hairy squamules under a lens), *whitish*, or rufescent at the base, inserted when young at least with a very small tubercle. **Gills** adnate, connected behind, *slightly distant, narrow, white.*

Gregarious, arid, *inodorous*.

On dead twigs. Common. Sept.–Nov.

Under a lens clothed with minute matted silkiness. *M. F. B.* Name—*ramus*, a branch. Growing on twigs. *Fr. Monogr.* ii. p. 225. *Hym. Eur.* p. 474. *Berk. Out.* p. 221. *C. Hbk.* n. 670. *S. Mycol. Scot.* n. 635. *Ag. Bull.* t. 336. *Mich.* t. 74. f. 7.

## II.—MYCENA.

### A. CHORDALES.

**22. M. alliaceus** Fr.—**Pileus** 2.5–4 cent. (1–1½ in.) broad, whitish inclining to fuscous, often milk-white when young, somewhat membranaceous, campanulate then expanded, somewhat umbonate, even, at length striate and sulcate, smooth, dry. **Stem**



as much as 20 cent. (8 in.) long, *horny*, rigid, fistulose, attenuated upwards, *pruinato-velvety*, *blackish*, rooted at the base where it is somewhat incurved and naked. **Gills** adnexed in the form of a ring then *free*, slightly ventricose, arid, slightly distant, fuscous-whitish, crisped when dry. Marasmius.

Odour strong, of garlic, persistent. *There is nothing of a reddish tinge in the whole plant.* The stem is not tomentose at the base as in the *Tergini*.

Among leaves and on rotten wood. Frequent. Aug.-Oct.

Name—*allium*, garlic. From the smell. *Fr. Monogr.* ii. p. 226. *Hym. Eur.* p. 475. *Berk. Out.* p. 223. *C. Hbk. n.* 671. *S. Mycol. Scot. n.* 636. *Ag. Jacq. Austr. t.* 82. *Fl. Dan. t.* 1251. *Mich. Gen. t.* 78. f. 4. *Paul. t.* 122. f. 1.

**23. M. caulicinalis** Fr.—**Pileus** 12 mm. ( $\frac{1}{2}$  in.) and more broad, dingy yellow becoming ferruginous, then ochraceous, membranaceous, thin, pliant, campanulate then convex, obtuse, at length plane, *never umbilicate*, even then striato-sulcate. **Stem** 2.5–5 cent. (1–2 in.) long, *fistulose with a pith*, rigid, tough, *floccoso-villous* and *bay-brown* below, *attenuated*, *paler* and *mealy* upwards, somewhat inserted at the base. **Gills** *adnato-decurrent*, somewhat distant, connected by a network of veins, *pallid light yellow*.

It differs widely from *Ag. campanella*, with which it agrees in stature and habit. *A. caulicinalis* Bull. is entirely different.

Among leaves. Rare. Nov.

It occurs paler in colour. Spores  $4.5 \times 2.3$  mk. *B.* Name—*caulis*, a stalk, stem. Growing on stems. *Fr. Monogr.* ii. p. 227. *Hym. Eur.* p. 476. *B. & Br. n.* 1136. *C. Hbk. n.* 672. *S. Mycol. Scot. n.* 637. *Ag. With.—Sow. t.* 163.

## B. ROTULÆ.

\* *Stem very smooth, shining.*

**24. M. rotula** Fr.—**Pileus** 6–8 mm. (3–4 lin.) broad, whitish, unicolorous, or with the umbilicus becoming fuscous, arid, membranaceous, slightly convex, *umbilicate*, plicate, undulato-crenulated at the margin. **Stem** variable in length, *horny*, fistulose, equal, shining, *very smooth*, *blackish*, manifestly striate when dry. **Gills** *adnate behind to a collar encircling the stem but removed from it*, broad, few (often equal), very distant, white.

Gregarious, often cæspitose, with root-like branchlets.

On dead twigs. Common. July–Nov.

Stem deep shining brown at the base. *M. J. B.* Name—*rota*, a wheel. Like a little wheel. *Fr. Monogr.* ii. p. 229. *Hym. Eur.* p. 477. *Berk. Out. p.* 222. t. 14. f. 7. *C. Hbk. n.* 673. *S. Mycol. Scot. n.* 638. *Gonn. &*

Marasmius. *Rab. t.* 8. *f.* 8. *Ag. Scop.*—*Sow. t.* 95. *Bull. t.* 64, 569. *f.* 3. *Fl. Dan. t.* 1134. *Mich. t.* 74. *f.* 5.

**25. M. graminum** Berk.—**Pileus** scarcely exceeding 6 mm. (3 lin.) broad, very pale rufous, the furrows paler, umbo brown, nearly plane, *umbonate, sulcate*. **Stem** quite *smooth*, shining, *black*, white above. **Gills** attached to a free collar, few, somewhat ventricose, even, with veiny interstices, cream-coloured.

A most elegant species, and quite distinct from *M. rotula*.

On grass. Uncommon. July–Sept.

Stem 2.5 cent. (1 in.) and more long. Gills 6–8 only. *Fr.* Name—*gramen*, grass. On grass. *Berk. Out. p.* 222. *t.* 14. *f.* 8. *C. Hbk. n.* 674. *S. Mycol. Scot. n.* 639. *Fr. Hym. Eur. p.* 477. *Ag. Lib. exs.* 119.

**26. M. androsaceus** Fr.—**Pileus** 4–6 mm. (2–3 lin.) broad, whitish or somewhat fuscous, membranaceous, arid, somewhat umbilicate, smooth, striate. **Stem** 4 cent. (1½ in.) and more long, filiform, horny, fistulose, very tough, equal, *very smooth, black*, contorted and striate when dry. **Gills** *adnate to the stem* (without a collar), *simple, distinct*, distant, whitish.

There are two forms: on deciduous leaves the pileus is whitish, deeply umbilicate, plicate, the mycelium (on oak-leaves) creeping within the substance of the leaf; on perennial leaves (*Pinus silvestris*, Juniper, here and there on bark) the pileus is scarcely umbilicate, more even, somewhat fuscous, and the mycelium commonly free.

On leaves, twigs, &c. Common. April–Nov.

The umbilicus is darker. Name—*androsaces*, a zoophyte: *madrepora acetabulum* Linn. Perhaps from its form. *Fr. Monogr. ii. p.* 229. *Hym. Eur. p.* 477. *Berk. Out. p.* 222. *C. Hbk. n.* 675. *S. Mycol. Scot. n.* 640. *Ag. Linn.*—*Fl. Dan. t.* 1551. *f.* 1. *Bolt. t.* 32. *Sow. t.* 94. *Bull. t.* 569. *f.* 2. *Bocc. Mus. t.* 104.

**27. M. splachnoides** Fr.—**Pileus** somewhat membranaceous, convex then expanded and umbilicate, smooth, striate. **Stem** horny, fistulose, smooth, shining, *red* ("becoming fuscous"). **Gills** somewhat decurrent, crowded, simple and anastomosing, white.

Inodorous. Allied most closely to *M. androsaceus*; differing however in the red stem, and somewhat decurrent, more crowded gills.

Among pine-leaves. Cabalva, Foxley. Oct.

By an error in 'Hym. Eur.' it is represented as being of the habit of *M. perforans*. See 'Epicr.' and 'Monogr.' Name—*splachnum*, a moss (σπλάχνον, applied to a cryptogamic plant); εἶδος, appearance. *Splachnum*-like. Resemblance not apparent. *Fr. Hym. Eur. p.* 478. *Monogr. ii. p.* 230. *Grevillea, vol. viii. p.* 110.

**28. M. Curreyi** B. & Br.—**Pileus** pallid rufous, furrows paler, Marasmius. umbo fuscous, somewhat plane, sulcate, somewhat radiate. **Stem** inserted, very smooth, shining, black, white at the apex. **Gills** attached to a collar, few, somewhat ventricose, cream-coloured, interstices veined, elsewhere very even.

On leaves of grasses. Fineshade, Norths. Aug.

Name—after F. Currey. *B. & Br. n. 1794.*

**\*\* Stem velvety or pilose.**

**29. M. perforans** Fr.—**Pileus** 8 mm. (4 lin.) broad, whitish, becoming pale-rufescent, somewhat membranaceous, *rather plane*, rarely depressed, not umbilicate, *without striæ*, at length *slightly wrinkled*. **Stem** 2.5 cent. (1 in.) or little more long, fistulose, not very horny, but tough, equal, even, *everywhere velvety, bay-brown then black*, inserted at the base. **Gills** adnate, numerous, *simple, unequal* (the alternate ones shorter), not very distant.

*Odour fetid*, not of garlic. In appearance like the *Calopodes*.

On fir-leaves. Rare. Aug.

Name—*perforo*, to pierce. Perforating with its stem the leaves on which it grows. *Fr. Monogr. ii. p. 230. Hym. Eur. p. 478. Berk. Out. p. 223. C. Hbk. n. 676. S. Mycol. Scot. n. 641. Ag. Hoffm. Nomencl. t. 4. f. 2. Schæff. t. 239. Batsch f. 10.*

**30. M. insititius** Fr.—**Pileus** 12 mm. (½ in.) broad, white, membranaceous, pliant, convexo-plane, somewhat umbilicate, unpolished, even when young, then plicato-sulcate. **Stem** scarcely 2.5 cent. (1 in.) long, horny, fistulose, *attenuated downwards* into a simple (not tubercular) inserted base, *floccoso-furfuraceous*, rufous inclining to fuscous. **Gills** broadly adnate to the stem and attenuated to the margin, *distant, simple, unequal*, pale to white.

Inodorous. Gregarious, arid. A very remarkable species. The habit is that of *M. Vaillantii*.

On leaves, decayed grass, &c. Rare. Aug.

Variable. Sometimes the stem is paler and the pileus less plicate. Sometimes the disc is reticulated. *M. J. B.* Name—*insero*, to insert or graft. From the way in which the stem is engrafted on the host plant. *Fr. Monogr. ii. p. 230. Hym. Eur. p. 478. Berk. Out. p. 223. t. 14. f. 6 (white-stemmed var.) B. & Br. n. 1792. C. Hbk. n. 677. S. Mycol. Scot. n. 642.*

**31. M. Hudsoni** Fr.—**Pileus** scarcely 4 mm. (2 lin.) broad, pale fuscous-rufescent, membranaceous, hemispherical, slightly

**Marasmius.** wrinkled, beset with scattered purple hairs. **Stem** horny, filiform, dark purple, with the same hairs as the pileus. **Gills** adnexed, narrow, single, white, the alternate ones dimidiate.

Inodorous.

On fallen holly-leaves. Frequent in England. Rare in Scotland. June-Dec.

The whole of the outer surface of the pileus is clothed with echinulate processes. *B. & Br.* Spores fusiform, 11 mk., with a central nucleus very different from those of any other Agaric or Marasmius. *B. & Br.* Name—after Hudson, author of 'Flora Anglica.' *Fr. Hym. Eur. p. 478. Berk. Out. p. 223. C. Hbk. n. 678. S. Mycol. Scot. n. 643. B. & Br. n. 708\*. t. 15. f. 3 (spores). Sow. t. 164.*

**32. M. epichloe** Fr.—**Pileus** whitish, bay-brown-fuscos in the centre, thin, plano-convex, somewhat papillate, without striæ. **Stem** bay-brown, opaque, sulcato-striate, the striae slightly bristly, paler at the base. **Gills** rounded, somewhat crowded, broader behind.

On the base of grasses. Hereford.

Name—ἐπί, upon; χλόη, grass. *Fr. Hym. Eur. p. 479. B. & Br. n. 1566. M. gramineus Leveill.*

**33. M. saccharinus** Fr.—**Pileus** wholly white, membranaceous, convex, somewhat papillate, smooth, sulcate and plicate. **Stem** very thin, filiform or attenuated upwards, flocculose then becoming smooth, inserted obliquely, reddish, pale at the apex. **Gills** broadly adnate, narrow, thick, very distant, reticulato-united, whitish.

Stem often rufous-fuscos below. Tending more to the form of *Mycena* than the rest.

On dead twigs. Rare. King's Cliffe, &c.

Name—saccharon, sugar. *Granulate.* From the papillate pileus. *Fr. Hym. Eur. p. 479. Monogr. ii. p. 231. Berk. Out. p. 224. B. & Br. n. 1793. C. Hbk. n. 680. S. Mycol. Scot. n. 644. Ag. Batsch f. 83.*

**34. M. epiphyllus** Fr.—**Pileus** 4-10 mm. (2-5 lin.) broad, milk-white, membranaceous, very thin, convex then plane, at length umbilicate, smooth, at length plicato-rugose. **Stem** 1-2.5 cent. ( $\frac{1}{2}$ -1 in.) long, very thin, somewhat horny, fistulose, equal, velvety under a lens, date-brown, whitish at the apex, inserted without mycelium at the base. **Gills** adnate, few (4-10), very distant, entire, veined, branched, white.

Gregarious, very slender but persistent, inodorous. When dried it revives with moisture. Sometimes there are only wrinkles in place of gills. *Batt. t. 28. D. Batsch f. 84 (very young).* It occurs also wholly even beneath. *Pers. Ic. & descr. t. 9. f. 7.*

On dead leaves, twigs, &c. Common. Sept.—Oct.

Marasmius.

In large specimens the gills form a close collar round the stem, margin of the collar cream-coloured. *M. J. B.* Name—ἐπί, upon; φύλλον, a leaf. *Fr. Monogr.* ii. p. 231. *Hym. Eur.* p. 479. *Berk. Out.* p. 224. *C. Hbk. n.* 679. *S. Mycol. Scot. n.* 645. *Ag. Tratt. Austr. f.* 22. *Fl. Dan. t.* 1194. *f. 1. Sow. t.* 93.

### III.—APUS.

**35. *M. spodoleucus* Berk.**—**Pileus** 4 mm. (2 lin.) broad, cinereous, pulverulent or slightly furfuraceous, *conchate, resupinate*, margin free, arched. **Stem** none. Hymenium white, very even. **Gills** few, narrow, entire, so short as to have a naked space at the base, white, interstices even.

On dead elm-twigs. Batheaston.

Spores very small, 1–2 mk. *W. G. S.* Name—σποδός, wood-ash; λευκός, white. From the cinereous pileus and white hymenium and gills. *B. & Br. n.* 803. *Berk. Out.* p. 224. *C. Hbk. n.* 681.

**36. *M. Broomei* Berk.**—Half-resupinate. Pallid brown then striate black; hymenium shining white. **Gills** distant, veined; interstices even.

On dead twigs. Batheaston. Jan.

Name—after C. E. Broome. *B. & Br. n.* 1795.

GENUS XV.—**Lentinus** (*lentus*, tough or pliant). *Fr. Epicr.* p. 387. *Lentinus*.

Pileus fleshy-coriaceous, pliant, or in fleshy species becoming hard when old, persistent. Hymenophore continuous with the stem or the base of the pileus when sessile. Gills concrete with the hymenophore, thin, unequal, membranaceous, *with the edge serrated or torn in a toothed manner*. Spores somewhat round, even, white. *With the exception of one species, growing on wood, polymorphous, somewhat irregular, very numerous in warmer countries, decreasing in number and not becoming so hard towards the north.* *Fr. Hym. Eur. p.* 480.

A very natural, though very poly-



LIV. *Lentinus cochleatus*. One-fourth natural size.

Lentinus. morphic genus. The serrated or torn edge of the gills is a distinctive feature.

I. MESOPODES (μεσός, middle; πούς, a foot). Pileus nearly entire. Stem distinct.

\* Lepidei (λεπίς, a scale). *Pileus scaly*. More or less manifestly veiled. (In none of the following groups is there a veil.)

\*\* Pulverulenti (*pulvis*, dust). *Pileus villous* or *pulverulent*.

\*\*\* Cochleati (*cochlea*, a snail-shell, lobed and twisted in shape). *Pileus smooth*.

II. PLEUROTİ (πλευρόν, a side; οὖς, an ear). Dimidiate, sessile or furnished with a somewhat lateral stem.

### I.—MESOPODES.

\* Lepidei. *Pileus scaly*, &c.

1. **L. tigrinus** Fr.—**Pileus** commonly 5 cent. (2 in.) broad, white, *variegated* with somewhat adpressed, *blackish*, *hairy squamules*, fleshy-coriaceous, *thin*, commonly orbicular and central, at first convexo-plane, *umbilicate*, *at length infundibuliform*, often split at the margin when dry. **Stem** about 5 cent. (2 in.) long, *thin*, solid, very hard, commonly attenuated downwards, minutely *squamulose*, whitish, often ascending and becoming fuscous at the base, at first furnished *at the apex with an entire reflexed ring, which soon falls off*. **Gills** decurrent (*by no means sinuate*), narrow, crowded, unequal, toothed like a saw, white.

Somewhat gregarious, even cæspitose, thinner and more coriaceous and regular than *L. lepideus*. B. wholly *blackish* with squamules.

On old stumps. Rare.

When fresh very tender and easily torn, when dry coriaceous. *Sow.* Smell strong, acrid, like that of some *Lactarii*. *M. J. B.* Spores  $3 \times 7$  mk. *W. G. S.* Name—*tigrinus*, tiger-like. From the marking on the pileus. *Fr. Monogr.* ii. p. 233. *Hym. Eur.* p. 481. *Berk. Out.* p. 224. *C. Hbk. n.* 682. *Ag. Bull. t.* 70. *Sow. t.* 68. *Batt. t.* 12. f. B-D.

\* **L. Dunalii** Fr.—**Pileus** pallid, *with adpressed, separating, spot-like scales*, fleshy-coriaceous, thin, umbilicate, irregularly shaped. **Stem** short, *somewhat silky*. **Gills** decurrent, crowded, pallid.

When the scales separate the pileus is even. Veil scarcely manifest. Very much allied to *L. tigrinus*.

On ash-trees. Rare.

Cæspitose. Pileus 2 in. broad, margin deflexed, sometimes variously split and sinuated, yellow-white, with brownish rather close scales, margin nearly smooth: gills dichotomous, crenate. Stem 1 in. high, 3 lin. thick, tough, the lower part clothed with dark, nearly square, adpressed scales, the upper half not scaly, white, resembling the under side of white kid leather. Odour sub-

acid, farinaceous. *M. F. B.* Name—after Dunal. *Fr. Hym. Eur. p. 481.* Lentinus. *Berk. Out. p. 225. t. 15. f. 2.* *C. Hbk. n. 683.* *Batt. t. 12. A.*

**2. L. lepidus** Fr.—**Pileus** 5–10 cent. (2–4 in.) broad, pallid-ochraceous, variegated with adpressed, *darker, spot-like scales, fleshy*, very *compact* and firm, *irregular*, commonly excentric, convex then depressed, but not truly umbilicate, sometimes broken up into cracks; flesh pliant, white. **Stem** short, commonly 2.5 cent. (1 in.) long, solid, *stout*, very *irregularly formed*, almost woody, tomentoso-scaly, whitish, rooted at the base, *at the first furnished with a cortina towards the apex.* **Gills** decurrent, but *sinuate behind*, crowded, broad, transversely striate, whitish, edge torn into teeth.

Odour pleasant. Almost always solitary; earlier than the rest, often appearing quite in spring. Very variable in stature and size. Cortina very fugacious. Gills very remarkable in being *sinuate behind* while decurrent. Often pierced by larvæ when old. *Sow. t. 382* is a monstrous form of this.

On pine. Rare. July.

\* Fries refers to various monstrous forms found in close places, cellars, aqueducts, &c. I have one such from under a floor; more than 2 feet in height, the very thick woody stem springing from a dense leathery stratum, which covered yards of the wood, then branched and united in fantastic forms, like branched coral, with here and there a small infundibuliform pileus on the end of a branch. Spores very white. *Fr. ; 11 × 5 mk. W. G. S.* Name—*λεπίς*, a scale. From the pileus. *Fr. Monogr. ii. p. 233.* *Hym. Eur. p. 481.* *Berk. Out. p. 225.* *C. Hbk. n. 684.* *S. Mycol. Scot. n. 646.* *Ag. Schaff. t. 29, 30.* *Buxb. C. iv. t. 25.*

\*\* Pulverulent. *Pileus villous or pulverulent.*

**3. L. leontopodius** Schulz.—**Pileus** tan-clay-colour, fleshy-coriaceous, tough, irregular, *delicately tomentose*, disc depressed, the deflexed margin slightly lobed. **Stem** thick, woody, unpolished, pulverulent, pale chestnut, *blackish downwards.* **Gills** decurrent, connected by veins, wrinkled at the sides, serrated at the edge.

Large, handsome, with a pleasant taste. Pileus here and there cinnamon with the down.

On decayed willow. Menmuir, Forfarshire.

Name—*λέων*, a lion; *πούς*, a foot. From the appearance of the stem. *Schulz.—Kalchbr. in litt.* *Fr. Hym. Eur. p. 482.* *S. Mycol. Scot. n. 647.*

**4. L. pulverulentus** Fr.—**Pileus** yellow, *mealy with white dust*, fleshy-pliant, convex. **Stem** stout, elongated, equal, rigid, *mealy with white dust.* **Gills** slightly toothed, white.

Cæspitose.

Lentinus.

On dead trunks. Glamis, 1875. Oct.

Tufted, at first infundibuliform, then lateral flabelliform, fuliginous, floccoso-pulverulent, with little umber particles; stem elongated, at length smooth; gills thick, pallid, deeply decurrent, their edge crenulate but not torn. Pileus 2 inches across, stem 3 inches high. *B. & Br.* Name—*pulvis*, dust. Pulverulent. *Fr. Hym. Eur. p.* 483. *B. & Br. n.* 1567. *S. Mycol. Scot. n.* 648. *Ag. Scop. Carn. p.* 434.

\* **L. resinaceus** Fr.—**Pileus** *ochraceous-cinnamon*, somewhat fleshy, pliant, excentric, *somewhat gummy-villous*. **Stem** unequal, tomentose. **Gills** crowded, serrated, shining white.

Somewhat *cæspitose*. Covered with a dry, tenacious, somewhat resinous gluten arising from the quasi-glandular villous down.

On trunks. Forres.

Name—*resina*, resin. From the *resinous* covering. *Fr. Hym. Eur. p.* 483. *B. & Br. n.* 1283. *S. Mycol. Scot. n.* 649. *Ag. Trog. Fl.* 1832. *p.* 525.

5. **L. adhærens** Fr.—**Pileus** dingy pallid, somewhat fleshy, pliant, irregular, convexo-umbonate then depressed and infundibuliform, lacunose, somewhat pulverulent, *coated with a glutinous varnish*. **Stem** *somewhat hollow*, rooted, of the same colour as the pileus and like it *coated with a glutinous varnish*. **Gills** decurrent in lines, very thin, torn, white.

Small, unequal, taste at length astringent.

In pine woods. Rare.

Name—*ad*, and *hæreo*, to stick. From the glutinous varnish. *Fr. Hym. Eur. p.* 483. *C. Hbk. n.* 685. *S. Mycol. Scot. n.* 649. *Ag. A. & S.—A. adhærens* With.

\*\*\* Cochleati. *Pileus smooth.*

6. **L. cochleatus** Fr.—**Pileus** 5–7.5 cent. (2–3 in.) broad, flesh-colour, but becoming pale, somewhat tan, fleshy-pliant, thin, commonly excentric, imbricated, very unequal, somewhat lobed or contorted, sometimes plane, sometimes infundibuliform-umbilicate, but not pervious, *smooth*. **Stem** solid, firm, sometimes central, most frequently excentric, sometimes wholly lateral, *always sulcate, smooth*, flesh-coloured upwards, rufous-fuscon downwards. **Gills** decurrent, crowded, serrated, white-flesh-colour.

Pliant, tough, *flaccid*, very changeable in form, sometimes solitary, sometimes *cæspitose*, imbricated, growing into each other. From very small forms, which are commonly solitary, with the stem and pileus scarcely 2.5 cent. (1 in.) it ranges to 7.5 cent. (3 in.)

On stumps. Frequent. Aug.–Oct.



According to Fries the odour is weak, of anise; but it is generally strong *Lentinus*. and very pleasant. Spores sphaeroid or ellipsoid-sphaeroid, uniguttate, 4-6 mk. *K.*; almost globular, 4 mk. *W.G.S.* Name—*cochlea*, a snail-shell. From the shape. *Fr. Monogr.* ii. p. 235. *Hym. Eur.* p. 484. *Berk. Out.* p. 226. *t.* 19. *f.* 4. *C. Hbk.* n. 686. *S. Mycol. Scot.* n. 650. *Price t.* 125. *Ag. Sow. t.* 168.

## II.—PLEUROT. Dimidiate, &c.

**7. *L. scoticus*** B. & Br.—**Pileus** 1-4 cent. ( $\frac{1}{2}$ -1 $\frac{1}{2}$  in.) broad, *pallid*, at length brownish, hygrophanous, *umbilicate*, sometimes infundibuliform, at length flattened; extremely variable in form, either quite stemless and *reniform*, or variously stipitate, lobed at the margin and sinuate, or plicate. **Stem** when present varying from 4 mm. (2 lin.) to 5 cent. (2 in.) long, *cylindrical*, darker, *pulverulent*, *springing from a brown, fibrillose mycelium*. **Gills** decurrent when the stem is developed, rather distant, strongly toothed.

Solitary or caespitose; sometimes 2 pilei are joined. Inodorous, or at any rate without any odour of aniseed. The nearest ally to this curious species is *L. omphalodes*.

On decayed *ulix*, birch and spruce. Menmuir. Glamis. Nov.-Jan.

The teeth of the gills are very irregularly torn. Name—*Scottish*. *B. & Br.* n. 1423. *S. Mycol. Scot.* n. 651. *Fr. Hym. Eur.* p. 485.

**8. *L. fimbriatus*** Curr.—**Pileus** 1-2.5 cent. ( $\frac{1}{2}$ -1 in.) broad, fawn-colour, covered with darker floccose scales, somewhat dimidiate, somewhat coriaceous, thin (not fleshy), depressed, sometimes very much so and almost cyathiform, margin slightly involute, almost strigose. **Stem**  $\frac{1}{8}$ - $\frac{1}{4}$  in. long, lateral, rough with somewhat reflexed scales of the same colour as the gills, or rather paler. **Gills** descending, but not decurrent, irregularly serrated and torn at the margin, pale brown.

In young specimens a delicate white fimbriate collar or fringe (the remains of the ruptured veil) separates the gills from the stem. Pilei two or three together, one above another in an imbricated manner. Some of the pilei tinged here and there with pink stains.

On a stump standing in a pond. Lewes. Sept.

Name—*fimbriæ*, fringe. From the *fimbriate* collar. *Currey in Trans. Linn. Soc.* xxiv. p. 152. *t.* 25. *f.* 2. *C. Hbk.* n. 688. *Fr. Hym. Eur.* p. 485.

**9. *L. vulpinus*** Fr. Sessile, *many times imbricated*.—**Pilei** *tan*, fleshy, very pliant, *reniform-conchate*, very convex, margin deflexed and almost perpendicular, hence concave beneath; *surface* wholly peculiar, *with raised longitudinal ribs*, which are

Lentinus.

broken up into scales or fibrous teeth towards the margin, hence entirely rough and corrugated, and white-warty behind. **Gills** extended to the base, *broad*, crowded, torn into teeth.

Pilei like steps of a ladder (the intermediate ones largest, 7.5 cent. (3 in.) broad), horizontal, connate behind in a dense cluster, the common base piercing the trunks of trees. Substance white. Margin thin, incurved, becoming fuscous in older specimens. When dry the pileus often gapes open in longitudinal cracks. The gills at the base are concrete with the substance of the pileus. It sheds *very plentiful, white spores*, hence the clusters when old are very white-mealy.

On stumps. Rare.

Smell very strong and overpowering, somewhat resembling that of field mint. *M. J. B.* Spores almost globular, very small, 2 mk. *W. G. S.* Name—*vulpes*, a fox; foxy. Meaning obscure. *Fr. Monogr.* ii. p. 238. *Hym. Eur.* p. 486. *Icon. t.* 176. *f.* 1. *Berk. Out.* p. 226. *C. Hbk. n.* 687. *Sow. t.* 351.

**10. L. flabelliformis** Fr. Somewhat sessile.—**Pileus** not exceeding 2.5 cent. (1 in.) broad, pallid fawn-colour, *membranaceous*, pliant, *reniform*, *plane*, even, *smooth*, *margin fimbriato-toothed*. **Stem** commonly *rudimentary*, thin, 4–6 mm. (2–3 lin.) long. **Gills** broad, somewhat distant, rather thick, pallid or whitish, torn into teeth at the edge.

Commonly *exactly lateral with a rudimentary stem*; much smaller and *thinner* than any of the rest, at length crisped. Var. *herbarum* very small, the membranaceous pileus becoming pale.

On stumps. Rare.

Perhaps Bolton's plant may be only *Ag. salignus*. *M. J. B.* Name—*flabellum*, a small fan; *forma*, form. From the shape. *Fr. Monogr.* ii. p. 239. *Hym. Eur.* p. 487. *Berk. Out.* p. 226. *C. Hbk. n.* 689. *Ag. Bolt. t.* 157?

Panus

GENUS XVI.—**Panus** (a name given to an arboreal fungus by Pliny.) *Fr. Epicr.* p. 396.

Whole fungus fleshy-coriaceous, tough, drying up, of fibrous texture, which radiates into the hymenium. Gills concrete with the hymenophore, unequal, at length *coriaceous*, *edge quite entire*. Spores even, white, somewhat cylindrical in species which have been examined. *Growing on wood, various in form, lasting long.*

A genus which must be inserted in this series on account of its flesh which is pliant and somewhat coriaceous (even in the gills), allied to the *Lentini*, but differing from them in the firmer, *coriaceous and very entire* gills. Either poisonous, or owing to the toughness of the substance not suitable for eating. *Fr. Hym. Eur.* p. 487.

\* *Pileus irregular, stem excentric.*

\*\* *Stem definitely lateral.*

\*\*\* *Pileus resupinate, sessile or extended behind.*

Panus.

\* *Pileus irregular, stem excentric.*

1. **P. conchatus** Fr.—**Pileus** 5–10 cent. (2–4 in.) broad, *cinnamon then becoming pale*, fleshy-pliant, thin, unequal, *excentric*, even *dimidiate*, *flaccid*, *squamulose* when old. **Stem** about 12 mm. ( $\frac{1}{2}$  in.) long, 8 mm. (4 lin.) thick, unequal, often compressed, *pubescent at the base*. **Gills** deeply decurrent in parallel lines, by no means anastomosing, but here and there branched and unequal, at first whitish or pale flesh-colour, at length ochraceous-wood-colour, crisped when dry.

Caespitose, often imbricated and growing into each other. No form is constant. So much allied to *P. torulosus* that the real difference is not apparent. It is thinner, more conchate and more lobed than that species.



LV. *Panus torulosus*. One-fourth natural size.

On trunks. Rare. June–Oct.

Name—*concha*, a shell. Shell-shaped. *Fr. Monogr.* ii. p. 240. *Hym. Eur.* p. 488. *Berk. Out.* p. 227. *C. Hbk. n.* 691. *S. Mycol. Scot.* n. 652. *Krombh. t.* 42. f. 1, 2. *Schæff. t.* 43, 44. *Bull. t.* 298, 517. f. O. P.

2. **P. torulosus** Fr.—**Pileus** 5–7.5 cent. (2–3 in.) broad, somewhat flesh-colour, but varying rufescent-livid and becoming violet, *entire*, but very excentric, fleshy, somewhat compact when young, *plano-infundibuliform*, *even*, smooth; flesh pallid. **Stem** short, commonly 2.5 cent. (1 in.), solid, oblique, tough, firm, commonly with grey, but often violaceous *down*. **Gills** decurrent, somewhat distant, simple, separate behind, reddish then tan-colour.

Very changeable in form, at first fleshy-pliant, at length coriaceous. In the covering of the stem it approaches *Paxillus atro-tomentosus*, but there is no affinity between them.

On old stumps. Frequent.

Spores  $6 \times 3$  mk. *W.G.S.* Name—*torulus*, a tuft of hair. From the hairy down on the stem. *Fr. Monogr.* ii. p. 240. *Hym. Eur.* p. 489. *Berk. Out.* p. 226. *C. Hbk. n.* 690. *S. Mycol. Scot. Supp. Scot. Nat.* 1881, p. 37. *Ag. Krombh. t.* 42. f. 3–5. *Bolt. t.* 146 (var.) *Gonn. & Rab. t.* 12. f. 4. *Batsch f.* 33. *Paul. t.* 26. f. 3, 4. *Nees Syst. f.* 176.

Panus.

**\*\* Stem definitely lateral.**

**3. P. stipticus** Fr.—**Pileus** 1-2.5 cent. ( $\frac{1}{2}$ -1 in.) broad, cinnamon becoming pale, arid, thin, but not membranaceous, *reniform*, pruinose, the *cuticle separating into furfuraceous scales*. **Stem** not reaching 2.5 cent. (1 in.) long, solid, *definitely lateral*, compressed, *dilated upwards*, ascending, pruinose, paler than the gills. **Gills** ending determinately (not decurrent), thin, very narrow, crowded, *elegantly connected by veins*, cinnamon.

Gregarious, caespitose, remarkable for its *astringent taste*. The pileus sometimes has an infundibuliform appearance with lobes all round.

On stumps, &c. Common. Aug.-Feb.

Reckoned poisonous. Spores obovoid-sphaeroid,  $2-3 \times 1-2$  mk. *K.*;  $3 \times 4$  mk. *W.G.S.* Name—*stypticus*, astringent. *Fr. Monogr.* ii. p. 241. *Hym. Eur.* p. 489. *Berk. Out.* p. 227. *C. Hbk.* n. 692. *S. Mycol. Scot.* n. 653. *Ag. Bull.* t. 140, 557, f. 1. *Schæff.* t. 208. *Sow.* t. 109. *Fl. Dan.* t. 832, f. 1, 1292, f. 1. *Tratt. Austr.* t. 2. *Krombh.* t. 44, f. 13-17. *Buxb. C.v.t.* 10, f. 1.

**4. P. farinaceus** Schum.—**Pileus** cinnamon-umber, somewhat coriaceous, flexuous, *cuticle separating into whitish-bluish-grey scurf*. **Stem** short, lateral, of the same colour as the pileus. **Gills** determinately free, *distinct*, paler.

The habit is that of *P. stipticus*.

On fir pole. Glamis, 1873. Dec.

Name—*farina*, meal. From the scurf on the pileus. *Schum.*—*Fr. Hym. Eur.* p. 490. *S. Mycol. Scot.* n. 654.

**\*\*\* Pileus resupinate, sessile or extended behind.**

**5. P. patellaris** Fr.—**Pileus** 12 mm. ( $\frac{1}{2}$  in.) or a little more broad, pallid externally, *resupinate*, coriaceous, orbicular, *planocup-shaped*, adnate by the sessile vertex, the free margin involute, furfuraceous-villous externally. **Gills** concurrent in a central point, crowded, quite entire, arid, dark ochraceous.

Very remarkable, nearest to *P. stipticus*.

On cherry. Forres. March.

Name—*patella*, a small dish. *Fr. Monogr.* i. p. 242. *Hym. Eur.* p. 490. *Icon.* t. 176, f. 3. *B. & Br.* n. 1680. *S. Mycol. Scot.* n. 655.

**6. P. Stevensoni** B. & Br.—**Pileus** olivaceous-light yellow, spatulate; flesh greenish-yellow. **Stem** dilated upwards, convex and golden, slightly hispid. **Gills** narrow, entire.

On oak. Glamis, 1877. Penicuik, 1878. Sept.-Oct.

A very striking species. Name—after Rev. John Stevenson. *B. & Br. n.* 1796. *S. Mycol. Scot. n.* 656.

GENUS XVII.—**Xerotus** (ξηρός, dry; ὄς, an ear).

Xerotus.

Fr. Elench. i. p. 48.

Hymenophore continuous with the stem, descending into the trama which is homogeneous with the *coriaceous pileus*. Gills coriaceous, broadly plicæform, dichotomous, edge quite entire, obtuse. *Rigid, persistent, analogous with the Cantharelli, but differing in the whole structure.* Fr. *Hym. Eur. p.* 491.



LVI. *Xerotus degener*. One-half natural size.

1. **X. degener** Fr.—**Pileus** date-brown-grey when moist, when dry grey, *somewhat zoned*, coriaceous-membranaceous, very thin, but very tough, plano-infundibuliform, *striate when moist*, flocculose when dry. **Stem** not 2.5 cent. (1 in.), often only 4–6 mm. (2–3 lin.) long, 2 mm. (1 lin.) thick, very tough, solid or floccoso-stuffed at the apex, equal, fuscous, somewhat white-velvety. **Gills** decurrent, few, very distant, when rightly developed thin, rather broad, whitish-grey, edge acute, flaccid.

When dry the pileus is zoneless in smaller specimens. When the gills are less fully developed, they resemble the ribs of *Cladoderis*, the edge being very obtuse. They vary, sometimes all equal, simple, sometimes simply dichotomous, with shorter alternate ones. They are the most widely distant of any of the species of *Agaricini*.


On peaty soil among short grass. Glamis, Warren, 1877. Jan.

This extremely rare species has hitherto been recorded as British by Sowerby only. My specimens, in the opinion of Berkeley were nearest *Schaff. t.* 243. Spores elliptical, irregular, 8–12 × 4–6 mk. *B.* Name—*degener*, degenerate. Fr. *Monogr. ii. p.* 243. *Hym. Eur. p.* 491. *Berk. Out. p.* 227. *C. Hbk. n.* 693. *S. Mycol. Scot. n.* 657. *Ag. Schaff. t.* 243. *A. turfusus Sow. t.* 210.

Trogia.

GENUS XVIII.—**Trogia** (after Trog, a Swiss botanist).  
Fr. Epicr. p. 402.

Gills fold-like, edge longitudinally channelled (in the single European species only crisped). In other respects agreeing with *Xerotus*. *Soft, flaccid, but arid and persistent, texture fibrillose.* Fr. Hym. Eur. p. 491.



LVIII. *Trogia crispa*.  
Natural size.

Reviving when wet. Spores white.

1. **T. crispa** Fr.—**Pileus** 1–2.5 cent. ( $\frac{1}{2}$ –1 in.), light yellow-rufescent behind, whitish at the margin, sessile with or without a vertical stem, then reflexed, horizontal, very irregularly shaped, lobed, delicately villous. **Gills** in the form of veins, *dichotomous*, narrow, *crisped*, very much swollen, edge very obtuse but not channelled, whitish or bluish-grey.

Substance and texture as described in the generic characters. Very gregarious, commonly imbricated. When young pezizoid or cup-shaped.

On logs, birch, &c. Uncommon. Oct.–Nov.

It occurs almost white. Spores elongated or cylindrical,  $4 \times 1$  mk. *K.* Name—*crispus*, crisped or curled. Fr. *Monogr.* ii. p. 244. *Hym. Eur.* p. 492. *C. Hbk.* n. 694. *S. Mycol. Scot.* n. 638. *Merulius Pers. Ic. descr.* t. 8. f. 7. *Cantharellus Fl. Dan.* t. 1759. *B. & Br.* n. 1135.

Schizophyl-  
lum.


GENUS XIX.—**Schizophyllum** ( $\sigma\chi\acute{\iota}\zeta\omega$ , to split;  $\phi\acute{\upsilon}\lambda\lambda\omicron\nu$ , a leaf).  
Fr. Obs. i. p. 103.

Pileus fleshless, arid. Gills coriaceous, fan-wise branched, united above by the tomentose pellicle, bifid, split longitudinally at the edge. Spores somewhat round, white. Fr. *Hym. Eur.* p. 492.

The two lips of the split edge of the gills are commonly revolute. The farthest removed of all the *Agaricini* from the type. Growing on wood.

1. **S. commune** Fr.—**Pileus** scarcely exceeding 2.5 cent. (1 in.),

very arid, pendulous, commonly extended behind into a stem-like base, entire or lobed, covered with whitish-grey down. **Gills** fuscous-grey then purplish, villous, revolute at the edge. Schizophyllum.



On dead wood, logs, &c. Rare in Britain.

White or greyish zoned, sometimes resupinate. *M. J. B.* Spores very small, almost globular, 3 mk. *W. G. S.* In Britain it is usually found and is common on foreign logs. In the 'Scottish Cryptogamic Flora' Greville records it as having been gathered on Dundas Hill, near Edinburgh. I have examined his specimen in the Edinburgh Herbarium marked "Dundas Hill." From the locality and from Greville's well-known accuracy, there can be no doubt that it was indigenous. Berkeley and Broome state that "undoubtedly indigenous specimens have occurred both in Buckinghamshire and Kent in 1878." Name—*communis*, common. Common in most countries. *Fr. Monogr.* ii. p. 244. *Hym. Eur.* p. 492. *Berk. Out.* p. 228. *B. & Br. n.* 1796\*. *C. Hbk. n.* 695. *S. Mycol. Scot. n.* 659. *Grev. t.* 61. *Krombh. t.* 4. f. 14-16. *Ag. Linn.*—*Batsch f.* 126. *Bull. t.* 346, 581. f. 1. *Sow. t.* 183.

LVIII. *Schizophyllum commune*.  
Natural size. Section ten times  
natural size.

## GENUS XX.—**Lenzites** (after Lenz, a German botanist).

Lenzites.

*Fr. Epicr.* p. 403.

Pileus corky or coriaceous, texture arid and floccose. Gills coriaceous, firm, sometimes simple and unequal, sometimes anastomosing and forming pores behind, trama floccose and similar to the pileus, edge somewhat acute. The European species are dimidiate, sessile, persistent, growing on wood, quite resembling *Dædalea*. *Fr. Hym. Eur.* p. 492.

Allied most nearly to *Trametes* and *Dædalea* and forming as it were the transition from *Agaricini* to *Polyporei*. In tropical countries they are more woody in texture.

\* *On wood of deciduous trees.*

1. **L. betulina** *Fr.*—**Pileus** 2.5–5 cent. (1–2 in.) broad, pale, grey-whitish, *corky-coriaceous*, firm and *rigid*, becoming plane, *tomentose*, commonly *obsoletely zoned*, zones sometimes darker; flesh floccose, white. **Gills** reaching the base, straight, some-


Lenzites.

times simple, sometimes branched and often anastomosing, about 6 mm. (3 lin.) broad, *dingy white, edge acute*.



Dimidiate, sessile. It varies solitary and caespitose-imbricated. The margin is of the same colour.

Stumps, logs, &c., chiefly birch. Common in England. Rare in Scotland. Perennial.



Deeply grooved concentrically. Often green with minute *Algæ*. Often quite resupinate, and then very deceptive. *M. f. B.* Name—*betula*, birch. *Fr. Monogr.* ii. p. 246. *Hym. Eur.* p. 493. *Berk. Out.* p. 228. *t.* 15. *f.* 3. *C. Hbk.* n. 696. *S. Mycol. Scot.* n. 660. *Ag. Linn.—Fl. Dan.* t. 1555. *B.* with darker gills. *Sow. t.* 182. *Schæff. t.* 57 is a handsome form.

LIX. *Lenzites betulina*. One-fourth natural size.

## 2. *L. flaccida* Fr.—**Pileus** at first

whitish, then dingy, *with zones wholly of the same colour*, coriaceous, thin, scarcely 2 mm. (1 lin.) thick, unequal, *easily bent, strigoso-hairy*. **Gills** thick, firm, straight, very broad, crowded, simple or branched, with shorter ones intermixed, at first shining white, then becoming pale.

The gills are never anastomosing, but here and there obfurcate, *i.e.*, towards the base. Commonly taken for *L. betulina*, but wholly different.

On stumps. Frequent in England.

Name—*flaccidus*, flaccid. *Fr. Monogr.* p. 246. *Hym. Eur.* p. 493. *Berk. Out.* p. 228. *C. Hbk.* n. 697. *Ag. Bull.* t. 394. *Bolt.* t. 158.

**\*\* On wood of trees with leaves like needles.**

3. *L. sæpiaria* Fr.—**Pileus** 2.5–7.5 cent. (1–3 in.) broad, *yellow-tawny when young* (remaining so at the margin when full grown), *date brown* when full grown, becoming black when old, corky-coriaceous, hard, convex becoming plane, sometimes orbicular, more frequently extended longitudinally, zoned, *strigoso-tomentose*, at length squamulose and pitted; *flesh tawny*. **Gills** extended to the base, very rigid and firm, branched, more or less anastomosing, 2–4 mm. (1–2 lin.) broad, *yellowish* when in vigour, but umber when old, the edge entire or slightly toothed.

Dimidiate, lateral. The gills are like those of *Dædalea*. It varies resupinate, forming orbicular placenta-like patches. There are many remarkable monstrous forms; occasionally the hymenium is wholly that of *Polyporus*.

On pine. Uncommon. Jan.–Dec.



Name—*sæpis*, a hedge, fence. Growing on old fences. *Fr. Monogr.* ii. p. Lenzites. 247. *Hym. Eur.* p. 494. *Berk. Out.* p. 228. *C. Hbk.* n. 698. *S. Mycol. Scot.* n. 661. *Dædalea*—*Schæff. t.* 76. *Ag. Sow. t.* 418. *Buxb. C. v. t.* 6.—*Vaill. t. i. f.* 1-3 (resupinate var.)

**4. L. abietina** Fr.—*Pileus* *umber-tomentose*, then becoming smooth, hoary, coriaceous, thin and comparatively *soft*, zoneless, *effuso-reflexed*; flesh very thin, of the same colour as the pileus. **Gills** decurrent in the effused base, distant, simple, unequal, not anastomosing, but here and there interrupted, or torn into teeth, *becoming glaucous with dense pruina*.

Nearly always lengthened out even to as much as 30 cent. (12 in.), but always remaining narrow, about 12 mm. ( $\frac{1}{2}$  in.) broad. *Effuso-reflexed*, but sometimes remaining resupinate.

Chiefly on dressed fir. Rare. Glasgow. Perennial.

Perhaps imported. Name—*abies*, fir. *Fr. Monogr.* ii. p. 247. *Hym. Eur.* p. 495. *Berk. Out.* p. 229. *C. Hbk.* n. 699. *S. Mycol. Scot.* n. 662. *Ag. Bull. t.* 442. f. 2, 541. f. 1. *Ventur. t.* 60. f. 3-5.

## ORDER II.—POLYPOREI.

Hymenophore inferior, facing the ground; hymenium consisting of tubes with poriform mouths which are round or angular, sometimes sinuous or torn, lined within with 4-spored sporophores and cystidia.

*Fleshy, coriaceous, or woody fungi, most abundant and luxuriant in warm countries. Intermediate between the Agaricini and the Hydnei, connected with the former by Dædalea and Lenzites, and with the latter by Fistulina and Irpex.*

Boletus.

GENUS XXI.—**Boletus**. Dill. Fr. Obs. i. p. 109. (The name of a fungus considered a great delicacy among the Romans, derived from βῶλος, a clod, probably to denote the round figure of the plant.)

Hymenium wholly composed of small tubes, connected together in a stratum the surface of which is dotted with their poriform mouths, and which is distinct from the hymenophore on account of the latter not descending into a trama. Tubes packed close together, easily separating from the hymenophore and from one another. Pores or mouths of the tubes round or angular (in the subgenus *Gyrodon* sinuous or gyroso-plicate). Spores normally fusiform, rarely oval or somewhat round. *Growing on the ground, fleshy, putrescent, with central stems. Mostly edible, and of importance as articles of food; a few poisonous.*



LX. *Boletus edulis*. One-fourth natural size.

A well-marked and sharply defined genus, although some of the species depart so much from the type that they have been referred to distinct genera. *Fr. Hym. Eur. p. 495. Compare Epicr. p. 408 and Syst. Myc. i. p. 385.*

The colour of the spores of many of the species is unknown,

and where it is known, it is found to differ in closely allied Boletus. species; and hence in arranging them it is necessary to take the colour of the tubes (and not as in the *Agaricini* the colour of the spores) as the basis of the primary groups. *Fr. Hym. Eur. p.* 496, &c.

*Ser. I. EUCHROI.* (εὖ, well; χρῶς, colour). Tubes bright-coloured, commonly yellow, not white or grey.

A. Viscipelles (*viscum*, bird-lime; *pellis*, a skin). *Pileus covered over with a viscous pellicle.* Stem solid, not bulbous, nor reticulated with veins. Tubes adnate to the stem behind, rarely sinuate, of one colour. Spores of such as are sufficiently well known yellowish.

\* *Genuini*, forming a continuous series.

\*\* *Departing from the type, of doubtful affinity.*

B. Subtomentosi (*sub*, tomentum, down). *Pileus destitute of a viscid pellicle, villous when young, rarely at length becoming smooth.* Stem at the first extended, not bulbous, nor reticulated with veins, though here and there rugose or striate. Flesh rarely changing colour. Tubes of one colour, adnate.

C. Subpruinosi (*sub*, pruina, hoar-frost). Tubes adnate to the stem, yellowish. Stem equal, even, not bulbous nor reticulated. Pileus smooth, but most frequently pruinose.

D. Calopodes (καλός, beautiful; πούς, a foot). Stem stout, at the first bulbous, in typical species reticulated with veins. Tubes adnate, pores not reddish.

\* *Genuini. Stem reticulated with veins, commonly of a beautiful red.*

\*\* *Departing from the type, of uncertain affinity. Stem clavate, even.*

E. Edules (*edulis*, edible). *Stratum of tubes, which at the first are not reddish at the orifice, but have usually a white stuffing, rounded and depressed about the stem, somewhat free.* Stem stout, bulbous, resembling that of the *Luridi*, only (except in *B. edulis*) it is neither reticulated, nor dotted with small scales, nor red. Flesh scarcely changing colour, taste pleasant. Pre-eminent for their esculent qualities.

F. Luridi (*luridus*, lurid). *Stratum of tubes rounded towards the stem and free; pores at the first stopped up, red.* Pileus compact, then soft, pulvinate; flesh juicy, changing colour. Stem stout, at first curt, bulb-shaped, then elongated and nearly equal, somewhat reticulated or dotted. Growing chiefly in deciduous woods. Very poisonous.

*Ser. II. TEPHROLEUCI* (τεφρός, ash-coloured; λευκός, white). Tubes at first white or grey.

G. Favosi (*favus*, honeycomb. From the size and shape of the tubes). Tubes large, angular, unequal, adnate to the stem, but often shortened around it, not however forming a rounded-free stratum. Spores, as far as known, brown.

H. Versipelles (*verto*, to change; *pellis*, a skin. From the typical species *B. versipellis*). Tubes minute, round, equal, crowded together in a convex stratum, which is free from the stem. Spores ferruginous.

I. Hyperhodii (ὑπό, somewhat; ῥόδον, a rose. From the colour of the spores). Tubes adnate to the stem, whitish; spores rosy then white-flesh-colour.

K. Cariosi (*caries*, rottenness). Stem externally never reticulated, internally stuffed with a spongy pith, at length commonly hollowed out. Tubes

**Boletus.** at the first white, then often light yellowish; pores minute, round. Spores white.

*Subg. I. GYRODON* Opatowski (γυρός, a circle; ὀδός, a tooth). Pores sinuous or gyroso-plicate; tubes very short, scarcely 2 mm. (1 lin.) long, slightly adnate to the hymenophore.

*Subg. II. BOLETINUS* Kalchbr. Enum. p. 286. (From *Boletus* to denote its relation to that genus.) "Hymenophore not even as in the true *Boleti*, but with projecting points which descend like a trama between the tubes." Stem annulate.

**Ser. I. EUCHROI.** Tubes bright-coloured, &c.

**A. Viscipelles.** *Pileus covered over with a viscous pellicle, &c.*

\* *Genuini*, forming a continuous series.

**1. B. luteus** Linn.—**Pileus** about 5–10 cent. (2–4 in.) broad, yellowish, *smear'd with fuscous separating gluten*, gibbous then pulvinate; flesh white. **Stem** about 5 cent. (2 in.) long, 12 mm. (½ in.) and more thick, equal, firm, whitish, *rough with dots* and light yellowish *above the large, membranaceous, white-fuscous ring*. **Tubes** adnate; pores minute, simple, yellow.

The pileus becomes paler and virgate-spotted when the gluten separates.

In pine woods. Common. July–Nov.

Edible and highly esteemed like other Boleti. The flesh is very tender. The tubes should be scraped away from all the species before being cooked. Spores spindle-shaped, yellowish-brown, 8 × 3 mk. *W.G.S.*; 6–8 × 3 mk. *K.* Name—*luteus*, yellow. *Linn.*—*Fr. Hym. Eur. p. 497.* *Syst. Myc. i. p. 386.* *Sv. ätl. Sv. t. 22.* *Berk. Out. p. 229.* *C. Hbk. n. 700.* *S. Mycol. Scot. n. 663.* *Scheff. t. 114.* *Fl. Dan. t. 1135.* *Rostk. Sturm. 4. t. 1.* *Kl. Fl. Bor. t. 377.* *Barla t. 31. f. 1–3.* *Harz. t. 6.* *Gonn. & Rab. vii. t. 6. f. 2 and t. 4.* —*Krombh. t. 33.*

**2. B. elegans** Schum.—**Pileus** golden or inclining to ferruginous, convexo-plane, viscous; flesh light yellowish. **Stem** firm, unequal, golden then rufescent, dotted *above the white then light yellowish*, fugacious ring. **Tubes** decurrent; pores minute, simple, golden-sulphur-yellow.

Gills at length darker.

In woods, especially larch. Common. June–Oct.

About the size of *B. luteus*. Name—*elegans*, elegant. *Schum.*—*Fr. Hym. Eur. p. 497.* *Sv. ätl. Sv. t. 76.* *Berk. Out. p. 229.* *C. Hbk. n. 701.* *S. Mycol. Scot. n. 664.* *Hussey ii. t. 12.* *Gonn. & Rab. t. 5. f. 2.* *Price f. 110.* *Grev. t. 183.* *Ventur. t. 47. f. 1, 2.* *Bull. t. 332.* *Krombh. t. 34. f. 1–10.*

**3. B. flavus** With.—**Pileus** light yellow with fuscous separating

gluten, pulvinate, compact. **Stem** yellow becoming fuscous, Boletus. *reticulated above the dirty light yellow*, membranaceous, fugacious *ring*. **Tubes** adnate; pores *angular*, yellow.

Firm. Distinguished from *B. luteus*, which is like it, by the larger angular pores.

In mixed woods. Common. June–Oct.

Pileus 5–7.5 cent. (2–3 in.) broad. Stem 5–7.5 cent. (2–3 in.) long, 12–18 mm. (6–9 lin.) thick. Spores spindle-shaped, yellowish-brown,  $8 \times 4$  mk. *W.G.S.* Name—*flavus*, light yellow. *With.*—*Fr. Hym. Eur. p. 497. Berk. Out. p. 230. C. Hbk. n. 702. S. Mycol. Scot. n. 665. Bolt. t. 169. Sow. t. 265. Kl. Linn. vii. t. 198.*

**4. B. flavidus** Fr.—Pileus 2.5–5 cent. (1–2 in.) broad, livid-light-yellowish, gibbous then plane, viscous; flesh pallid. **Stem** 5–7.5 cent. (2–3 in.) long, thin, 4–6 mm. (2–3 lin.) thick, somewhat equal, pallid, *sprinkled with fugacious glandules above the entirely viscous ring*. **Tubes** decurrent; pores large, angular, compound, dirty light yellow.

*Thin.* Quite distinct from its allies, more slender, and differing from all of them in the veil, which is entirely glutinous.

In pine woods. Rannoch. Sept.

Spores elongato-ellipsoid, straight, subhyaline,  $8-10 \times 3.5-4$  mk. *K.* Name—*flavidus*, light yellowish. *Fr. Hym. Eur. p. 498. Syst. Myc. i. p. 387. S. Mycol. Scot. n. 666.—Pers. Myc. Eur. 2. t. 20. f. 1-3. Krombh. t. 4. f. 35-37.*

**5. B. collinitus** Fr.—Pileus becoming pale *when the fuscous gluten separates*, pulvinate, even; flesh white. **Stem** firm, attenuated downwards, *ringless*, white becoming fuscous, *somewhat reticulated with adpressed squamules*. **Tubes** adnate, elongated; pores *divided in two*, pallid then yellow, naked.

It resembles *B. luteus* in stature and colour.

In fir woods. Ascot. Nov.

Name—*collino*, to besmear. From the gluten. *Fr. Hym. Eur. p. 498. B. & Br. n. 1284.—Krombh. t. 76. f. 10, 11?*

**6. B. granulatus** Linn.—Pileus 5–7.5 cent. (2–3 in.) broad, *fuscous-ferruginous with the gluten with which it is smeared*, yellowish when the gluten separates, convexo-expanded; flesh light yellowish. **Stem** 5 cent. (2 in.) and more long, 6–12 mm. (3–6 lin.) thick, *ringless*, light yellowish, *dotted with granules upwards*. **Tubes** adnate, short; pores *simple, granulated*.

Commonly gregarious. The stem is often attenuated upwards.

Boletus.

In woods. Rare. July–Sept.

Orifices of tubes at first dripping with a milky fluid. *M. F. B.* This dries in the form of sugary granules. Edible. Spores spindle-shaped, yellowish-orange,  $8 \times 3$  mk. *W. G. S.*; oblong,  $8-10 \times 3$  mk. *K.* Name—*granula*, a granule. Granulated. *Linn.*—*Fr. Hym. Eur.* p. 498. *Syst. Myc.* i. p. 387. *Sv. atl. Sv. t.* 23. *Berk. Out. p.* 230. *C. Hbk. n.* 704. *S. Mycol. Scot. n.* 667. *Lenz f.* 31. *Letell. t.* 604. *Barla t.* 31. *f.* 4-12. *Rostk. t.* 3. *Gonn. & Rab. vii. t.* 6. *f.* 1.—*Schaff. t.* 123. *Krombh. t.* 34. *f.* 11-14. *Sow. t.* 420.

**7. B. bovinus** Linn.—**Pileus** 5 cent. (2 in.) broad, pale yellowish, rather plane, smooth, viscid; flesh white. **Stem** 5 cent. (2 in.) long, equal, even, unicolorous. **Tubes** somewhat decurrent; pores *compound*, *grey-light-yellow* becoming ferruginous.

The gluten on the pileus, which is at length repand, is thin, not of a different colour and not separating. Somewhat caespitose.

In woods, chiefly pine. Uncommon. Sept.–Oct.

Tubes resembling the pores of *Merulius lachrymans*, very shallow ( $\frac{1}{2}$  in.) Smell like *Marasmius oreades*. *M. F. B.* Spores spindle-shaped, dingy green-ochre,  $8 \times 4$  mk. *W. G. S.*; oblong-ellipsoid,  $8-10 \times 3$  mk. *K.* Name—*bos*, an ox. Perhaps from its gregarious habit; the synonym of 'Fl. Dan.' is *B. gregarius*. It might also refer to colour as *vaccinus* does. *Linn.*—*Fr. Hym. Eur. p.* 499. *Syst. Myc. i. p.* 388. *Berk. Out. p.* 230. *C. Hbk. n.* 705. *S. Mycol. Scot. n.* 668. *Lenz f.* 38 (old). *Krombh. t.* 75. *f.* 1-6. *Kl. Bor. t.* 378. *Gonn. & Rab. vii. t.* 2. *f.* 1. *Hussey i. t.* 34.—*Fl. Dan. t.* 1018.

**8. B. badius** Fr.—**Pileus** 7.5–15 cent. (3–6 in.) broad, *bay-brown-tawny*, pulvinate, soft, viscous; flesh 2.5–5 cent. (1–2 in.) thick, becoming azure-blue at the tubes. **Stem** 7.5 cent. (3 in.) long, 2.5 cent. (1 in.) thick, solid, somewhat equal, even, paler, *brown-pruinata*. **Tubes** long, adnate, sinuato-depressed; pores rather large, dingy, *white-light-yellow becoming green*, angular.

The stem is attenuated, sometimes upwards, sometimes downwards, but never bulbous. Pileus commonly glutinous, but shining when dry, or when young as if villous-viscous, always even. Very distinct.

In woods, chiefly pine. Frequent. Aug.–Oct.

Spores fusoid-oblong, light yellow,  $15-20 \times 5-6$  mk. *K.* Name—*badius*, bay-brown. *Fr. Hym. Eur. p.* 499. *Elench. p.* 126. *Sv. atl. Sv. t.* 50. *Berk. Out. p.* 231. *C. Hbk. n.* 706. *S. Mycol. Scot. n.* 669. *Rostk. t.* 5. *Fl. Bat. t.* 804. *Kl. Bor. t.* 379. *Lenz f.* 35.—*Krombh. t.* 36. *f.* 12-18.

**9. B. sanguineus** With.—**Pileus** *blood-red*, convexo-plane, even, smooth, *viscid*. **Stem** equal, even, *variegated yellow and blood-red*. **Tubes** adnate, orange-yellow; pores large, unequal.

Colour remarkably glistening. Easily confounded with forms of *B. chrysenteron*.

In woods. Rare.

**Pileus** 18 mm.-3 cent. ( $\frac{3}{4}$ -1 $\frac{1}{4}$ ) in. broad, when old nearly 7.5 cent. (3 in.) **Boletus**. Stem 2.5-6 cent. (1-2 $\frac{1}{2}$  in.) long, nearly 9 mm. ( $\frac{3}{8}$  in.) thick. Pileus when old rich brown. Flesh white, a little tinged with crimson next to the skin, changing slowly to a bluish cast when wounded. In the larger specimens the base is bulbous. *With.* Name—*sanguis*, blood. Blood-red. *With.*—*Fr. Hym. Eur. p. 500. Berk. Out. p. 231. C. Hbk. n. 707. Sow. t. 225. Leveill. in Paul. t. 181. f. 3, 4.*

**10. B. piperatus** Bull.—**Pileus** 2.5-7.5 cent. (1-3 in.) broad, yellow to pale yellowish, convexo-plane, smooth, slightly viscid; flesh yellowish. **Stem** 4 cent. (1 $\frac{1}{2}$  in.) long, thin, 6-8 mm. (3-4 lin.) thick, even, fragile, of the same colour as the pileus, containing yellow milk at the base. **Tubes** decurrent, ferruginous; pores large, angular.

Smaller than the rest, occasionally rimoso-squamulose. Easily distinguished by its peppery taste.

In woods. Frequent. Aug.—Nov.

Reckoned poisonous. Spores oval, brown, 8×4 mk. *W.G.S.*; ellipsoid-oblong, yellowish, 8-9×3-4 mk. *K.* Name—*piper*, pepper. From the taste. *Bull. t. 451. f. 2. Fr. Hym. Eur. p. 500. Syst. Myc. i. p. 388. Sv. atl. Sv. t. 67. Berk. Out. p. 231. C. Hbk. n. 708. S. Mycol. Scot. n. 670. Sow. t. 34. Fl. Dan. t. 1850. f. 2. Krombh. t. 37. f. 16-20. Rostk. t. 6. Barla t. 32. f. 5-10. Corda Sturm xi. t. 60.—Batsch t. 128.*

**\*\* Departing from the type, of doubtful affinity.**

No British species.

B. Subtomentosi. *Pileus destitute of a viscid pellicle, &c.*

**11. B. variegatus** Sw.—**Pileus** 7.5-12.5 cent. (3-5 in.) broad, dark yellow, convex then plane, obtuse, slightly moist, sprinkled with superficial, fasciculate-hairy squamules, the acute margin at the first flocculose; flesh yellow, here and there becoming azure-blue. **Stem** 5-7.5 cent. (2-3 in.) long, 12 mm. ( $\frac{1}{2}$  in.) thick, firm, equal, even, dark yellow, sometimes reddish. **Tubes** adnate, unequal, pores minute, brown then cinnamon.

The scales on the pileus are separating.

In woods, chiefly pine. Frequent. Sept.—Oct.

Smell unpleasant, taste not so. *M.F.B.* Spores oval, greenish-ochre, 3×2 mk. *W.G.S.*; oblong-ellipsoid, hyaline or very pale yellow, 8-11×3-4 mk. *K.* Name—*variegatus*. Of the different colours. *Swartz.*—*Fr. Hym. Eur. p. 501. Syst. Myc. i. p. 388. Sv. atl. Sv. t. 66. Berk. Out. p. 231. C. Hbk. n. 711. S. Mycol. Scot. n. 671. Lenz f. 39. Krombh. t. 34. f. 15-18, and t. 75. f. 7-14. Harz. t. 15. Rostk. t. 16.—B. Golden. Schaff. t. 115.*

**12. B. sulphureus** Fr.—**Pileus** sulphur-yellow, compact, convex then plane, silky-tomentose with innate flocci; flesh light

Boletus.

yellow, becoming more or less azure-blue when broken, but golden when exposed to the air, here and there reddish under the tubes. **Stem** 2.5–5 cent. (1–2 in.) long, 2.5 cent. (1 in.) thick, firm, ventricose, *even*, smooth, *sulphur-yellow*, at length becoming dingy ferruginous. **Tubes** adnato-decurrent, short, 2–4 mm. (1–2 lin.) long, adhering more closely than usual, changeable on being touched, at length with ferruginous spots; pores minute, compound, sulphur-yellow, at length becoming green.

Springing from a broadly expanded, golden, woolly mycelium, it is very caespitose, with the appearance of *Ag. spectabilis*.

Among sawdust. Loch-an-Eilan, Rothiemurchus. Sept.

Spores light yellow then ochraceous. *Fr.* Name—*sulphur*, brimstone. Sulphur-yellow. *Fr. Hym. Eur. p.* 502. *B. & Br. n.* 1424. *S. Mycol. Scot. n.* 672.

**13. B. striæpes** Secr.—**Pileus** olivaceous, *the cuticle ferruginous within*, convex then plane, soft, silky. **Stem** firm, curved, yellow, with fuscous-black *striae*, fuscous-rufescent at the base. **Tubes** adnate, angular, greenish; pores minute, yellow.

Flesh white, yellow at the tubes, and at the base of the stem which is externally of the same colour and velvety.

In woods. Rare.

Flesh sparingly changing to blue. *M. f. B.* Name—*stria*, a line; *pes*, a foot. From the striate stem. *Secr.*—*Fr. Hym. Eur. p.* 502. *Berk. Out. p.* 232. *C. Hbk. n.* 712. *Battar. t.* 29. *C.*

**14. B. chrysenteron** Fr.—**Pileus** fuscous or somewhat brick-colour, convexo-plane, soft, *floccoso-squamulose*; *flesh yellow, red beneath the cuticle*. **Stem** somewhat equal, rigid, fibroso-striate, scarlet or light yellow. **Tubes** somewhat adnate; pores rather large, angular, unequal, greenish-yellow.

The pileus is often cracked into patches, and then the cracks are scarlet. Varying in stature like its allies, with the pileus becoming smooth and more brightly coloured, with the flesh becoming somewhat azure-blue, and with the tubes depressed round the stem.

In woods and open ground. Common. July–Nov.

Edible. Pileus 5–7.5 cent. (2–3 in.) or more broad. Stem 7.5 cent. (3 in.) long, 6–12 mm. ( $\frac{1}{4}$ – $\frac{1}{2}$  in.) thick, easily distinguished by the red cracks. Spores spindle-shaped, pale warm brown, 14–2 mk. *W. G. S.* Name—*χρυσός*, gold; *έντερον*, inward parts. From its yellow flesh. *Fr. Hym. Eur. p.* 502. *Berk. Out. p.* 232. *C. Hbk. n.* 713. *S. Mycol. Scot. n.* 673. *Bull. t.* 490. *f.* 3. *Hussey i. t.* 5. *Quel. t.* 16. *f.* 4.

**15. B. subtomentosus** Linn.—**Pileus** somewhat olivaceous, *of the same colour beneath the cuticle*, pulvinato-expanded, soft, dry,



*villosa-tomentose*. **Stem** stout, attenuated downwards, *somewhat* Boletus.  
*ribbed-sulcate*, slightly rough with dots under a lens. **Tubes**  
adnate; *pores large, angular, yellow*.

Flesh white or pallid, not red beneath the cuticle. When the pileus is cracked the *interstices become yellow*. The colour of the pileus is brighter when the olivaceous down disappears.

In woods. Common. Aug.—Nov.

About same size as *B. chrysenteron*. Often growing on beech-nuts. *M. J. B.* Easily distinguished by its yellow cracks. Spores oval, yellowish-brown,  $14 \times 5$  mk. *W. G. S.*; fusoid-oblond, hyaline or very pale yellow,  $11-13 \times 4-5$  mk. *K.* Name—*sub*, and *tomentosus*, downy. *Linn.*—*Fr. Hym. Eur. p. 503.* *Berk. Out. p. 232.* *C. Hbk. n. 714.* *S. Mycol. Scot. n. 674.* *Price f. 2.* *Vivian. t. 37.* *Gonn. & Rab. vii. t. 5. f. 1.* *Bolt. t. 84. middle fig.*—*Schæff. t. 112.* *Krombh. t. 37. f. 8-11.* *Bull. t. 393.* *Fl. Dan. t. 1074.* *Battar. t. 30. f. F.* *Krombh. t. 48. f. 1-6* represents a beautiful var., somewhat æruginous.

**16. B. spadiceus** Schæff.—**Pileus** date-brown, opaque, pulvinato-expanded, moderately compact, dry, tomentose, then widely cracked; the white flesh unchangeable, fuscous reddish above. **Stem** firm, clavate, *even, flocculoso-furfuraceous*, yellow inclining to fuscous. **Tubes** adnate, yellow; pores minute, somewhat round.

The stem is compact, not reddish, yellowish-white internally.

In mixed woods. Glamis, 1875. Aug.

Pileus 7.5–10 cent. (3–4 in.) broad. Stem very thick. Name—*spadix*, a palm-branch. Date-brown. *Schæff. t. 126.* *Fr. Hym. Eur. p. 503.* *B. & Br. n. 1797.* *S. Mycol. Scot. n. 675.*—*Krombh. t. 36. f. 19, 20.*

**17. B. radicans** Pers.—**Pileus** olivaceous-cinereous then pale yellowish, pulvinate, dry, somewhat tomentose, the thin margin involute; flesh light yellow, becoming immediately dark azure-blue. **Stem** *attenuato-rooted*, even, light yellow, *flocculose with reddish pruina*, naked and dark when touched. **Tubes** adnate; pores unequal, large, lemon-yellow.

Taste bitterish. The pores are those of *B. subtomentosus*, the habit that of *B. badius*.

In woods. Epping Forest. Staplehurst, &c.

Spores spindle-shaped, very pale ochre, almost white,  $6 \times 3$  mk. *W. G. S.* Name—*radix*, a root. Rooting. *Pers.*—*Fr. Hym. Eur. p. 503.* *Journ. Bot. 1873, p. 337.* *C. Hbk. n. 714.* *Opatowski Bolet. t. 1.*

**18. B. rubinus** Smith.—**Pileus** 5–7.5 cent. (2–3 in.) broad, yellow-fuscous, pulvinate-gibbous then plane, dry, somewhat tomentose, slightly cracked; flesh vivid yellow, wholly unchange-

Boletus.

able. **Stem** 5-7.5 cent. (2-3 in.) long, yellow, smeared with crimson, irregular. **Tubes** somewhat decurrent, compound, of medium size, *wholly* carmine.

It differs from all other British species in the wholly carmine tubes, together with the vivid yellow, wholly unchangeable flesh.

In mixed woods. Rare. Aug.-Sept.

Spores oval, almost round, pale warm brown,  $6 \times 5$  mk. *W.G.S.* Name—*ruber*, red. From the tubes. *Smith Seem. Journ.* 1868, p. 33. t. 75. f. 1-4. *Fr. Hym. Eur.* p. 504. *C. Hbk. n.* 709. *S. Mycol. Scot. n.* 676.

C. Subpruinosi. Pileus smooth, but most frequently pruinose, &c.

**19. B. pruinatus** Fr.—**Pileus** 5 cent. (2 in.) and more broad, *purplish-bay-brown, umber-pruinose*, convex then plane, rigid, *dry*; flesh white, but also obsoletely green or azure-bluish. **Stem** 5 cent. (2 in.) long, firm, somewhat ventricose, even, smooth, *variegated yellow and reddish*. **Tubes** adnate, light yellow; pores minute, round.

On grassy ground. Kew, &c.

Name—*pruina*, hoar-frost. From the pruinose pileus. *Fr. Hym. Eur.* p. 504. *Épicr. p.* 414. *B. & Br. n.* 1285. *Schaff. t.* 133?—*Bull. t.* 393. f. B. C.

**20. B. parasiticus** Bull.—**Pileus** 2.5-5 cent. (1-2 in.) broad, dingy yellow, convex then plane, *silky becoming smooth*, dry, *soon cracked in a tessellated manner*. **Stem** 5 cent. (2 in.) long, thin, rigid, incurved, externally and internally yellow. **Tubes** decurrent, of medium size, compound, golden.

The colour of the pileus is variable.

On species of *Scleroderma*. Rare. Clifton. Kew.

Tubes labyrinthiform. Not viscid in any stage of growth. *B. & Br.* Spores spindle-shaped, elongated, pale brown,  $14 \times 4$  mk. *W.G.S.* Name—*parasitus*, a guest. *Parasitic. Bull. t.* 451. f. 1. *Fr. Hym. Eur. p.* 505. *Syst. Myc. i. p.* 389. *Berk. Out. p.* 231. *C. Hbk. n.* 710. *Saund. & Sm. t.* 43. *Rostk. t.* 7 (monstrous).

D. Calopodes. Stem stout, at the first bulbous, &c.

\* Genuini. *Stem reticulated with veins, &c.*

**21. B. variegator** B. & Br.—**Pileus** olivaceous, convex, somewhat tomentose, *margin involute*; flesh dark purple under the cuticle. **Stem** bulbous, attenuated upwards, reticulated at the

apex, yellowish downwards, rufescent and delicately pubescent *Boletus* upwards. **Tubes** minute, free, yellow.

The flesh of the pileus and stem is pale, here and there inclining to yellow, and partially marbled. Approaching *B. subtomentosus* in habit, but with the bulbous reticulated stem of *Calopodes*.

In woods, &c. Uncommon. Aug.—Sept.

Name—*varius* and *color*. From the various colours. *B. & Br. n. 1020. t. 13. f. 3. Fr. Hym. Eur. p. 506. C. Hbk. n. 715. S. Mycol. Scot. n. 677.*

**22. B. calopus** Fr.—**Pileus** *olivaceous*, globose then pulvinate, unpolished, *somewhat tomentose*; flesh pallid, becoming somewhat azure-blue. **Stem** firm, conical then somewhat equal, reticulated, *scarlet*, throughout or *at the apex*. **Tubes** adnate; pores minute, angular, yellow.

Stem at length of the same colour as the pileus downwards and elongated. Pileus not involute, somewhat repand.

In mixed woods. Frequent. July—Sept.

Spores spindle-shaped, yellowish-brown,  $8 \times 3$  mk. *W.G.S.* Names—*καλός*, beautiful; *πούς*, a foot. With beautiful stem. *Fr. Hym. Eur. p. 506. Sv. atl. Sv. t. 69. Berk. Out. p. 232. C. Hbk. n. 716. S. Mycol. Scot. n. 678. Krombh. t. 37. f. 1-7. Rostk. t. 27. Harz. t. 69. Saund. & Sm. t. 14. —Schæff. t. 315.*

**23. B. olivaceus** Schæff.—**Pileus** *olivaceous-fuscous*, convex, even, becoming smooth, *margin at first inflexed*; flesh azure blue, then white. **Stem** firm, clavate, bulbous, red, *yellowish at the apex*, with blood-red reticulations and dots. **Tubes** adnate, curt; pores minute, unequal, light yellow-olivaceous.

Stature curt, robust.

In beech woods. Uncommon. Sept.

Pileus 5-7.5 cent. (2-3 in.) and more broad. Stem short. Name—*oliva*, an olive. Olivaceous in colour. *Schæff. t. 105. Fr. Hym. Eur. p. 506. Berk. Out. p. 233. C. Hbk. n. 717. S. Mycol. Scot. n. 679. Ventur. t. 36. f. 3, 4. —Rostk. t. 32.*

**24. B. pachypus** Fr.—**Pileus** 10-20 cent. (4-8 in.) broad, fuscous then pallid-tan, pulvinate, somewhat tomentose; flesh thick, whitish, becoming slightly azure-blue. **Stem** 5-12.5 cent. (2-5 in.) long, always thick, firm, curt, ovato-bulbous, then elongated, equal, reticulated, *variegated light yellow and red*, often wholly intensely blood-red. **Tubes** somewhat elongated, *shortened round the stem*; pores round, light yellow, at length somewhat green.

Larger than preceding species, handsomer, approaching *Edules* in habit.

In woods, chiefly pine. Uncommon. Sept.—Oct.

Boletus.

Monstrous forms of *B. edulis* are often mistaken for it. Spores oval, yellowish ochre,  $14 \times 5$  mk. *W.G.S.* Name—*παχύς*, thick; *πούς*, a foot. Thick-stemmed. *Fr. Hym. Eur. p.* 506. *Syst. Myc. i. p.* 390. *Sv. ätl. Sv. t.* 68. *Berk. Out. p.* 233. *C. Hbk. n.* 718. *S. Mycol. Scot. n.* 680. *Lenz t.* 60. *Krombh. t.* 35. *f.* 13-15.—*Ventur. t.* 64. *f.* 1, 2. *B. pachypus Krombh. t.* 35. *f.* 10-12. *Saund. & Sm. t.* 17 (wholly white) is perhaps a new species.

*\*\* Departing from the type, of uncertain affinity, &c.*

No British species.

E. Edules. *Stratum of tubes rounded and depressed about the stem, &c.*

**25. B. edulis Bull.**—*Pileus somewhat brown, pulvinate, smooth, moist; flesh white, reddish beneath the cuticle, compact then softer. Stem stout, reticulated, pallid-fusca. Tubes semi-free, elongated, minute, at first white, then yellow and greenish.*

The pileus varies white, grey-rufescent, &c. The stem is sometimes obsoletely reticulated, sometimes curt and bulbous, sometimes elongated and equal, also rooting.

In woods, especially beech. Common. July–Nov.

Pileus 15 cent. (6 in.) and more broad. Stem 10 cent. (4 in.) or more long. 5 cent. (2 in.) thick. Spores fusoid-obl., bright yellowish,  $14-16 \times 4-5.5$  mk. *K.* Name—*edulis*, edible. Reckoned one of the most delicious, but the flesh is rather soft when cooked. *Bull. t.* 60, 494. *Fr. Hym. Eur. p.* 508. *Sv. ätl. Sv. t.* 13. *Berk. Out. p.* 234. *C. Hbk. n.* 719. *S. Mycol. Scot. n.* 681. *Sow. t.* 111. *Sv. Bot. t.* 197. *Lenz f.* 34. *Tratt. Austr. f.* 34. *Krombh. t.* 31. *Vittad. t.* 22. *Ventur. t.* 8. *Vivian. t.* 25. *Barla t.* 34. *Hussey i. t.* 81. *Badh. i. t.* 3. *ii. t.* 3. *f.* 1, 2. *Price t.* 63. *Hogg & Johnst. t.* 11. *Harz. t.* 40, 41. *Gonn. & Rab. vii. t.* 1.—*Schæff. t.* 134. *Rostk. t.* 37.

**26. B. æreus Bull.**—*Pileus olivaceous-fusca, somewhat blackish, pulvinate, smooth, somewhat pelliculose; flesh white, becoming yellow on exposure to the air. Stem stout, somewhat reticulated, yellowish, becoming fuscous at the base. Tubes minute, somewhat free, sulphur-yellow.*

Pileus small. The stem when young is oblongate not bulbous.

In woods. Surrey.

Spores oblong, oblique at the base,  $11-14 \times 6$  mk. *B. & Br.* Name—*æs*, copper. From the colour. *Bull. t.* 321. *Fr. Hym. Eur. p.* 508. *B. & Br. n.* 1425. *Krombh. t.* 36. *f.* 1-7. *Quel. t.* 16. *f.* 2.—*Rostk. t.* 15. *Bull. t.* 385 var.

**27. B. fragrans Vitt.**—*Pileus fuscous-umber, pulvinate, repand, somewhat tomentose, margin inflexed; flesh yellow, sometimes unchangeable, sometimes becoming green or azure-blue, at length becoming red. Stem stout, at first ovato-bulbous, often fusiform*

at the base, *even, variegated yellowish and red*. **Tubes** semi-Boletus. free; pores minute, round, yellow, becoming green.

The pileus varies unequal, and here and there spotted or margined with purple.

In woods, under oak, &c. Sept.—Oct.

Spores pale yellowish-green, ovate, with an apiculus at one end,  $11 \times 4$  mk. *W.G.S.*; elongato-pruniform, nucleate, 10 mk. *Q.* Edible. Name—*fragrans*, fragrant. *Vittad. t.* 19. *Fr. Hym. Eur. p.* 509. *C. Hbk. n.* 720. *Ventur. t.* 53. *f.* 3-5.—*Krombh. t.* 75. *f.* 15-21.

**28. B. impolitus** Fr.—**Pileus** *tawny-brown*, pulvinate, dilated, *flocculose*, at length granuloso-rivulose, unpolished; flesh thick, light yellowish beneath the cuticle. **Stem** stout, somewhat bulbous, *even, pubescent*, light yellow. **Tubes** free; pores minute, yellow.

Handsome, obese, pleasant in odour and taste. Sometimes there is a reddish zone near the apex of the stem.

In mixed woods. Rare. Sept.

Pileus 10 cent. (4 in.) or more broad. Stem about 5 cent. (2 in.) long. Spores oval or spindle-shaped, pale greenish-brown,  $14 \times 5$  mk. *W.G.S.*; ellipsoid, papillate, 16-18 mk. *Q.* Edible; reckoned among the most delicious. Name—*impolitus*, unpolished. *Fr. Hym. Eur. p.* 509. *Sv. atl. Sv. t.* 42. *Berk. Out. p.* 234. *C. Hbk. n.* 721. *S. Mycol. Scot. n.* 682.—*Harz. t.* 51. *B. pileus areolato-rimose. Schæff. t.* 108. *Letell. t.* 614. *Rostk. t.* 36.

**29. B. æstivalis** Fr.—**Pileus** *whitish*, pulvinate, somewhat repand, *even, smooth*, granular in dry weather; flesh yellow beneath, white above, reddish in the base of the stem. **Stem** very thick, bulbous, *even, smooth, light yellow*. **Tubes** somewhat free; pores equal, minute, yellow.

Stature almost that of *B. impolitus*, equally pleasant, tasty.

In woodland pastures. Uncommon.

Pileus sometimes 15-20 cent. (6-8 in.) broad. Stem 5 cent. (2 in.) thick. Spores elongated oval, greenish-brown, rather dark,  $12 \times 4$  mk. *W.G.S.* Edible. Name—*æstivalis*, pertaining to summer. *Fr. Hym. Eur. p.* 510. *Sv. atl. Sv. t.* 43. *Berk. Out. p.* 234. *C. Hbk. n.* 722. *Hussey i. t.* 25. *Hogg & Johnst. t.* 13.

**F. Luridi.** Stratum of tubes rounded towards the stem, &c.

**30. B. satanas** Lenz.—**Pileus** *fuscous-tan* then *whitish*, pulvinate, *smooth*, somewhat viscous; flesh whitish, becoming red or violet when broken. **Stem** obese, *ovato-ventricose*, with blood-red reticulations above. **Tubes** free, yellow; pores minute, at first blood-red-scarlet, orange when old.

Boletus.

Handsome, robust, mild, not bitterish.

In woods. Rare. Aug.-Sept.

Pileus sometimes 20 cent. (8 in.) broad. Stem 5-7.5 cent. (2-3 in.) long. Spores earthy-yellow. *Fr.*; rich brown, oval or spindle-shaped,  $12 \times 5$  mk. *W.G.S.* *Very poisonous*. Name—*Satan*. Satanic. From its poisonous and dangerous qualities. *Lenz f.* 31. *Fr. Hym. Eur. p.* 510. *Berk. Out. p.* 233. *C. Hbk. n.* 723. *S. Mycol. Scot. n.* 683. *Fl. Bat. t.* 1040. *Hussey i. t.* 7. *Quel. t.* 15. *f.* 1.—*Rocq. t.* 6. *Krombh. t.* 38. *f.* 1-6. *Vivian. t.* 40.

**31. B. luridus** Schæff.—**Pileus** often 20 cent. (8 in.) but occurring 5-7.5 cent. (2-3 in.) broad, tomentose, *umber-olivaceous*, then *somewhat viscous*, fuliginous, pulvinate; flesh yellow, becoming azure-blue when broken. **Stem** 5 cent. (2 in.) and much more long, stout, *vermilion-red*, marked with reticulations or dots. **Tubes** free, yellow, at length becoming green; pores round, vermilion then *orange*.

Stem somewhat orange at the apex. Taste pleasant.

In mixed woods, &amp;c. Common. July-Nov.

Spores oval, greenish slate-colour,  $17 \times 9$  mk. *W.G.S.* *Very poisonous*. Name—*luridus*, lurid in colour. *Schæff. t.* 107. *Fr. Hym. Eur. p.* 511. *Syst. Myc. i. p.* 391. *Sv. atl. Sv. t.* 12. *Berk. Out. p.* 233. *t.* 15. *f.* 5. *C. Hbk. n.* 724. *S. Mycol. Scot. n.* 684. *Grev. t.* 121. *Barla t.* 33. *f.* 1-5. *Krombh. t.* 38. *f.* 11-17.—*Bull. t.* 100. *Rostk. t.* 31. B. stem curt, bulbous, scarcely reticulated *Bolt. t.* 85.

\* **B. erythropus** Pers.—**Stem** thinner, cylindrical, not reticulated but *dotted with squamules*, internally somewhat red.

The pileus is often tawny-rufescent. Smaller than preceding species. The flesh, according to Persoon, is red-blood-colour, according to *Fl. Dan. t.* 1792 yellow.

In woods and shrubbery. Uncommon. July-Aug.

Name—*ἐρυθρός*, red; *πούς*, a foot. Red-stemmed. *Pers.*—*Fr. Hym. Eur. p.* 511. *Berk. Out. p.* 233. *C. Hbk. n.* 725. *S. Mycol. Scot. n.* 685. *Letell. t.* 612. *Barla t.* 33. *f.* 6, 7. *Harz. t.* 56.

**32. B. purpureus** Fr.—**Pileus** *purplish-red*, pulvinate, somewhat velvety, opaque, dry; flesh azure-blue only when young, then dark yellow. **Stem** stout, *yellow*, variegated with purple veins and dots. **Tubes** somewhat free, greenish-yellow; pores minute, *purple-orange*.

Very handsome. The stem varies reticulated only at the apex, becoming red internally, chiefly at the base.

In woods. Rare.

Spores somewhat spindle-shaped, green-sepia-brown,  $11 \times 6$  mk. *W.G.S.* Name—*purpureus*, purple. *Fr. Hym. Eur. p.* 511. *Sv. atl. Sv. t.* 41. *Berk.*

*Out. p.* 234. *C. Hbk. n.* 726. *Barla t.* 33. *f.* 8-10. *Saund. & Sm. t.* 43.—*Boletus. Krombh. t.* 37. *f.* 12-15. *Clus. Pern. gen.* xix. 3.

*Ser. II. TEPHROLEUCI.* Tubes at first white or grey.

G. Favosi. Tubes large, angular, unequal, &c.

**33. B. laricinus** Berk.—**Pileus** 5-7.5 cent. (2-3 in.) broad, *dirty white*, with livid stains, covered at first with dirty-yellow or brownish evanescent slime, *somewhat scaly*; flesh white, very slightly tinged with yellow. **Stem** 5 cent. (2 in.) or more long, 12-16 mm. ( $\frac{1}{2}$ - $\frac{3}{4}$  in.) thick, cribose above the ring, pitted below, dirty white. **Tubes** adnate, somewhat decurrent, compound, at first nearly white.

In larch woods. Frequent. Aug.-Sept.

The squamules on the pileus are the fragments of a veil. Spores oblong, brownish clay-coloured. *M. J. B.*; spindle-shaped, pale-brown, 10 × 4 mk. *W. G. S.* Name—*larix*, larch. From habitat. *Berk. Eng. Fl. v. p.* 148. *Out. p.* 230. *C. Hbk. n.* 703. *S. Mycol. Scot. n.* 686. *Fr. Hym. Eur. p.* 513. *Hussey i. t.* 25.

**34. B. viscidus** Linn.—**Pileus** dingy yellowish, pulvinate, soft, smooth, viscid; *veil somewhat annular, torn*, white, somewhat appendiculate. **Stem** viscous, white becoming yellow, thickened at the base, *reticulated above*. **Tubes** adnate; pores large, unequal, livid.

Perhaps too nearly allied to *B. laricinus*. The stem often appears torn from the agglutinated fragments of the veil.

In woods. Rare.

Spores fuscous-black. *Fr.*; spindle-shaped, elongated, brownish green, 9 × 3 mk. *W. G. S.* Name—*viscum*, bird-lime. Viscid. *Linn.—Fr. Hym. Eur. p.* 513. *Icon. t.* 178. *f.* 3. *Berk. Out. p.* 235. *C. Hbk. n.* 727.

H. Versipelles. Tubes minute, round, equal, &c.

**35. B. versipellis** Fr.—**Pileus** rufous, pulvinate, *dry, at first compact and tomentose*, then scaly and becoming even, *appendiculate with the annular, membranaceous, fugacious veil*. **Stem** solid, attenuated upwards, wrinkled-scaly. **Tubes** free, plane, minute, dingy white; pores grey.

Truly distinct from *B. scaber*, but it is difficult to define exactly the difference between them on account of analogy in colour and variation in stature. The colour of *B. versipellis* is more constantly rufous.

In woods and heath. Frequent. Aug.-Oct.

Boletus.

Spores spindle-shaped, pale, ochraceous,  $15 \times 5$  mk. *W.G.S.*; fusoid-obl.,  $17-20 \times 6-7.5$  mk. *K.* Name—*verto*, to turn; *pellis*, the skin. From its changeable appearance. *Fr. Hym. Eur. p.* 515. *Berk. Out. p.* 235. *C. Hbk. n.* 728. *S. Mycol. Scot. n.* 687.—A. pileus rufescent. *Schæff. t.* 103. *Krombh. t.* 32. *Sow. t.* 110. *Rostk. t.* 39. B. pileus fuliginous. *Batt. t.* 30. *f. A.*

**36. B. scaber** Fr.—**Pileus** variable in colour, red or orange, &c., pulvinate, *smooth, viscid when moist*, at length rugulose or rivulose, margin at first furnished with a cortina. **Stem** solid, attenuated upwards, *roughened with fibrous scales*. **Tubes** free, convex, white then dingy; pores minute, round.

Sometimes robust, sometimes slender. The chief varieties are as follows: A. pileus red or orange. *Bull. t.* 489. *f.* 2. B. pileus fuliginous or cinereous. *Fl. Dan. t.* 833. *Bull. t.* 489. *f.* 1. *Sow. t.* 175. *Krombh. t.* 35. *f.* 1-6. C. pileus fuscous-black, stem floccoso-scaly, tubes shining white. *Schæff. t.* 104. D. pileus fuscous-olivaceous. *Bolt. t.* 86. E. pileus wholly white. *Rostk. t.* 48.

In woods, &c. Common. July–Nov.

Pileus 5–17.5 cent. (2–7 in.) or more broad. Stem 10–15 cent. (4–6 in.) or more long. Scales on the stem black. Spores spindle-shaped, pale brown,  $14 \times 5$  mk. *W.G.S.*; fusoid-obl., light yellowish,  $14-18 \times 6$  mk. *K.* Edible. Name—*scaber*, rough. From the stem. *Fr. Hym. Eur. p.* 515. *Sv. atl. Sv. t.* 14. *Berk. Out. p.* 235. *C. Hbk. n.* 729. *S. Mycol. Scot. n.* 688. *Vittad. t.* 28. *Vent. t.* 9, 10. *Barla t.* 35. *f.* 6-12. *Rostk. t.* 40. *Harz. t.* 2. *Gonn. & Rab. vii. t.* 3. *Badh. i. t.* 7. *f.* 1, ii. *t.* 6. *f.* 1, 2. *Hussey i. t.* 57. *Hogg & Johnst. t.* 22.

I. Hyporhodii. Tubes adnate to the stem, whitish, &c.

**37. B. felleus** Bull.—**Pileus** 7.5–10 cent. (3–4 in.) broad, somewhat pale yellowish, pulvinate then expanded, soft, *smooth*, even; flesh when broken white then flesh-colour. **Stem** 7.5 cent. (3 in.) and more long, solid, attenuated and reticulated upwards. **Tubes** adnate, convex, elongated; pores angular, white then flesh-colour.

Taste bitter. The colour of the pileus varies chestnut, pale yellowish and tawny. Stem of the same colour, here and there becoming olive, more rarely dotted with squamules as in *B. erythropus*.

In woods. Uncommon.

Spores spindle-shaped, salmon-colour,  $16 \times 6$  mk. *W.G.S.*; fusoid-obl., hyaline,  $12-16 \times 4.5-5.4$  mk. *K.* Poisonous. Name—*fel*, gall. Bitter. *Bull. t.* 379. *Fr. Hym. Eur. p.* 516. *Syst. Myc. i. p.* 394. *Sv. atl. Sv. t.* 52. *Berk. Out. p.* 236. *C. Hbk. n.* 731. *Rostk. t.* 43. *Krombh. t.* 74. *f.* 1-7.

**38. B. alutarius** Fr.—**Pileus** *fuscous-tan*, pulvinate then expanded, soft, *velvety*, becoming smooth. **Stem** solid, bulbous, rather even, rugged at the apex. **Tubes** depressed round the



stem, short, plane; pores round, white, becoming fuscous when bruised.

Very much allied to *B. felleus*, but easily distinguished by its somewhat unchangeable flesh, and mild watery taste.

In woodland pastures. Rare.

Name—*aluta*, tanned leather. From the colour. *Fr. Hym. Eur. p.* 516. *Berk. Out. p.* 235. *C. Hbk. n.* 730.

K. Cariosi. Stem externally never reticulated, &c.

**39. B. cyanescens** Bull.—**Pileus** 5–12.5 cent. (2–5 in.) broad, somewhat fuscous or tan, convexo-expanded, adpressedly tomentose, floccoso-scaly, opaque; *flesh rigid, white, soon becoming deep blue when broken.* **Stem** stuffed with a spongy pith, at length cavernous, ventricose, villosopruinose, of the same colour as the pileus, constricted at the apex where it is even, white. **Tubes** free, white, at length light yellow; pores minute, round, when young scarcely 2 mm. (1 lin.) long.

Very remarkable; when compressed pouring out azure-blue juice.

In mixed woods. Rare. Aug.–Sept.

Spores at first milk-white, at length pale lemon-yellow. *Fr.*; spindle-shaped, 14 × 8 mk. *W.G.S.* Becoming *instantly* intense blue when broken. The down on the pileus is very matted and floccosely scaly. Name—*cyaneus*, deep blue. From the flesh becoming deep blue. *Bull. t.* 369. *Fr. Hym. Eur. p.* 517. *Syst. Myc. i. p.* 395. *Sv. atl. Sv. t.* 80. *Berk. Out. p.* 236. *C. Hbk. n.* 732. *S. Mycol. Scot. n.* 689. *Letell. t.* 654. *Krombh. t.* 35. f. 7-9. *Barla t.* 37. *Rostk. t.* 44. *Harz. t.* 71. *Saund. & Sm. t.* 47 (dark).

**40. B. castaneus** Bull.—**Pileus** 5–7.5 cent. (2–3 in.) broad, *cinnamon*, opaque, convexo-expanded or depressed, firm, even, *velvety*; flesh white, unchangeable. **Stem** 5–7.5 cent. (2–3 in.) long, stuffed then hollow, attenuated from the somewhat bulbous base, even, *velvety*, opaque, *cinnamon*. **Tubes** free, curt, white; pores minute, round.

Smaller than *B. cyanescens*, not juicy. Pileus at first embracing the base of the stem, whence the stem when young is obsoletely somewhat marginatobulbous, not constricted at the apex as in *B. cyanescens*.

In woods. Rare.

Name—*castanea*, chestnut. Chestnut-coloured. *Bull. t.* 328. *Fr. Hym. Eur. p.* 517. *Syst. Myc. i. p.* 392. *Berk. Out. p.* 236. *C. Hbk. n.* 733. *Barla t.* 32. f. 11-15. *Krombh. t.* 4. f. 28-30. *Hussey, ii. t.* 17.

### Subg. I. GYRODON.

No British species.

Boletus.

## Subg. II. BOLETINUS.

No British species.

Of uncertain affinity.


**41. C. carnosus** Rostk. Compact.—**Pileus** fuscous, pulvinate, smooth; flesh pallid, dirty yellowish. **Stem** short, firm, somewhat striate, rufescent light yellow. **Tubes** adnate (according to the fig. depressed round the stem), dark yellow; pores rather large, angular, of the same colour.

According to the fig. the pileus is purplish-fuscous. In woods. Stoke Poges. Name—*carnosus*, fleshy. *Rostk. t.* 14. *Fr. Hym. Eur.* p. 520. *B. & Br. n.* 1426.

Strobilomyces.

GENUS XXII.—**Strobilomyces** Berk. (στροβίλος, a pine-cone; μύκης, a fungus).

Hymenophore quite distinct from the hymenium. **Pileus** fleshy, at length tough. Spores globose or broadly elliptic, minutely rough. *Berk. Out.* p. 236.



LXI. *Strobilomyces strobilaceus*.  
One-sixth natural size.

**1. S. strobilaceus** Berk. Blackish-umber.—**Pileus** pulvinate, rough with thick floccose scales. **Stem** equal, veiled, sulcate above. **Tubes** adnate, white, angular, whitish-brown.

**Pileus** tessellated or cracked like the cone of the Scotch fir.

In fir woods. Very rare. Bullstrode. Ludlow. Haywood Forest, Herefordshire. Oct.

**Pileus** 5–10 cent. (2–4 in.) broad, brown-umber, tough, pulvinate, broken up into large, thick, projecting scales, merging into a thick, floccose, ragged and pendulous, white veil at margin. **Stem** 7.5–15 cent. (3–6 in.) long, 12 mm. ( $\frac{1}{2}$  in.) or more thick, solid, equal, coarsely fibrillose, brown at the base, white at the deeply sulcato-reticulated apex, which runs gradually into the tubes. **Tubes** white, very large, adnate, or with a decurrent tooth, anastomosing. The whole plant turns deep sienna-red when cut or bruised. *W.G.S.* Spores very dark. *M. f. B.*; oval, nearly globose, stalked, blackish-brown,  $9 \times 13$  mk. *W.G.S.* *Berk. Out.* p. 236. *C. Hbk. n.* 734. *Boletus strobilaceus* *Scop. Ann. Nat. Hist.* iv. t. 1. f. 1. *Berk. Eng. Fl.* v. p. 154.

*Fr. Hym. Eur. p. 513. Dicks. Crypt. Brit. t. 3. f. 2. Krombh. t. 74. f. 12, 13. Rostk. t. 38. Quel. t. 16. f. 1.—Vent. t. 43. f. 1, 2. Pers. Myc. Eur. 2. t. 19.*


GENUS XXIII.—**Fistulina** (*fistula*, a pipe. From the pipe-like character of the tubes). Bull. Champ. p. 314. Fistulina.

Hymenium inferior, at the first dotted over with warts, then protruding cylindrical tubes, which are free and distinct from one another. *Somewhat fleshy fungi, growing on wood, intermediate between Polyporei and Hydnei.* *Fr. Hym. Eur. p. 522.*

1. **F. hepatica** Fr. Juicy-fleshy, not rooting.—**Pileus** entire, blood-red; flesh thick, soft, viscid above, traversed with tenacious fibres, hence variegated-red. **Tubes** at first pallid.

Changeable in form, sessile or extended into a lateral stem.

On living oak. Common in England. Aug.—Oct.



LXII. *Fistulina hepatica*. One-fifth natural size. Section two times natural size.

Pileus roundish, dimidiate or subspathulate. Substance thick and juicy, marbled like beet-root, distilling a red pellucid juice; hymenium convex. Taste rather acid. It attains sometimes an enormous size, even nearly 30 lb. in weight. *M. J. B.* Spores salmon-colour, nearly round with an oblique apiculus, 3 mk. *W. G. S.* Edible and nourishing, but rather coarse, and not of a very pleasant flavour. Known as the beef-steak fungus. Slices exactly like beef-steak. Name—*ἥπατικός*, of the liver; diseased in the liver. From its appearance. *Fr. Hym. Eur. p. 522. Sv. atl. Sv. t. 25. Berk. Out. p. 257. t. 17. f. 1. C. Hbk. n. 841. S. Mycol. Scot. n. 690. Grev. t. 270. Ventur. t. 36. f. 1, 2. Barla t. 30. f. 4-7. Hogg & Johnst. t. 7. Hussey i. t. 65. Badh. i. t. 12. f. 4; ii. t. 12. f. 2. Price f. 22.—Huds. Schæff. t. 116-120. Bolt. t. 79. Sow. t. 58. Fl. Dan. t. 1039. Bull. t. 74, 464, 497.*

GENUS XXIV.—**Polyporus**. Fr. Obs. Myc. i. p. 121.  
(*πολύς*, many; *πόρος*, a pore.)

Polyporus.

Hymenophore descending and forming a trama between the pores. Hence the pores are connate with the substance of the


Polyporus. pileus, and not separable from one another; they are at first rudimental or altogether wanting, then round, angular, or torn.

*Persistent fungi, most of them growing on wood, not fully formed before expansion like Boleti, but growing gradually and indefinitely. Odour, if any, slightly acid.*

A genus holding the like central place in this order as *Agaricus* does in the former one, of immense extent, and specially abundant in the number of its species in warm countries on account of the greater variety of trees.

*Fr. Hym. Eur. p. 522.*

A few are edible, but not recommended.



LXIII. *Polyporus perennis*. One-third natural size.

I. MESOPUS (μεσός, middle; πούς, a foot). Pileus entire or excentric. Stem distinct, vertical, simple, of the same colour at the base, and not definitely blackish.

A. Carnosi (*caro*, flesh). Pileus fleshy, fragile or pliant (but not coriaceous), zoneless. White-spored, growing on the ground, edible, autumnal.

\* Pileus unpolished, scaly or floccose.

\*\* Pileus polished, smooth.

B. Lenti (*lentus*, tough, pliant). Pileus fleshy-pliant then coriaceous, zoneless. White-spored, growing on trunks, persistent.

\* Pileus unpolished, scaly or villous.

\*\* Pileus even, smooth.

C. Spongiosi (*spongia*, a sponge). Pileus at first spongy-soft, absorbing water, tomentose, then corky or coriaceous. Stem curt, irregularly shaped. Pores irregularly shaped, pruinose, changing colour, but the spores (of all of them?) white.—A very varied group, of striking appearance, mostly growing on the ground. At first they appear as a rooting, somewhat crustaceous mass, effused and encrusting, shapeless, spongy and juicy, tomentose; from this is evolved a pileus, which is sometimes remarkably regular, with a central stem, and of large size, sometimes by growing into one another irregular and lobed, nay dimidiate. They only last in a dead state to the following spring, but thereafter are covered over with a new stratum of pores. In all these respects they are analogous to *Thelephoræ* (e.g. *Th. terrestris*). Most of them rare, confined to certain localities.

D. Subcoriacei (*corium*, a hide, leather). Pileus at the first indurated, arid, when full grown corky or coriaceous. Stem definite. Pores at first sprinkled over with a white bloom. Substance of a ferruginous colour. Growing on the ground or trunks, more rigid than the *Spongiosi*, more regular and persistent, but not reviving.

\* Pileus tomentose, velvety.

\*\* Pileus very smooth.

II. PLEUROPUS (πλευρόν, the ribs or side; πούς, a foot). Pileus pliant or corky, horizontal, not circular. Stem simple, ascending, corticate, either defi-

*nitely lateral or excentric, always black at the base. Growing on wood, and Polyporus. not on the ground.*

A well-defined group. Species with excentric stems occur both in this and in the former group, but the *definitely black base* at once distinguishes them.

A. Lenti. Pileus *fleshy-pliant*, zoneless. Stem *excentric, blackish at the base*. Tubes short. Substance pallid, somewhat fibrous.

\* *Pileus scaly or floccose.*

\*\* *Pileus even, very smooth.*

B. Suberoso-lignosi (corky-woody). Pileus thick, hard, zoneless. Stem stout, vertical, of the same colour at the base. Pores elongated. *Species not closely allied to one another nor to the preceding group.*

III. MERISMA (μερίζω, to divide). Syst. Myc. i. p. 354. From a common trunk or tubercle proceed very many pileoli, arising from the subdivision of the primary pileus, which are at first, at least potentially, porous all over, but the side which is turned to the light is sterile. (In dark underground places they often remain club-shaped and porous.) *Very handsome fungi, becoming free, by far the largest of all fungi, composed of many small pilei cohering the one with the other.*

A very distinct group, well marked off from the rest, and only verging back by imperfect forms to the *Mesopodes*.

A. Carnosi. Pileus *fleshy*, firm, floccoso-fibrous, zoneless, not coriaceous-indurated. Tufts central, stipitate, springing from a common base, more or less concrete, so that in slender forms they occasionally resemble a central, simple, lobed pileus. Pores separating from the pileus. *White-spored, partially growing on the ground, autumnal, all certainly edible.*

B. Lenti. Pileus *fleshy-pliant* then somewhat coriaceous, more or less zoned, fibrous within. Pores adnate. *Tufts lateral, somewhat stipitate, in many imbricated layers, stems more or less connate, or growing from a common tuber. White-spored, not edible, fragile when old, autumnal and not lasting till the following spring, occurring at the base of trunks.*

C. Caseosi (caseus, cheese). Pileus *cheesy*, at first soft and juicy, then arid, fragile, without a pellicle, zoneless; pores separating. *Tufts sessile on the stems of trees, commonly dimidiate, but in a horizontal situation expanded on all sides, central, at the first evolved from a single shapeless tubercle into numerous pileoli. Acid. Occurring from spring through the whole summer, but soon decaying.*

D. Suberosi (suber, cork). Pileus *corky* or coriaceous, persistent, tough; substance floccose, somewhat soft, suitable for tinder, pores adnate. *Tufts somewhat sessile, lateral or central according to situation, but the pileoli are drawn together and free at the base and not effused. On trunks close to the ground. Not edible.*

IV. APUS (α, without; πούς, a foot. Stemless). Pileus *sessile*, normally adnate by a thickened and dilated base and dimidiate, or entire and attached behind (commonly by an umbo), more rarely attenuated and sessile (a transition to the *Pleuropodes*), more frequently wholly resupinate (always abnormal states; the true *Resupinati* are very different having no pileus). *Growing on wood, very abundant, with an endless variety of forms.*

A. Anodermei (ἀν, δέρμα, skin. Without a cuticle). Pileus *without a cuticle*, consequently the naked surface is broken up into flocci or fibres, zoneless, but within transversely zoned, or more or less fibrous. *Always annual, somewhat fleshy, and not reviving.*

I. Carnosi. Pileus *cheesy*, at first of a watery softness, fragile, flocculose

Polyporus. and not bristly-hispid, pores white. *When more fully grown sometimes soft, sometimes hardened, but soon putrefying and not lasting through the winter.*

\* Eupolypori. Pores round, entire, obtuse, without small teeth.

\*\* Pores elongated, flexuous, acute, and torn. Species in this section do not become hard.

2. Lenti. *Pileus fleshy-pliant* (tough even in younger specimens), soft, elastic, villosa-tomentose on account of the fibroso-floccose texture of the substance; pores somewhat adnate, coloured. *Always soft, somewhat flexile, lasting to the following spring*, joined by intermediate forms to the preceding group, but manifestly presenting a different type.

\* Substance coloured.

\*\* Substance white.

3. Spongiosi. At first and while fresh moist, spongy, when dry firm, elastic, for the most part bristly-hispid, internally fibrous. The flesh commonly presents an intermediate stratum, more compact than the exterior. *Firm, but annual, autumnal, the finest specimens on living trunks.* Commonly mixed up with *P. stipposi* which are always dry and rigid not spongy.

\* Substance coloured.

\*\* Substance and spores white.

B. Placodermei (πλάξ, anything flat; δέρμα, skin). Pileus clothed with a continuous crust, zoneless or concentrically sulcate. *Persistent.*

4. Suberosi. Pileus at the first somewhat fleshy and juicy, then hardened, covered with a rather thin crust; pores slender, at length somewhat separating. *Autumnal, in some cases lasting to the following spring, but never reviving so as to produce strata. Differing from the foregoing by the cuticle, from all the following by the pileus being at first fleshy and juicy.*

\* Substance coloured.

\*\* Substance, which is not zoned, and spores white.

5. Fomentarii (*fomentum*, fuel. From some species, e.g., *P. fomentarius*, being used as tinder). Pileus floccose, of the nature of tinder, juiceless from the first, not fleshy or spongy, covered with a hard, horny crust, the pores at length stratosed. *Growing on wood, somewhat pulvinate, truly perennial, adding new strata year by year.* In specimens from 10 to 20 years of age narrow zones corresponding with warm and dry years, and thick and broad ones corresponding with rainy years can be observed. Care must be taken not to be deceived by a specimen of the first season, without strata. In old specimens the cuticle is here and there broken into chinks or wasted away.

\* Substance and pores umber or fuscous.

\*\* Substance and pores somewhat ferruginous.

\*\*\* Substance white or pallid.

6. Lignosi (*lignum*, wood). Pileus from the first hard, woody, juiceless, covered with a thin, somewhat varnished crust, which is smooth or becomes so, somewhat effused at the base; pores not stratosed. *Perennial* (not putrefying) but not stratosed, flatter than the former group or effuso-reflexed.

C. Inodermei (ἴσ, a fibre; δέρμα, skin). Pileus from the first arid and firm, with a thin fibrous cuticle. *Annual or biennial fungi, not reviving.*

7. Stipposi (*stippa*, flax). Pileus flocculose then smooth, or adpressedly villous, uneven, zoneless, texture fibrous.

\* Substance coloured.

\*\* Substance white.

8. Coriacei (*corium*, a hide). Pileus coriaceous, villous, banded with concentric zones which are generally of different colours.

V. RESUPINATI. Pileus none, consequently the fungus is absolutely resupinate; pores immediately seated on wood or on the mycelium without the

intervention of any subiculum (except in the case of *P. medullapanis* and *P. Polyporus mucidus*).

The species of this group must be cautiously examined. For many species of *Apodes* are found in a resupinate state when growing in a horizontal situation, e.g., *P. mollis*, *destructor*, *croceus*, *nidulans*, *amorphus*, *adustus*, *igniarius*, *radiatus*, *conchatus*, *abietinus*.

In a horizontal situation the pores in this group are straight, in a vertical one oblique, nay gaping at one side.

\* Pores fuscous or blackish.

\*\* Pores ferruginous or cinnamon.

\*\*\* Pores purple or violaceous.

\*\*\*\* Pores red.

\*\*\*\*\* Pores yellowish. Compare *P. vulgaris*.

\*\*\*\*\* Pores white then changing colour.

\*\*\*\*\* Pores persistently white, minute, crowded, in a plane situation round, equal.

\*\*\*\*\* Pores persistently white, unequal, angular, close to one another, commonly rather large.

\*\*\*\*\* Pores white, superficial, distant, punctiform.

## I. MESOPUS. Pileus entire, &c.

### A. Carnosi.

\* *Pileus unpolished.*

1. **P. leucomelas** Fr.—Pileus 5-7.5 cent. (2-3 in.) broad, black-fuliginous, fleshy, somewhat fragile, irregularly shaped, slightly silky-squamulose; flesh white, reddish when broken. **Stem** 2.5-7.5 cent. (1-3 in.) long, stout, unequal, somewhat tomentose, of the same colour as the pileus. **Pores** rather large, unequal, cinereous, whitish.

The stem is sometimes curt, 2.5 cent. (1 in.) and tuberous, sometimes 5-7.5 cent. (2-3 in.) and equal, also becoming black internally. The pores become black in dried specimens.

In fir wood. Rothiemurchus. Sept.

Name—λευκός, white; μέλας, black. From the colours. *Fr. Hym. Eur. p. 524. Icon. t. 179. f. 1. B. & Br. n. 1682. S. Mycol. Scot. n. 691.—Pers. Syn. p. 515.*

\*\* *Pileus polished, smooth.*

No British species.

### B. Lenti. Pileus fleshy-pliant then coriaceous, &c.

\* *Pileus unpolished, scaly or villous.*

2. **P. lentus** Berk.—Pileus 4 cent. (1½ in.) broad, reddish-brown at length ochraceous, fleshy, but tough, umbilicate, minutely scaly, especially at first. **Stem** 12 mm.-2.5 cent. (½-1 in.) long, 9 mm. (⅓ in.) thick, rather slender, incurved,

*Polyporus*. central or excentric, hispid, furfuraceous, nearly of the colour of the pileus. **Pores** irregular, large, decurrent, white.

The stem is covered with pores to the very base, only the lower ones are abortive. Allied to *P. squamosus*.

On *Ulex*, &c. Frequent. April.

Name—*lentus*, tough, pliant. *Berk. Out. p. 237. t. 16. f. 1. Eng. Fl. v. p. 134. C. Hbk. n. 738. S. Mycol. Scot. n. 692. Fr. Hym. Eur. p. 526.*

**3. P. brumalis** Fr.—**Pileus** 2.5–10 cent. (1–4 in.) broad, fleshy-pliant then coriaceous, somewhat umbilicate, zoneless, in the first year *villous, fuliginous*, in the second squamulose becoming smooth, becoming pale; flesh white. **Stem** 2.5–5 cent. (1–2 in.) long, 4–8 mm. (2–4 lin.) thick, *hirsute*, squamulose. **Pores** *oblong* and angular, thin, acute, toothed, white, at length becoming yellow.

The margin of the pileus is somewhat fimbriato-ciliate. In hollow trunks, &c., there is a very small form, somewhat pulverulent.

On dead branches. Frequent. Oct.–March.

Pores oblong, slightly curved, hyaline, 6×2 mk. *K*. Name—*bruma*, winter. *Fr. Hym. Eur. p. 526. Syst. Myc. i. p. 348. Berk. Out. p. 237. C. Hbk. n. 735. S. Mycol. Scot. n. 693. Pers. Batsch f. 42 (first year). Fl. Dan. t. 1297 (second year). P. floccipes Rostk. 28. t. 13.*

**\*\* Pileus even, smooth.**

**4. P. fuscidulus** Fr.—**Pileus** 2.5–5 cent. (1–2 in.) broad, *fuscous-yellowish*, fleshy, pliant, somewhat coriaceous, convexo-plane, zoneless, smooth; flesh white-yellowish. **Stem** 2.5–5 cent. (1–2 in.) long, 4 mm. (2 lin.) thick, equal or thickened at both ends, *smooth*, fuscous becoming yellow. **Pores** somewhat rounded-angular, obtuse, quite entire, yellowish.

Among chips, &c. Rare. Darlington.

Name—*fuscus*, fuscous, brown. Slightly fuscous. *Fr. Hym. Eur. p. 528. Berk. Out. p. 237. C. Hbk. n. 736. Schrad. Bot. t. 170.*

**5. P. leptcephalus** Fr. Small.—**Pileus** under 2.5 cent. (1 in.) broad, *pale then fawn-colour*, pliant, then coriaceous, convexo-plane, thin, smooth, even, *zoneless*; flesh white. **Stem** short, scarcely 2.5 cent. (1 in.) smooth, pallid. **Pores** adnate, minute, somewhat round, obtuse, *whitish*.

The margin of the pileus is somewhat repand.

On trunks. Rare.

Stem thick as a crow-quill. *With.* Name—*λεπτός*, thin; *κεφαλή*, head.



With thin pileus. *Fr. Hym. Eur. p. 528. Syst. Myc. i. p. 349. Berk. Out. Polyporus. p. 237. C. Hbk. n. 737. S. Mycol. Scot. n. 694.—Jacq. Misc. i. t. 12. Mich. t. 70. f. 7. Paul. t. 164. f. 12.*

C. Spongiosi. Pileus at first *spongy-soft*, &c.

**6. P. Schweinizii** Fr.—Pileus 20 cent. (8 in.) and more broad, *date-brown*, thick, spongy-corky, *strigoso-tomentose*, *rugged*. **Stem** thick, very short or obsolete, ferruginous. **Pores** large, variable in size and torn, *sulphur-greenish*.

Very large, sometimes regular and plano-cup-shaped, sometimes irregularly shaped and dimidiate from growing into each other and incrusting, at first tawny-tomentose, soon only at the margin, then when old wholly fuscous. The substance when the plant is in fullest vigour is very soft-spongy, then hardened, fragile when dry, rhubarb-colour becoming fuscous.

In pine woods about roots. Rare. July–Oct.

It occurs with a central stem and also imbricated. *B. & Br.* Spores oval, white or transparent,  $3 \times 8$  mk. *W.G.S.* Name—after Schweinitz. *Fr. Hym. Eur. p. 529. Icon. t. 179. f. 3. Berk. Out. p. 238. C. Hbk. n. 739. S. Mycol. Scot. n. 695. Sv. Bot. t. 720.*

**7. P. rufescens** Fr. *Flesh-colour*.—Pileus 5–10 cent. (2–4 in.) broad, spongy, soft, unequal, *hairy*. **Stem** short, irregularly shaped. **Pores** large, *sinuated* and torn, white-flesh-colour.

Its growth is quite that of *P. Schweinizii*, but it is smaller, softer, and different in colour both internally and externally.

On stumps. Frequent. Nov.

Name—*rufesco*, to become red. *Fr. Hym. Eur. p. 529. Syst. Myc. i. p. 351. Berk. Out. p. 238. C. Hbk. n. 740. S. Mycol. Scot. n. 696.—Sow. t. 191.*

D. Subcoriacei. Pileus at the first indurated, &c.

\* *Pileus tomentose, velvety.*

**8. P. perennis** Fr.—Pileus *cinnamon* then *date-brown*, coriaceous, tough, plano-infundibuliform, velvety becoming smooth, *zoned*. **Stem** slightly firm, thickened downwards, velvety. **Pores** minute, angular, *acute*, at first sprinkled with a white bloom, then naked, *torn*.

The margin when the plant is in fullest vigour is fimbriate, then entire.

On the ground and stumps. Frequent. July–Jan.

Pileus 4–5 cent. ( $1\frac{1}{2}$ –2 in.) broad. Stem 2.5 cent. (1 in.) long, varying in thickness. *M. f. B.* Spores oval, white,  $3 \times 2$  mk. *W.G.S.* Name—*perennis*, perennial. *Fr. Hym. Eur. p. 531. Berk. Out. p. 238. C. Hbk. n. 741. S. Mycol. Scot. n. 697. Rostk. t. 6. Hussey i. t. 51.—Linn. Schæff. t. 125. Sow. t. 192. Fl. Dan. t. 1075. Bull. t. 28, 449. f. 2. Bolt. t. 87.*

Polyporus.

**\*\* *Pileus very smooth.***

No British species.

**II. PLEUROPUS.** Pileus pliant or corky &c.**A. Lenti.** Pileus *fleshy-pliant*, &c.**\* *Pileus scaly or floccose.***

**9. *P. squamosus* Fr.**—Pileus 3 in.—1½ ft. broad, somewhat ochraceous, *variegated with broad, adpressed, spot-like, centrifugal, darker scales*, fleshy pliant, fan-shaped, flattened. **Stem** excentric and lateral, obese, *reticulated* at the apex, blackish at the base. **Pores** thin, variable (at first minute), then large angular and torn, pallid.



LXIV. *Polyporus squamosus*.  
One-seventh natural size.

Handsome, commonly very large, somewhat central and umbilicate when young, at length lateral, very variable in shape.

On trunks and stumps, chiefly ash. Common. May–Nov.

When young a mere swollen blackish lump. Spores oval, white, 14 × 6 mk. *W.G.S.* Name—*squama*, a scale. Scaly. *Fr. Hym. Eur. p. 532. Syst. Myc. i. p. 343. Berk. Out. p. 238. C. Hbk. n. 742. S. Mycol. Scot. n. 698. Grev. t. 207. Rostk. t. 2. Ventur. t. 37. Hussey i. t. 33. —Huds. Bolt. t. 77. Schæff. t. 101, 102.*

*Bull. t. 19. Harz. t. 32. Fl. Dan. t. 983, 1196. Paul. t. 16. Sterb. t. 13, 14. B. pileus fuliginous.*

**10. *P. Michelii* Fr.**—Pileus 5–7.5 cent. (2–3 in.) broad, yellowish-white, fleshy-pliant, depressed, repand, *slightly silky, somewhat squamulose*. **Stem** 4 cent. (1½ in.) long, somewhat lateral, bulbous, rough, white, becoming fuscous at the base. **Pores** large, somewhat round-oblong, entire, white.

Allied to *P. squamosus*.

On stumps. Penzance.

Name—after Micheli. *Fr. Hym. Eur. p. 533. Syst. Myc. i. p. 343. B. & Br. n. 1963. Rostk. t. 1. —Mich. t. 61. f. 2.*

**11. *P. melanopus* Fr.**—Pileus 5–7.5 cent. (2–3 in.) broad, white then yellowish-fuscous, fleshy-pliant, plane then infundibuliform, *at first delicately flocculose*; flesh thick, white, soft, not

becoming woody. **Stem** 4 cent. (1½ in.) long, 6–8 mm. (3–4 lin.) *Polyporus*. thick, excentric, *somewhat velvety*, incurved, *thickened downwards*, black. **Pores** decurrent, curt, minute, obtuse, unequal, white.

The pileus is at length, and at first sight smooth. Stem diffused into the pileus.

On dead wood and roots. Uncommon. Aug.–Sept.

Name—μέλας, black; πούς, a foot. Black-stemmed. *Fr. Hym. Eur. p. 534. Syst. Myc. i. p. 543. B. & Br. n. 1286. S. Mycol. Scot. n. 699. Rostk. t. 4.—Swartz V. A. H.*

**\*\* Pileus even, very smooth.**

**12. P. Rostkovii** Fr.—**Pileus** *smoke-colour*, fleshy-pliant, imidiato-infundibuliform, *smooth, even*. **Stem** long, excentric, *reticulated*, abruptly black, thickened at the base. **Pores** decurrent, *large, pentagonal*, acute, toothed, white then dingy-yellowish.

In size it vies with *P. squamosus* to which it is allied. The stem is generally longer, incurved, connate and somewhat caespitose at the base. The pileus inclines to yellowish. The pores are deeply decurrent on the outer side of the stem.

On stumps. Rare.

Pileus 15 cent. (6 in.) broad. Stem long. Pores 4 mm. (2 lin.) or more long. *B. & Br.* Spores with two nuclei, 14×6 mk. *W.G.S.* Name—after Rostkovius. *Fr. Hym. Eur. p. 534. Berk. Out. p. 238. C. Hbk. n. 743. Rostk. t. 17.*

**13. P. picipes** Fr.—Pallid then chestnut, commonly pale yellowish-livid, with the disc chestnut. **Pileus** fleshy-coriaceous then rigid, tough, even, smooth, depressed at the disc or behind; flesh white. **Stem** excentric and lateral, equal, firm, *at first velvety, then naked, dotted, black up to the pores*. **Pores** decurrent, round, very small, rather slender, white, then slightly pale yellowish.

Imbricated, odour somewhat sweet. The pileus is depressed behind, commonly emarginate, infundibuliform with lobes all round.

On trunks, especially willow. Frequent. July–Dec.

Name—πίξ, pitch; πες, a foot. Black-stemmed. *Fr. Hym. Eur. p. 534. Berk. Out. p. 239. C. Hbk. n. 744. S. Mycol. Scot. n. 700.—Grev. t. 202. Pers. Ic. pict. t. 4. f. 1, 2.*

**14. P. varius** Fr.—**Pileus** commonly bay-brown, fleshy-pliant, *extenuated*, soon woody, smooth, somewhat virgate, irregularly shaped, depressed at the disc or behind. Stem excentric and

Polyporus. lateral, *even, smooth, gradually cinereous-blackish downwards*. **Pores** decurrent, minute, curt, round, unequal, whitish (internally watery) then cinnamon.

In stature, form, &c., it is not constant. The colour is commonly bay-brown (fuliginous and very evidently virgate when old) but (compare *Bolt. t. 168*) pale when young, and when full grown, as it is also internally, light yellowish. It varies sessile; also crowded in very large, irregularly shaped tufts, in habit quite as in *Schæff. t. 109, 110*.

On trunks and stumps, especially ash. Frequent. Aug.—Oct.

Pileus 12 mm.—10 cent. ( $\frac{1}{2}$ –4 in.) broad. *M. J. B.* Name—*varius*, variable (in size, form, colour, &c.) *Fr. Hym. Eur. p. 535. Berk. Out. p. 239. C. Hbk. n. 745. S. Mycol. Scot. n. 701. Batsch f. 129. Ventur. t. 57. f. 2.—Rostk. 28. t. 24, t. 20. Buxb. C. v. t. 15. f. 2.*

**15. P. elegans** Fr.—Pileus 5–10 cent. (2–4 in.) broad, unicolorous, pallid ochraceous or orange, shining, *equally fleshy*, then soon hardened, becoming woody, flattened, even, smooth; flesh white. **Stem** excentric or lateral, *even, smooth, pallid, at the first abruptly black and rooting at the base*. **Pores** plane, minute, somewhat round, yellowish-white, pallid.

White when young. The pileus is not streaked, nor is it infundibuliform, scarcely depressed. The flesh is thick to the margin.

On trunks, chiefly birch. Frequent. July–Nov.

Pileus 5–10 cent. (2–4 in.) broad. Name—*elegans*, elegant. *Fr. Hym. Eur. p. 535. Berk. Out. p. 239. C. Hbk. n. 746. S. Mycol. Scot. n. 702.—Bull. t. 46. Fl. Dan. t. 1075. f. 1. Rostk. t. 11. Bolt. t. 83.*

\* **P. nummularius** Fr. Smaller, thinner.—Pileus scarcely 2.5 cent. (1 in.) broad, *somewhat regular*. **Stem** equal, excentric.

It merges in *P. elegans* without definite line of demarcation. On trunks. Uncommon. Dec. Name—*nummulus*, a small coin. From fancied resemblance of shape. *Fr. Hym. Eur. p. 536. Berk. Out. p. 239. C. Hbk. n. 746. S. Mycol. Scot. n. 702. Rostk. t. 12. Bull. t. 124.*

**16. P. petaloides** Fr.—Pileus chestnut-fuscous, somewhat membranaceous, spathulate, *rugose, smooth, zoneless*, flaccid when moist. **Stem** lateral, ascending, compressed, smooth, not rooting, whitish, *adnate at the base which is dilated in the form of a shield*. **Pores** decurrent, very short, very small, shining white.

When dry it is involute and fragile, not becoming woody like preceding species.

On old stumps. Sibbertoft.

Name—from some fancied resemblance to the petal of a flower. *Fr. Hym. Eur. p. 536. B. & Br. n. 2023.*

B. Suberosi-lignosi. Pileus thick, hard, zoneless, &c.

Polyporus.

**17. P. lucidus** Fr.—Pileus light yellow then blood-red-chestnut, corky then woody, fan-shaped, sulcato-rugose. Stem lateral, equal, *varnished, shining*, of the same colour as the pileus. Pores determinate, long, minute, white then cinnamon.

On and about stumps. Frequent. Summer.

Pileus 5-15 cent. (2-6 in.) broad. Stem 15-25 cent. (6-10 in.) long, 2.5 cent. (1 in.) or more thick. *M. f. B.* Name—*lucidus*, bright, shining. From the stem. *Fr. Hym. Eur. p. 537. Berk. Out. p. 240. t. 16. f. 2. C. Hbk. n. 748. S. Mycol. Scot. n. 703. Krombh. t. 4. f. 22-24. Rostk. t. 13. Grev. t. 245. Ventur. t. 49.—Leys. Sow. t. 134. Fl. Dan. t. 1253. Schæff. t. 263. Batsch f. 225. Bull. t. 7, 459.*

### III. MERISMA. Very many pileoli, &c.

A. Carnosi. Pileus *fleshy*, firm, &c.

**18. P. umbellatus** Fr.—Very much branched, fibrous-fleshy, toughish. Pileoli very numerous, 1-4 cent. ( $\frac{1}{2}$ -1  $\frac{1}{2}$  in.) broad, fuliginous, rufous or pallid light yellow, *entire, umbilicate*. Stems elongated, separate, united at the base, white. Pores minute, white.

The pileoli have occurred white.

On stump. Epping. July.

Cluster about 17.5 x 12.5 cent. (7 x 5 in.) *M. C. C.* Name—*umbella*, a sunshade, parasol. From its spreading habit. *Fr. Hym. Eur. p. 537. Syst. Myc. i. p. 354. Grevillea, vol. xii. p. 36. Krombh. t. 52. f. 3-9. Lenz f. 44. Tratt. t. T. Quel. t. 18. f. 1.—Schæff. t. iii. 265, 266. Jacqu. Austr. t. 172. Fl. Dan. t. 1197.*

**19. P. frondosus** Fr.—Tuft 15-30 cent. ( $\frac{1}{2}$ -1 ft.) broad, very much branched, fibrous-fleshy, toughish. Pileoli very numerous, 1-5 cent. ( $\frac{1}{2}$ -2 in.), fuliginous-grey, *dimidiate, rugose*, lobed, intricately recurved; flesh white. Stems growing into each other, white. Pores *rather tender, very small, acute*, white.


Pores commonly round, but in an oblique position gaping open and torn.

On stumps and roots. Rare. Oct.

Name—*frons*, a leafy branch. Branched. *Fr. Hym. Eur. p. 538. Syst. Myc. i. p. 355. Sv. atl. Sv. t. 44. B. & Br. n. 1364. Krombh. t. 48. f. 17-20. Rostk. t. 18.—Fl. Dan. t. 952. Paul. t. 29. Sterb. t. 28.*

**20. P. intybaceus** Fr.—Very much branched, fleshy, somewhat fragile. Pileoli *very numerous*, pale yellowish inclining to fus-cous, *dimidiate, stretched out*, sinuate, at length spatulate.

Polyporus. **Stems** connate in a very short trunk. **Pores** firm, obtuse, white inclining to fuscous.



LXV. *Polyporus intybaceus*.  
One-sixth natural size.

*B. truncigenus* flaps elongated, conchate, extended laterally, undulated, paler. *Paul. t.* 130. Odour of mice.

On stumps, especially oak. Rare. Sept.

About same size as *P. frondosus* and larger. Spores oval, white,  $6 \times 3$  mk. *W.G.S.* Edible. Name—from its likeness in habit of growth to succory (*Cichorium intybus*). *Fr. Hym. Eur. p.* 538. *Berk. Out. p.* 240. *C. Hbk. n.* 749. *S. Mycol. Scot. n.* 704. *Hussey i. t.* 6.—*Fl. Dan. t.* 1793.

**21. P. cristatus** Fr. — Branched, firmly fleshy, fragile. **Pileoli** about 7.5 cent. (3 in.) broad, rufous-greenish, entire and dimidiate, imbricated, depressed, somewhat pulverulent-vil-  
lous, then rimosely scaly. **Stems** connate, irregularly shaped, white. **Pores** minute, angular and torn, whitish.

Very changeable in form, sometimes simple with an undulato-lobed, central pileus. *B. flabelliformis* Schæff. *t.* 113.

In beech woods. Rare.

Name—*crista*, a crest. Crested. *Fr. Hym. Eur. p.* 539. *Syst. Myc. i. p.* 356. *Berk. Out. p.* 240. *C. Hbk. n.* 750. *S. Mycol. Scot. n.* 705. *Krombh. t.* 48. *f.* 15-16. *Barla t.* 29. *f.* 4-7. *Rostk. t.* 16.—*Pers. Myc. Eur. ii. p.* 50.

*B. Lenti.* Pileus fleshy-pliant then somewhat coriaceous, &c.

**22. P. giganteus** Fr.—Tuft 1-2 ft. and more broad, in many imbricated layers, fleshy-pliant then somewhat coriaceous. **Pilei** date-brown, dimidiate, very broad, flaccid, somewhat zoned, rivulose, depressed behind. **Stems** connato-branched from a common tuber. **Pores** minute, somewhat round, pallid, at length torn.

The rigid cuticle separates into granules or fibrillose squamules. Pores becoming dark when touched.

On stumps and old roots under ground. Frequent. Aug.-Sept.

It occurs 3-4 ft. broad. Name—*gigas*, a giant. Gigantic. *Fr. Hym. Eur. p.* 540. *Syst. Myc. i. p.* 356. *Berk. Out. p.* 240. *C. Hbk. n.* 751. *S. Mycol. Scot. n.* 706. *Hussey i. t.* 82.—*Pers. Schæff. t.* 267. *Bolt. t.* 76. Var. *clavæformis*: *Holmsk. Ot. 2. t.* 13.

**23. P. acanthoides** Fr.—In many imbricated layers, pliant Polyporus. then coriaceous. **Pileoli** ferruginous, infundibuliform, incisedimidiolate, somewhat zoned, longitudinally rugose. **Stems** connato-branched, *white then rufescent*. **Pores** lamelloso-sinuate, *thin*, toothed at the edge, *white then rufescent*.

Thinner than *P. giganteus*, pale when young. Although it is somewhat coriaceous, it very soon hardens, and becomes fragile when old. *P. rufescens* differs sufficiently in the spongy, hairy pileus, in not being undulato-lobed, &c.

On trunks. Penzance.

Name—*ἀκανθος, εἶδος*. From the resemblance of the pileoli to *Acanthus* leaves. *Fr. Hym. Eur. p. 540. B. & Br. n. 1694.—Bull. t. 486. Pers. Ic. Pict. t. 6.*

C. Caseosi. Pileus cheesy, &c.

**24. P. sulphureus** Fr.—In many cæspitose layers, 1–2 ft. and more, *juicy-cheesy*. **Pilei** 20 cent. (8 in.) or more broad, *reddish-yellow*, imbricated, undulated, rather smooth; flesh light yellowish, then white, splitting open and not hardened when old. **Pores** minute, plane, *sulphur-yellow*.

Soon becoming pale. Commonly sessile, but varying with a stem (*Batt. t. 34. B. Paul. t. 14*), lateral on standing trees, but expanded on all sides on fallen ones; also club-shaped, porous throughout, *Sow*. In its fullest vigour it is filled with sulphur-yellow milk.

On living trees and stumps. Frequent. Aug.–Oct.

Whole mass sometimes 2–2½ ft. broad. A lovely plant, of delicate shades. Spores oval, white, minutely papillose, 8 × 5 mk. *W.G.S.* Edible. Name—*sulphur*, brimstone. Sulphur-yellow. *Fr. Hym. Eur. p. 542. Syst. Myc. i. p. 357. Sv. ätl. Sv. t. 88. Berk. Out. p. 241. t. 16. f. 3. C. Hbk. n. 752. S. Mycol. Scot. n. 707. Grev. t. 113. Ventur. t. 53. f. 6-7. Rostk. t. 20. Hussey i. t. 46.—Bull. t. 429. Sow. t. 135. Schæff. t. 131, 132. Fl. Dan. t. 1019.*

**25. P. Herbergii** Rostk.—Cæspitose, rather corky. **Pileoli** imbricated, bright bay, sulphury about the margin. **Pores** labyrinthiform, unequal, torn and toothed, pale cinereous.

On trunks. Edinburgh Fungus Show, 1878. Oct.

Referred with doubt to *P. spongia* by Fries. Berkeley considers it clearly the plant of Rostkovius. Name—after Herbergius. *Rostk. xxix. t. 18. B. & Br. n. 1805. Grevillea, viii. p. 6. P. spongia Fr. Hym. Eur. p. 542. S. Mycol. Scot. n. 708.*

**26. P. alligatus** Fr.—In many cæspitose layers, fibrous-fleshy, rigid-fragile. **Pilei** tan-isabelline, imbricated, unequal, *zoneless, villous*. **Pores** minute, soft, white, readily becoming stopped up with flocci.

Polyporus. Often clavate when young. Commonly wrapping round stipules and grasses.

On roots, &c. Rare.

Name—*alligo*, to bind to. From its habit of growth. *Fr. Hym. Eur. p. 543. Berk. Out. p. 241. C. Hbk. n. 753. S. Mycol. Scot. n. 709.—Sow. t. 422.*

D. Suberosi. Pileus corky or coriaceous, &c.

**27. P. heteroclitus** Fr.—In many cæspitose layers, coriaceous. **Pilei** 6 cent. (2½ in.) broad, *orange, sessile, expanded on all sides from a radical tubercle*, lobed, villous, zoneless. **Pores** irregularly shaped and elongated, golden-yellow.

On the ground under oak. Rare.

The flat pilei extend horizontally from the tubercle. Name—*ἑτερος*, one of two; *κλίνω*, to lean. Irregular, excentric. *Fr. Hym. Eur. p. 544. Syst. Myc. i. p. 344. Berk. Out. p. 241. C. Hbk. n. 754.—Bolt. t. 164.*

**28. P. salignus** Fr.—Cæspitose, coriaceous-soft, elastic. **Pilei** whitish, dimidiate, imbricated, dilated-reniform, adpressedly villous, *depresso-sulcate round the margin* which is somewhat lobed and swollen. **Pores** thin, crowded, elongated, intricately flexuous, white.

The tufts are commonly small, inodorous.

On willows. Uncommon.

Name—*salix*, willow. *Fr. Hym. Eur. p. 544. Berk. Out. p. 241. C. Hbk. n. 755.—Pers. Bolt. t. 78. Batt. t. 38. f. E.*

#### IV. APUS. Pileus *sessile*, &c.

A. Anodermei. Pileus *without a cuticle*, &c.

I. Carnosi. *Pileus cheesy*, &c.

\* Eupolyporei. *Pores round, entire*, &c.

**29. P. epileucus** Fr.—**Pileus** 7.5–10 cent. (3–4 in.) broad, 2.5–5 cent. (1–2 in.) thick, *whitish, similar internally*, cheesy-soft then firm, pulvinate, *villous-rugged*. **Pores** minute, round, quite entire, white.

Simple, semi-orbicular, concave beneath, not fibrous internally, scarcely zoned; pores at first scarcely discernible.

On stumps, chiefly fir and elm. Rare. Sept.–Oct.

Name—*ἐπί*, and *λευκός*, white. Whitish. *Fr. Hym. Eur. p. 545. Smith Seem. Journ. 1868, p. 34. B. & Br. n. 1287. C. Hbk. n. 756. S. Mycol. Scot. n. 710.—Fl. Dan. t. 1794.*



**30. P. alutaceus** Fr.—**Pileus** 2.5 cent. (1 in.) broad and thick, Polyporus. tan, fleshy, at length tough, reniform-dilated, *somewhat velvety*, obsoletely zoned within, *margin acute, even*. **Pores** very small, thin, somewhat round, whitish-tan-colour.

Somewhat imbricated, sometimes convex, sometimes flattened, occasionally hairy and rugose, plane beneath.

On decayed pine-stumps. Glamis, 1876. Sept.—Oct.

Tubes about 18 mm. ( $\frac{3}{4}$  in.) long. Name—*aluta*, tanned leather. Tan-colour. *Fr. Hym. Eur. p. 545. Syst. Myc. i. p. 360. B. & Br. n. 1799. S. Mycol. Scot. n. 711.—Rostk. t. 30.*

**31. P. pallescens** Fr.—**Pileus** 5 cent. (2 in.) broad, *yellowish*, the acute margin of the same colour, fleshy then corky, *even, smooth*, zoneless. **Pores** curt, minute, round, white, then yellowish.

Somewhat caespitose, commonly small and thin, of doubtful affinity.

On old stumps. Rare.

Name—*pallesco*, to grow pale. *Fr. Hym. Eur. p. 546. Syst. Myc. i. p. 369. Berk. Out. p. 244. C. Hbk. n. 773.—Sow. t. 230.*

**32. P. chioneus** Fr.—*White*. **Pileus** 2.5 cent. (1 in.) and more broad, fleshy, *soft, becoming even, smooth*, zoneless, often extended behind, margin inflexed. **Pores** curt, very small, round, equal, quite entire.

Always soft, fragile, hyaline-white when moist, shining white when dry. Odour acid. Without a cuticle.

On stumps, &c. Uncommon. July—Sept.

Spores white, oval,  $21 \times 3$  mk. *W.G.S.* Name— $\chi\omega\omega\nu$ , snow. Snow-white. *Fr. Hym. Eur. p. 546. Berk. Out. p. 241. C. Hbk. n. 757. S. Mycol. Scot. n. 712.—Pers. Myc. Eur. 2. t. 15. f. 2.*

**33. P. cerebrinus** B. & Br.—About 2.5 cent. (1 in.) across, snow-white, pulvinate, delicately tomentose, becoming smooth, margin crenate. **Pores** rounded, entire, dissepiments thick, obtuse.

Looks like a portion of white brain.

On fir. Glamis, 1877. Aug.

Name—*cerebrum*, brain. Brain-like. *B. & Br. n. 1800. S. Mycol. Scot. n. 713.*

**\*\* Pores elongated, flexuous, &c.**

**34. P. lacteus** Fr.—*Shining white*. **Pileus** fleshy-fibrous,

Polyporus. fragile, triangular, *pubescent*, zoneless externally and internally, margin inflexed, acute. **Pores** *thin, acute, toothed, at length labyrinthiform, torn.*

Commonly small, thin, but sometimes larger, transversely elongated, very much sloped downwards and gibbous behind, at length becoming smooth and uneven.

On dead laburnum and birch. Rare. Sept.-Nov.

Name—*lac*, milk. Milk-white. *Fr. Hym. Eur. p. 546. Icon. t. 182. f. 1. S. Mycol. Scot. n. 714. Rostk. t. 23.*

**35. P. fragilis** Fr.—Whitish, *spotted with fuscous when touched.* **Pileus** fleshy, fibrous, *fragile*, plano-depressed and reniform, *rugose*, convex beneath. **Pores** thin, elongato-flexuous, intricate.

Very distinct, although no form is constant. It varies stretched out in the form of a stem and pendulous.

On dead pine. Uncommon. Sept.-Nov.

Name—*fragilis*, fragile. *Fr. Hym. Eur. p. 546. Icon. t. 182. f. 2. Berk. Out. p. 242. C. Hbk. n. 758. S. Mycol. Scot. n. 715.*

**36. P. mollis** Fr.—**Pileus** *flesh-colour*, fleshy-fibrous, soft, effuso-reflexed, somewhat triangular, *rugose*, margin acute. **Pores** unequal, elongated, flexuous, soft, white, *spotted with red when touched.*

Not watery, but soft to the touch, seriato-elongated or resupinate.

On dead pine. Slough.

Name—*mollis*, soft. *Fr. Hym. Eur. p. 547. Icon. t. 182. f. 3. B. & Br. n. 1365.—Pers. Obs. i. p. 14.*

**37. P. cæsius** Fr.—White, *now and then tinged bluish-grey.* **Pileus** 1-5 cent. ( $\frac{1}{2}$ -2 in.) broad, fleshy, soft, tough, unequal, *silky.* **Pores** very small, unequal, elongato-flexuous and torn into teeth.

Small, variable in form, stipitate and resupinate.

On dead fir. Frequent. Sept.-Nov.

Spores pale blue, oval,  $3 \times 14$  mk. *W.G.S.* Name—*cæsius*, bluish-grey. *Fr. Hym. Eur. p. 547. Syst. Myc. i. p. 36c. Berk. Out. p. 242. C. Hbk. n. 759. S. Mycol. Scot. n. 716.—Schröd. Sow. t. 226.*

**38. P. trabeus** Rostk.—*White.* **Pileus** fleshy-fibrous, then firm, effuso-reflexed, transversely elongated, zoneless, pallid. **Pores** curt, minute, somewhat round or elongated, toothed, white.

Commonly conjoined with *P. destructor*. Both vary effused and reflexed, smooth and pubescent, but *P. trabeus* is more regular, of brighter colour, hyaline within when moist, obsoletely zoned.

On dead pine. Den of Dun, 1874. Menmuir. Feb.

Polyporus.

Name—*trabs*, timber, a vessel. Found in ships. *Rostk. t.* 28. *Fr. Hym. Eur. p.* 547. *B. & Br. n.* 1428. *S. Mycol. Scot. n.* 717.

**39. P. destructor** Fr.—*Pileus* fuscous-whitish, *watery, fleshy*, effuso-reflexed, fragile, rugose, somewhat undulated, zoned internally. Pores elongated, somewhat round, toothed or torn, white.

B. undulated, broadly expanded, marginate, whitish-date-brown. It softens and destroys wood like *Merulius lachrymans*.

On fir, larch and Scotch. Common. July-Dec.

Sometimes almost resupinate or effused. *M. J. B.* Name—*destruo*, to destroy. From its destroying wood. *Fr. Hym. Eur. p.* 547. *Berk. Out. p.* 242. *C. Hbk. n.* 760. *S. Mycol. Scot. n.* 718. *Krombh. t.* 5. f. 8.—*Schrad. Rostk. t.* 27.

2. Lenti. *Pileus fleshy pliant, &c.*

\* *Substance coloured.*

**40. P. nidulans** Fr.—*Pileus* 1-2.5 cent. ( $\frac{1}{2}$ -1 in.) thick, *somewhat pale yellowish*, of the same colour internally, fleshy, *very soft*, somewhat pulvinate, villous, becoming even, zoneless. **Pores** elongated, of medium size, unequal, angular, *tawny-brick-colour*.

Fragrant when dry, remarkable for its softness. Margin spreading, somewhat obtuse. There are two forms, one pale yellowish, the other somewhat flesh-colour.

On dead wood. Uncommon. Sept.-Nov.

Name—*nidus*, a nest. From its shape. *Fr. Hym. Eur. p.* 548. *Syst. Myc. i. p.* 364. *Berk. Out. p.* 242. *C. Hbk. n.* 761. *S. Mycol. Scot. n.* 719.—*P. rutilans Saund. & Sm. t.* 45.—*Bull. t.* 482.

**41. P. rutilans** Fr.—*Pileus tawny-cinnamon* becoming pale (tawny), of the same colour internally, fleshy-pliant, *thin*, at first villous, then becoming smooth, zoneless. **Pores** curt, minute, thin, equal, acute, cinnamon, somewhat shining.

*Pileus* at length juiceless, friable, the slight margin inflexed, but not incurved or fimbriate as in *P. cuticularis*. Variable in form, but extenuated, not pulvinate and convex on both sides like *P. nidulans*.

On fallen branches. Uncommon.

When fresh has a sweet scent like that of anise. *M. J. B.* Name—*rutilo*, to be reddish. *Fr. Hym. Eur. p.* 548. *Berk. Out. p.* 242. *C. Hbk. n.* 762.—*Pers. Ic. t.* 6. f. 4.

**42. P. gilvus** Schw.—*Pileus pale yellowish*, fleshy-pliant, effuso-reflexed, soft, even, *becoming smooth*, zoneless, margin spreading, thin, acute. **Pores** *very small*, naked, quite entire, pale yellowish-ferruginous, opaque.

**Polyporus.** Commonly longitudinally effused, flaxy in close places, soft to the touch, properly velvety, but becoming so even as to appear smooth.

On trunks. Rare.

Fries has no doubt that Sowerby's *Boletus impuber* is the same plant, and therefore it must be included as British. Name—*gilvus*, pale yellowish. *Schwein.*—*Fr. Hym. Eur. p.* 548.—*Sow. t.* 195.

**\*\* Substance white.**

**43. P. fumosus** Fr.—**Pileus** 5–10 cent. (2–4 in.) broad, *fuliginous-pallid*, fleshy-corky, *firm*, zoneless, silky then becoming smooth, adnate and dilated behind, *internally fibrous*, *somewhat zoned*. **Pores** minute, curt, round, entire, whitish-smoke-colour, darker when bruised.

Cæspitose, imbricated, moderately thick, attenuated towards the margin which becomes black. Colour more or less smoky.

On old stumps. Common. Oct.–Dec.

Smell oppressive. *M. F. B.* Name—*fumosus*, smoky. *Fr. Hym. Eur. p.* 549. *Syst. Myc. i. p.* 367. *Berk. Out. p.* 243. *C. Hbk. n.* 763. *S. Mycol. Scot. n.* 720. *Kl. Bor. t.* 392. *Rostk. t.* 42. *Pers. Tratten. t.* 3. f. 5.

**44. P. adustus** Fr.—**Pileus** cinereous-pallid, fleshy-pliant, thin, villous, *effuso-reflexed* behind, the *margin* which is tense and straight *becoming black*. **Pores** minute, curt, round, obtuse, whitish-pruinose soon *cinereous-fuscous*, the marginal ones obsolete.

Like *P. fumosus*, but thinner, darker; pileus spuriously zoned, slightly wrinkled. Pores at length blackish.

On old stumps. Frequent. Sept.–Jan.

Pores always leaving a whitish margin on the under side. Name—*aduro*, to burn. Appearing as if scorched. *Fr. Hym. Eur. p.* 549. *Berk. Out. p.* 243. *C. Hbk. n.* 764. *S. Mycol. Scot. n.* 721. *Fl. Dan. t.* 1850. f. 1. *Kl. Bor. t.* 472. *Rostk. t.* 38.—*Willd. Bull. t.* 501. f. 2. *Batsch f.* 226.

**45. P. crispus** Fr.—**Pileus** cinereous-blackish, fleshy-pliant then coriaceous, tough, rugose, effuso-reflexed behind, margin thin, *crisped*, at length black. **Pores** rather large, *unequal*, at length *labyrinthiform*, silvery-cinereous.

Small, pileus when young fuliginous-black, floccoso-rugose, zoneless, white at the margin, when full grown thinner, flexile, becoming pale, grey, becoming smooth with a fuscous marginal zone (occasionally with many zones).

On old stumps. Uncommon. Dec.

Name—*crispa*, a curl. From the crisped margin. *Fr. Hym. Eur. p.* 550. *Berk. Out. p.* 243. *C. Hbk. n.* 765. *S. Mycol. Scot. n.* 722.—*Pers. Obs. Myc. ii. p.* 8.

**46. P. amorphus** Fr.—**Pileus** shining white, fleshy-pliant, thin, Polyporus. effuso-reflexed, silky. **Pores** minute, unequal, golden.

Often cræspitose, imbricated, very irregular, also resupinate on ground covered with pine-needles.

On fir-stumps and among pine-leaves. Uncommon. Nov.—Jan.

Name— $\alpha$ , μορφή, form. Shapeless. *Fr. Hym. Eur. p. 550. Berk. Out. p. 243. C. Hbk. n. 767. S. Mycol. Scot. n. 723.—Sow. t. 423. Rostk. 27. t. 12. pores reddish.*

**47. P. adiposus** B. & Br.—*White, here and there acquiring a foxy tinge.* **Pileus** waxy-soft, shortly reflexed, obscurely tomentose; hymenium rather thick. **Pores** small, obtuse at the edge.

Very variable in form, sometimes fixed by the apex, sometimes resupinate. Losing much of its substance in age. Substance not zoned within. Pores when horizontal with the edges even, but often elongated and irregular, not stratose. Turns brownish in drying. Allied to *P. amorphus*, but very distinct.

On the ground beside stumps. Uncommon. Dec.

Name—*adeps*, fat. From its substance. *B. & Br. n. 711. Berk. Out. p. 243. C. Hbk. n. 766. S. Mycol. Scot. n. 724. Fr. Hym. Eur. p. 550.*

**48. P. Keithii** B. & Br.—Shell-shaped, stemless, decurrent behind. **Pileus** about 12 mm. ( $\frac{1}{2}$  in.) across, bright red-brown, rough with rigid tooth-shaped processes; hymenium pallid; dissepiments lacerated.

On fallen sticks. Dunphail.

Name—after Rev. Dr Keith. *B. & Br. n. 1430. S. Mycol. Scot. n. 725.*

### 3. Spongiosi. Spongy when fresh, &c.

\* *Substance coloured.*

**49. P. hispidus** Fr.—**Pileus** 10–15 cent. (4–6 in.) broad, somewhat ferruginous, compact, spongy-fleshy, dimidiate, pulvinate, hispid, internally composed of diverging fibres. **Pores** minute, rounded, somewhat separate, fimbriated, paler than the pileus.

Changing colour. Pores yellowish then fuscous.

On living ash, &c. Common. July–Nov.

**Pileus** sometimes 30 cent. (12 in.) or more broad, 10 cent. (4 in.) thick. *M. J. B.* Name—*hispidus*, hispid. *Fr. Hym. Eur. p. 551. Syst. Myc. i. p. 362. Berk. Out. p. 243. C. Hbk. n. 768. S. Mycol. Scot. n. 726. Grev. t. 14. Krombh. t. 48. f. 7–10. Hussey i. t. 29, 31.—Bull. t. 210, 493. Bolt. t. 161. Sow. t. 345.*

Polyporus.

**50. P. cuticularis** Fr.—**Pileus** ferruginous-fuscos then blackish, thin, spongy-fleshy, then juiceless, becoming plane, hairy, tomentose, *internally laxly composed of parallel fibres*, margin *fibroso-fimbriate, incurved*. **Pores** minute, long, pallid then ferruginous.

Allied to *P. hispidus*, but much thinner, soft, the pores longer than the depth of the flesh. Spores plentiful, ochraceous.

On trunks. Burnham Beeches.

Name—*cuticula*, a thin external skin. *Fr. Hym. Eur. p. 551. B. & Br. n. 1137. C. Hbk. n. 769.—Bull. t. 462.*

**\*\* Substance and spores white.**

**51. P. spumeus** Fr.—Whitish. **Pileus** 7.5 cent. (3 in.) broad, fleshy-spongy, compact, pulvinate, gibbous, rugoso-hispid, *with a stem-like base*, margin incurved; flesh zoned towards the margin. **Pores** separating, minute, round, acute, entire.

A very distinguished species.

On trunks and stumps. Rare. July.

Oozing out from the tree in a very soft mass. *M. J. B.* Name—*spuma*, froth. *Fr. Hym. Eur. p. 552. Syst. Myc. i. p. 358. Berk. Out. p. 244. t. 16. f. 4. C. Hbk. n. 770. S. Mycol. Scot. n. 727.—Sow. t. 211.*

**52. P. borealis** Fr.—White then yellowish, 5 cent. (2 in.) and more broad and thick. **Pileus** spongy then *corky*, compact, somewhat pulvinate, hairy, *internally composed of parallel fibres*, margin spreading. **Pores** adnate, unequal, sinuoso-flexuous and torn, white.

It is exceedingly distinct from those nearest to it; fleshy indeed when young, but when full grown corky. Otherwise it is very variable; the following are to be noted. *B. montanus* pileus fleshy, thick, hairy, margin obtuse; pores obtuse, entire. *C. spathulatus* pileus thin, villous, margin acute, extended into a short lateral stem; pores thin, very much torn.

On stumps. Slough. Balnamoon, Forfarsh. Strachan, Kin-cardinesh.

Name—*borealis*, northern. *Fr. Hym. Eur. p. 552. Syst. Myc. i. p. 366. B. & Br. n. 1429. S. Mycol. Scot. n. 728. Rostk. 4. t. 40.—Schæff. t. 314 (bad).*

*B. Placodermei.* Pileus clothed with a continuous crust, &c.

4. *Suberosi.* Pileus at the first somewhat fleshy and juicy, &c.

**\* Substance coloured.**

**53. P. dryadeus** Fr.—**Pileus** somewhat ferruginous, becoming fuscous, fleshy then corky, thick, pulvinate, cuticle thin, soft,

*rugged becoming even*, smooth; flesh somewhat zoned, ferruginous. **Pores** very long, thin, round, soft, ferruginous, paler at the orifice. Polyporus.

Among the largest and most distinguished, growing rapidly, exuding watery drops at the margin and hence confounded with *P. applanatus*.

On old oaks. Frequent. Aug.-Oct.

Pileus 7.5-17.5 cent. (3-7 in.) or more broad, 6 cent. (2½ in.) thick. Smell strong, subacid. *M. J. B.* Name—*δρύς*, oak. From its habitat. *Fr. Hym. Eur. p. 553. Berk. Out. p. 244. C. Hbk. n. 771. S. Mycol. Scot. Supp. Scot. Nat. 1881, p. 37. Hussey i. t. 21.—Pers. Bull. t. 458.*

**\*\* Substance, which is not zoned, and spores white.**

**54. *P. betulinus* Fr.**—Pileus fleshy then corky, hoof-shaped, obtuse, zoneless, smooth, *the oblique vertex in the form of an umbo*, pellicle thin, separating. **Pores** late of being developed, curt, minute, unequal, at length separating.

On living and dead birch. Common. May-Dec.

Pileus 7.5-15 cent. (3-6 in.) broad. The pileus is at first pale, then acquiring a brownish tinge. The edge is always very obtuse. Name—*betula*, birch. From its habitat. *Fr. Hym. Eur. p. 555. Berk. Out. p. 244. C. Hbk. n. 772. S. Mycol. Scot. n. 730. Grev. t. 246. Rostk. t. 22.—Bull. t. 312. Bolt. t. 159. Sow. t. 212. Fl. Dan. t. 1254.*

**55. *P. quercinus* Fr.**—Pileus tan-pallid, now and then reddish when touched, thick, corky, texture floccose, soft then hardened, tongue-shaped, convexo-plane, even, at first floccoso-granulated, *narrowed behind into a thick horizontal stem*. **Pores** curt, minute, whitish.

A remarkable species, analogous with *Fistulina*.

On old oak. Rare.

Name—*quercus*, oak. From its habitat. *Fr. Hym. Eur. p. 555. Berk. Out. p. 239. C. Hbk. n. 747. Hussey i. t. 52.—Krombh. t. 5. f. 3-5; t. 48. f. 11-14.—Schrader. Spic. p. 157.*

5. Fomentarii. Pileus floccose, of the nature of tinder, &c.

**\* Substance and pores umber or fuscous.**

**56. *P. vegetus* Fr.**—Pileus fuscous, dilated, smooth, opaque, concentrically sulcate, the annual zone broad, texture floccose, lax, very thin, *cuticle of the second season thick, separable*. **Pores** minute, *separating*, umber, those of each season distinct on account of an intermediate floccose stratum formed annually.

It is among the largest, growing rapidly, attaining to 1 ft. in the first year,

Polyporus. but the flesh is very thin, and from it both the cuticle and the very long pores can at length be separated. Plane beneath, at first whitish, margin very smooth, shining, barren, narrow. It has been confounded with *P. applanatus*, but differs by many marks.

On lime-trees. Rare.

Name—*vegetus*, of vigorous growth. *Fr. Hym. Eur. p. 556. Icon. t. 183. f. 1. Berk. Out. p. 245. C. Hbk. n. 774.*

**\*\* Substance and pores somewhat ferruginous.**

**57. P. applanatus Fr.—***Pileus cinnamon* becoming hoary, flattened, tubercular, obsoletely zoned, pulverulent or smooth, *cuticle crustaceous, rigid, at length fragile*, internally very soft, laxly floccose, margin swollen, white then cinnamon. **Pores** very small, somewhat ferruginous, orifice whitish, becoming fuscous when bruised.

Odour acid. When fresh much softer than *P. fomentarius*, &c., but not fleshy or juicy. The pores are softer and shorter.

On trunks, especially ash. Uncommon. Aug.—Oct.

Attaining a large size. Name—*plano*, to level. Flattened. *Fr. Hym. Eur. p. 557. Berk. Out. p. 245. C. Hbk. n. 775. S. Mycol. Scot. n. 731. Kl. Bor. t. 393.—Pers. Rostk. 27. t. 9. Batsch f. 130. Bull. t. 454 f. C?*

**58. P. fomentarius Fr.—***Pileus fuliginous* becoming hoary, hoof-shaped-pulvinate, thick, smooth, remotely and concentrically sulcate, *smooth, opaque, internally soft, floccose*, ferruginous-tawny, cuticle thick, very hard, persistent, margin at first glaucous-pruinose then ferruginous. **Pores** very long, minute, distinctly stratose, at first glaucous-pruinose then ferruginous.

Plane beneath, attenuated towards the margin from the very thick base, hence triangular indeed, but semi-orbicular at the circumference and on all sides sloping downwards. There are two monstrous forms; one stalked, cylindrical (*Batt. t. 37. E. Nees Nov. Act. Nat. Cur. x. with fig.*); another effused, resupinate. It used to be gathered and sold as making *excellent tinder*.

On trunks. Common. Aug.—Nov.

Pileus 10–12.5 cent. (4–5 in.) broad. Sometimes when old, especially when growing on birch, nearly white, occasionally tinged with bright yellow. Mass of pores generally concave. *M. f. B.* Name—*fomentum*, tuel, tinder. *Fr. Hym. Eur. p. 558. Sv. ätl. Sv. t. 62. Berk. Out. p. 245. C. Hbk. n. 776. S. Mycol. Scot. n. 732. Rostk. 4, t. 52.—Linn. Lenz f. 48. Sow. t. 133 (first year).*

**59. P. nigricans Fr.—***Pileus black*, pulvinate, very thick, densely and concentrically sulcate, *smooth, shining*, the very hard crustaceo-torn crust persistent, internally very hard, ferruginous, mar-



gin very obtuse, ferruginous. **Pores** very small, plane, confluent- Polyporus.  
stratose, naked, ferruginous.

The typical form with the pileus even, varnished, and shining, presents a most distinct species, but with this is confounded a trivial form with the pileus triangular, rugose, opaque, which approaches *P. igniarius*. *Rostk. t. 51. Quel. t. 19. f. 3.*

On living and dead birch. Uncommon. May-Oct.

5-15 cent. (2-6 in.) broad. Sometimes exactly the shape of a horse's hoof. Name—*niger*, black. Blackish. *Fr. Hym. Eur. p. 558. Icon. t. 184. f. 2. Berk. Out. p. 245. C. Hbk. n. 777. S. Mycol. Scot. n. 733.*

**60. P. igniarius** Fr.—**Pileus** at first tuberculoso-globose (immarginate), even, with a thin, flocculose, adpressed, hoary covering, then hoof-shaped, ferruginous then fuscous-blackish, opaque, cuticle concrete, roughly uneven, very hard; flesh zoned, ferruginous, very hard, margin rounded. **Pores** very small, convex, stratose, cinnamon, when old stuffed with a white substance, at the first hoary.

It differs from preceding species in its entire nature and life-history. The new growth, which is vernal, makes smaller specimens throughout, and larger ones only at the margin and on the hymenium (which is also at first destitute of pores) floccoso-hoary, then ferruginous, at length dark. It varies resupinate: *Mich. t. 62.* Unsuitable for tinder, but used for burning.

On ash, poplar, willow, plane, cherry. Common. May-Nov.

Pileus narrower and thicker than *P. fomentarius*, resembling a horse's hoof. Mass of tubes in general plane or very convex. *M. F. B.* Name—*ignis*, fire. From its being used for burning. *Fr. Hym. Eur. p. 559. Berk. Out. p. 246. C. Hbk. n. 778. S. Mycol. Scot. n. 734. Lenz f. 47. Ventur. t. 61. f. 4. Rostk. 4. t. 54.—Linn. Bull. t. 454. Bolt. t. 80. Sow. t. 132.*

**61. P. fulvus** Fr.—**Pileus** tawny, at length becoming hoary, woody-corky, very hard, exactly triangular, even (not concentrically sulcate), at first hairy or villous. **Pores** curt, round, minute, cinnamon, at first covered with cinereous-yellow pruina.

Pileus convex on both sides, adnate by a broad base, not very obtuse at the margin. Pores not distinctly stratose.

On decayed trunks. Rare.

Name—*fulvus*, tawny. *Fr. Hym. Eur. p. 559. Icon. t. 184. f. 3. B. & Br. n. 1138. C. Hbk. n. 779.*

**62. P. pectinatus** Kl.—**Pileus** ferruginous-fuscous, corky-woody, hard, triangular, concentrically lamellato-plicate above, tomentoso-scaly; margin pale-yellowish, naked. **Pores** very small, curt, obtuse, pale-yellowish, naked.

It is doubtful whether the margin and pores are always naked.

Polyporus. Habitat not recorded. Penzance.

Name—*pecten*, a comb. From the markings on the pileus. *Klotsch*.—*Fr. Hym. Eur. p.* 559. *B. & Br. n.* 1965.—*Quel. t.* 17. *f.* 5.

**63. P. conchatus** Fr.—**Pileus** *date-brown*, corky inclining to the nature of tinder, thin, effused, *somewhat shell-shaped-reflexed*, concentrically sulcate, *tomentose*, margin acute. **Pores** curt, very small, the thin trama of the same colour.

Pileus concave beneath. It varies wholly resupinate.

On trunks, willow, &c. Frequent.

Distinguished from *P. ribis* by its harder substance and smoother pileus. *M. J. B.* Name—*concha*, a shell. From its shape. *Fr. Hym. Eur. p.* 560. *Berk. Out. p.* 246. *C. Hbk. n.* 781. *S. Mycol. Scot. n.* 735.—*Pers. Myc. Eur. ii. p.* 85.

**64. P. ribis** Fr.—**Pileus** as much as 10 cent. (4 in.) broad, ferruginous then *umber*, corky-coriaceous, *becoming plane*, rather even, velvety, tawny internally and at the acute margin. **Pores** curt, 2 mm. (1 lin.) long, very small, naked, tawny.

Perennial. Imbricated, somewhat zoned.

On currant and gooseberry bushes. Common in England.

Name—*ribis*, currant. *Fr. Hym. Eur. p.* 560. *Syst. Myc. i. p.* 375. *Berk. Out. p.* 246. *C. Hbk. n.* 780. *Rostk. t.* 53. *Fl. Dan. t.* 1790. *f.* 2. *Corda Sturm. xi. t.* 62.—*Schum. Bull. t.* 454. *f.* E.

**65. P. salicinus** Fr.—**Pileus** *cinnamon then hoary*, woody, very hard, undulated, *smooth*, for the most part resupinate, margin short, obtuse, spreading. **Pores** *very small*, round, ferruginous-cinnamon.

Pileus 30 cent. (12 in.) and more, often interrupted. Allied to *P. fomentarius* or *igniarius*, with an ebony-black crust. From its situation wholly resupinate or in vertical places encircled above with a narrow, undulated, smooth margin.

On willows. Common. Autumn.

Name—*salix*, willow. *Fr. Hym. Eur. p.* 560. *Syst. Myc. i. p.* 376. *Icon. t.* 185. *f.* 1. *Berk. Out. p.* 246. *C. Hbk. n.* 782. *S. Mycol. Scot. n.* 736.

\*\*\* *Substance white or pallid* (except *P. roseus*).

**66. P. roseus** Fr.—**Pileus** 5–10 cent. (2–4 in.) broad, 1–2.5 cent. (½–1 in.) thick, *externally and internally rose-colour*, corky-woody, hard, triangular, even, somewhat banded, externally as if blown upon with cinereous-blackish powder, internally floccosofibrous. **Pores** minute, round, rose-colour.

Somewhat caespitose.

On dressed wood in conservatory. Glamis, 1877. Autumn. Polyporus.

Name—*roseus*, rose-colour. *Fr. Hym. Eur. p. 562. Syst. Myc. i. p. 372. B. & Br. n. 1801. S. Mycol. Scot. n. 737.—A. & S. p. 251.*

**67. P. ulmarius** Fr.—**Pileus** 7.5–10 cent. (3–4 in.) broad, *in the first season white*, corky-woody, very hard, incrustated, tubercular, *becoming smooth*, substance white. **Pores** round, *stratose, yellowish*.

A. *first season*, pileus white. *Sow. t. 88.* B. In succeeding seasons, pileus blackish, margin tawny. *Berk. Out. t. 16. f. 5.* The two figures are so unlike that one would scarcely suppose them to be allied.

On old elm. Frequent. Sept.

Name—*ulmus*, elm. *Fr. Hym. Eur. p. 562. Syst. Myc. i. p. 365. Berk. Out. p. 246. C. Hbk. n. 783. S. Mycol. Scot. n. 738. Hussey i. t. 64. Sow. & Berk. quoted above.*

**68. P. cytisinus** Berk.—Large, 30 cent. (12 in.) or more across, imbricated. **Pileus** coarsely tuberculated, hard, woody, margin subacute. **Pores** minute, nearly white.

Dimidiate, quite smooth at least when dry; substance slightly zoned, very thick and close, but composed of velvety fibres, pale, nearly white, evidently composed of two or three successive layers.

On laburnum. Rare.

Name—*cytisus*, laburnum. *Berk. Eng. Fl. v. p. 142. Out. p. 247. C. Hbk. n. 785. S. Mycol. Scot. n. 739. Fr. Hym. Eur. p. 562.—Sow. t. 288.*

**69. P. connatus** Fr.—**Pilei** white or cinereous, corky-woody, effuso-reflexed, *imbricated like the steps of a ladder, growing into each other, villous*, internally white. **Pores** *stratose*, minute, somewhat round, white.

Increasing from season to season and distinctly stratose, pileus commonly covered with mosses growing upon it.

On old trunks, elm, crab-tree, &c. Frequent. Autumn.

Running up trunks for 1 or 2 ft. *M. J. B.* Name—*con*, together; *natus*, born. Growing together. *Fr. Hym. Eur. p. 563. Icon. t. 185. f. 2. Berk. Out. p. 248. C. Hbk. n. 789. S. Mycol. Scot. n. 740. Batt. t. 37. f. G.*

6. Lignosi. Pileus from the first hard, woody, &c.

**70. P. fraxineus** Fr.—**Pileus** white then rubiginous and fuscous, corky-woody, smooth, somewhat plane, zoneless, at first even, then concentrically sulcato-plicate, pallid within. **Pores** minute, curt, *rufous-ferruginous, at first, as well as the margin, covered over with white greasy villous down.*

Variable in form, softer when young, but not fleshy, truly perennial.

Polyporus. On old ash. Frequent. May-Nov.

Solitary or imbricated. Smell strong and penetrating. *M. J. B.* Spores ovoid-sphaeroid, 6-7 mk. *Q.* Name—*fraxinus*, ash-tree. *Fr. Hym. Eur. p. 563. Berk. Out. p. 247. C. Hbk. n. 784. S. Mycol. Scot. n. 741.—Bull. t. 433. f. 2.*

**71. P. variegatus** Secr.—**Pileus** variegated orange and bay-brown, corky-woody, somewhat plane, even, smooth, *zoneless*, *shining*, pallid within. **Pores** round, minute, curt, unequal and torn, *yellowish*.

With the habit of *P. varius*, but decurrent at the base as is usual in this section. Sowerby compares it with *P. lucidus* as regards its substance.

On trunks. Rare.

Name—*variegatus*, variegated. From the colours. *Secr. n. 45. Fr. Hym. Eur. p. 563. Berk. Out. p. 247. C. Hbk. n. 786.—Sow. t. 368.*

**72. P. carneus** Nees.—**Pileus** *flesh-colour*, of the same colour internally, effuso-reflexed, woody, hard, thin, rugose, *smooth*, *zoneless*. **Pores** minute, round, decurrent at the base.

Smaller and thinner than *P. fraxineus* and *P. variegatus*, but hard and perennial.

On old stump. Welshpool, 1871. Nov.

Name—*caro*, flesh. *Flesh-colour. Nees Nov. Act. Nat. Curios. xlii. t. 3. Fr. Hym. Eur. p. 563. B. & Br. n. 1366.*

**73. P. annosus** Fr.—**Pileus** woody, convex then becoming plane, *rugoso-tubercular*, in the first season brown and silky, in the second and subsequent seasons covered with a rigid, smooth, blackish crust, white within. **Pores** of medium size, whitish.

A. **Pileus** reflexed, pores obtuse. B. **Pileus** wholly resupinate, pores unequal. Very variable in form, but very distinct.

About roots of stumps, &c. Very common. April-Dec.

Forming masses 3 in.—1½ ft. wide of imbricated, rugged, or radiato-striate brown pilei, marked with strong vaulted zones. *M. J. B.* Name—*annus*, a year. From its growing yearly. *Fr. Hym. Eur. p. 564. Icon. t. 186. f. 2. Berk. Out. p. 247. C. Hbk. n. 788. S. Mycol. Scot. n. 742.—Rostk. 4. t. 29.*

**74. P. populinus** Fr.—*White. Pileus* 12 mm. (½ in.) broad, corky-woody, rigid, *zoneless*, *villous*, of the same colour internally, *margin obtuse*. **Pores** minute, curt, round.

Imbricated, growing into each other at the decurrent base, transversely dilated. Not stratose.

On trunks of poplar. Uffington, Lincolnshire.

Very much the habit of *P. connatus*. *B. & Br.* Name—*populus*, poplar. *Fr. Hym. Eur. p. 564. Syst. Myc. i. p. 367. B. & Br. n. 1288.*

C. Inodermei. Pileus from the first arid and firm, &c.

Polyporus.

7. Stupposi. Pileus flocculose then smooth, &c.

\* *Substance coloured.*

**75. P. radiatus** Fr.—**Pileus** 2.5 cent. (1 in.) broad, corky, coriaceous, rigid, *radiato-rugose*, at first *velvety*, *tawny*, then becoming smooth, ferruginous-fuscos, margin spreading, repand. **Pores** minute, pallid, silvery-glistening, at length ferruginous.

Very much imbricated, growing into each other, becoming woody, but not perennial.

On hazel, alder, &c. Frequent. Oct.—Feb.

Name—*radiatus*, radiate. From the wrinkles on the pileus. *Fr. Hym. Eur. p. 565. Syst. Myc. i. p. 369. Berk. Out. p. 248. C. Hbk. n. 790. S. Mycol. Scot. n. 743. Kl. t. 461.—Sow. t. 196. Batt. t. 39. C.*

**76. P. polymorphus** Rostk.—**Pileus** *umber*, resupinate, effused, coriaceous, margin reflexed, *crisped*, *smooth*. **Pores** rather large, angular and torn, pallid.

The above description is from the figure. From the description of the author it is wholly bay-brown-pallid.

On fir in a fence. Forres. Aug.

Name—*πολύς*, many; *μορφή*, form. *Rostk. 4. t. 56. Fr. Hym. Eur. p. 566. B. & Br. n. 1802. S. Mycol. Scot. n. 744.*

**77. P. cryptarum** Fr.—**Pileus** *rufescent-ferruginous*, corky-cottony, effuso-reflexed and variable, zoneless, *adpressedly* silky, paler internally. **Pores** very long, minute, round, paler than the pileus.

Exceedingly changeable in form and colour, becoming paler with the weather.

On fir-trees. Frequent.

Nearly related to *P. ferruginosus*, and perhaps a mere state of it. *B. & Br.* Name—*crypta*, a vault, cellar. Probably first found in cellar. *Fr. Hym. Eur. p. 566. B. & Br. n. 1804. Pers. Myc. Eur. 2. t. 16. f. 3.—Bull. t. 478.*

\*\* *Substance white.*

**78. P. gossypinus** Lev.—*White*. **Pileus** coriaceous, effuso-reflexed, becoming plane, *tomentose*, zoneless, substance of the same colour. **Pores** at first *labyrinthine*, then angular, pallid *cinereous*, dissepiments thin, slightly toothed.

The pores are three or four times longer than the flesh of the pileus is thick.

Polyporus. On old stumps of *Ulex*. Sibbertoft.

Name—*gossypium*, cotton. *Leveill. Ann. Sc. Nat.* 1843, p. 124.—*Fr. Hym. Eur.* p. 566. *B. & Br. n.* 2024.

**79. P. fibula** Fr.—Whitish. **Pileus** coriaceous, *soft*, tough, *velvety-hairy*, zoneless, often radiato-rugose, internally snow-white, margin entire, acute. **Pores** small, rounded, acute, at length torn, yellowish.

Small, adnate behind, everywhere free at the circumference, sometimes forming orbicular shields, sometimes variable in form from growing into each other.

On the door of a wine-cellar. Rare.

Name—*fibula*, a button. From its shape. *Fr. Hym. Eur.* p. 567. *Berk. Out.* p. 248. *C. Hbk. n.* 791. *Sow. t.* 387. f. 8.

8. Coriacei. **Pileus** coriaceous, villous, &c.

**80. P. hirsutus** Fr.—**Pileus** *unicolorous*, but zoned with concentric furrows, whitish, corky-coriaceous, convexo-plane, *shaggy with rigid hairs*. **Pores** round, obtuse, whitish, becoming somewhat fuscous.

B. Pores wholly white. C. Pores yellow at the orifice, but white within. D. Pores obtuse, angular. E. Margin of the pileus ferruginous (*Fl. Dan. t.* 2079. f. 1). F. pileus blackish.

On dead trunks. Near Twycross.

Name—*hirsutus*, hairy. *Fr. Hym. Eur.* p. 567. *B. & Br. n.* 1139. *C. Hbk. n.* 792.—*Wulf. & Jacq. Coll. ii.* p. 149.

**81. P. velutinus** Fr.—**Pileus** about 5 cent. (2 in.) broad, *white* at length yellowish, corky-coriaceous, plane on both sides, *velvety-soft, slightly zoned*, the extenuated margin acute. **Pores** round, minute, thin, white.

Damp when young, then hardened, not shining like *P. versicolor*, the zones only a little darker.

On stumps, branches, &c. Uncommon. Nov.

Sometimes entirely white, and not unfrequently the zones are mere depressions. *M. f. B.* Name—*vellutetum* (low Latin). *Velvety.* *Fr. Hym. Eur.* p. 568. *Syst. Myc. i.* p. 368. *Berk. Out.* p. 248. *C. Hbk. n.* 793. *S. Mycol. Scot. n.* 745.


**82. P. versicolor** Fr.—**Pileus** *variegated with differently coloured zones*, coriaceous, thin, rigid, becoming plane, depressed behind, *becoming even, velvety, shining*. **Pores** minute, round, acute and torn, white, becoming pale (somewhat yellowish).

Varying endlessly both in form and colour.

On dead wood. Exceedingly common. April-Dec.

Polyporus.

More or less imbricated. Pileus 5-7.5 cent. (2-3 in.) broad. Name—*versicolor*, changeable in colour. *Fr. Hym. Eur. p. 568. Berk. Out. p. 248. C. Hbk n. 794. S. Mycol. Scot. n. 746. Hussey i. t. 24.—Linn. Bull. t. 86. Bolt. t. 81. Sow. t. 229. Fl. Dan. t. 1554. Schæff. t. 263. Sterb. t. 27. K. Batt. t. 35. A.*



LXVI. *Polyporus versicolor*.  
One-half natural size.

**83. P. abietinus** Fr.—Pileus cinereous-white, coriaceous, thin, effuso-reflexed, villous, obsoletely zoned. Pores unequal, torn, violaceous becoming pale.

There is no species more easily distinguished, though it is sometimes wholly resupinate, sometimes reflexed, if only once it has been rightly identified when in full vigour, and if distinct species are not sought for in old (washed out) and dry specimens.

On fir. Common. July-Nov.

Fries says it is "the mother of vast confusion." Pileus 2.5-5 cent. (1-2 in.) broad. Often stained with minute green *Algæ*. *M. J. B.* Name—*abies*, fir. *Fr. Hym. Eur. p. 569. Berk. Out. p. 249. C. Hbk n. 795. S. Mycol. Scot. n. 747. Grev. t. 226. Fl. Dan. t. 2079. f. 2.—Dicks. Crypt. 3. t. 9. f. 9. Fl. Dan. t. 1298.*

**84. P. Wynnei** B. & Br.—Confluent, variable in form. Pileus tan-colour, adnate behind, effuso-reflexed, marked with silky raised lines. Pores small, angular, white.

Incrusting various substances, with the margin more or less broadly reflected. Pores  $\frac{1}{8}$  in. across, acquiring a slight tint like that of the pileus in drying. Somewhat of the habit of *P. amorphus*, but not so fleshy in texture.

Running over twigs, grass, &c. Coed Coch, &c.

Name—after Mrs Lloyd Wynne. *B. & Br. n. 807. Berk. Out. p. 249. C. Hbk n. 796. Fr. Hym. Eur. p. 569.*

V. RESUPINATI. Pileus none, &c.

\* Pores fuscous or blackish.

No British species.

\*\* Pores ferruginous or cinnamon.

**85. P. umbrinus** Fr.—Rufous-umber, resupinate, determinate, undulato-tubercular, defined by a smooth, paler margin. Pores

*Polyporus*. minute, somewhat round, unequal, of the same colour as the pileus.

Clearly different from *P. ferruginosus* and following species.

On trunks. Knowle Park, Burchell.

Name—*umbrinus*, umber-coloured. *Fr. Hym. Eur. p. 571. B. & Br. n. 1890.—Rostk. 27. t. 6.*

**86. *P. ferruginosus* Fr.**—As much as 2.5 cent. (1 in.) thick, *tawny*, date-brown-ferruginous when old, effused, firm, unequal, *circumference sterile*. **Pores** of medium size, very long, somewhat round and torn, cinnamon.

Composed chiefly of the elongated pores.

On logs, posts, &c. Common. May–Nov.

Mycelium ochraceous. At first appearing under the form of a confervoid stratum, which gradually thickens and acquires pores in the centre. *M. f. B.* Name—*ferrum*, iron. Rust-coloured. *Fr. Hym. Eur. p. 571. Berk. Out. p. 249. C. Hbk. n. 798. S. Mycol. Scot. n. 748. Grev. t. 155.—Schrader. Spic. p. 172.*

**87. *P. contiguus* Fr.**—Almost 7.5 cent. (3 in.) long, 2.5 cent. (1 in.) broad, 12 mm. ( $\frac{1}{2}$  in.) thick, cinnamon when young, effused, firm, *smooth, somewhat marginate*, margin at the first villous. **Pores rather large**, equal, obtuse, entire.

Smaller than *P. ferruginosus*; distinguished by its ochraceous mycelium from resupinate forms of *P. igniarius* (of which the mycelium is white according to Berkeley).

On rotten wood and sticks. Uncommon.

Name—*contiguus*, continuous, uninterrupted. *Fr. Hym. Eur. p. 571. Syst. Myc. i. p. 378. Berk. Out. p. 249. C. Hbk. n. 797.—Pers. Syn. p. 544.*

\*\*\* *Pores purple or violaceous.*

**88. *P. violaceus* Fr.**—*Violaceous*, effused, determinate, thin, short, even, *smooth*, closely adnate, destitute of a distinct subiculum. **Pores** short, *cellular or veined*, quite entire.

It approaches the *Merulii* in the curt, cellular pores.

On fir poles and stumps. Frequent. Jan.–May.

Spores ovoid, punctate, rose-yellow, 7 mk. *Q.* Name—*violaceus*, violet-colour. *Fr. Hym. Eur. p. 572. B. & Br. n. 1021. C. Hbk. n. 803. S. Mycol. Scot. n. 749.—Rostk. 27. t. 3.*

**89. *P. purpureus* Fr.**—Very broadly and widely effused, 10–30 cent. (4–12 in.) long and broad, *the mucedinous, flocculose, white mycelium* creeping over the surface of rotten wood. **Pores** curt,



scarcely 2 mm. (1 lin.) long, minute, unequal, *interruptedly* Polyporus,  
*scattered* or conglomerate, purple-lilac.

On decayed willow. Cotterstock, Norths.

Name—*purpureus*, purple. *Fr. Hym. Eur. p. 572. Syst. Myc. i. p. 379. Berk. Out. p. 250. C. Hbk. n. 804.*

\*\*\*\* *Pores red.*

**90. P. incarnatus** Fr.—Flesh-colour, effused, 2.5–10 cent. (1–4 in.) long, scarcely 2.5 cent. (1 in.) broad, *corky-coriaceous*, persistent, firm, smooth. **Pores** *elongated*, unequal, commonly oblique.

The superior margin here and there reflexed. Pores oblique from position.

On fir. Rare. June–Nov.

Sometimes small cottony protuberances occur amongst the pores which have the appearance of small pilei, with tubes underneath. *Grev.* Name—*incarnatus*, flesh-colour. *Fr. Hym. Eur. p. 573. Syst. Myc. i. p. 379. Icon. t. 189. f. 1. Berk. Out. p. 250. C. Hbk. n. 802. S. Mycol. Scot. n. 750. A. & S. p. 250.*

**91. P. rufus** Fr.—*Blood-red-rufous*, effused, coriaceous, thin, *adnate*, even, smooth, *determinate*. **Pores** very small, thin, acute.

On broom. Strachan, Kincardineshire.

Name—*rufus*, reddish. *Fr. Hym. Eur. p. 573. S. Mycol. Scot. n. 751. Rostk. 4. t. 62.*

**92. P. rhodellus** Fr.—*White-flesh-colour*, effused, 5 cent. (2 in.) and more long, thin, *adnate*, *soft*, *margin* *determinate*, *naked*. **Pores** minute, somewhat round, continuous.

Thinner than preceding species, apparently nearest to *P. vulgaris*.

In Scotch fir wood. Glamis, 1874. Aug.

Name—*ῥόδον*, a rose. Slightly rose-colour. *Fr. Hym. Eur. p. 573. Syst. Myc. i. p. 380. Icon. t. 189. f. 2. B. & Br. n. 1806. S. Mycol. Scot. n. 752. Bull. t. 442. f. D.*

**93. P. micans** Fr.—*Whitish-flesh-colour*, effused, thin, *adnate*, *soft*, *circumference flaxy and white*. **Pores** very thin, *resembling honeycomb*, angular, somewhat crenate.

Somewhat orbicular, confluent, fugacious. In the structure of the pores it approaches *P. bombycinus*.

On dead wood, ash. Leighwood, Bristol. Altyre. Nov.

Name—*mico*, to glitter. *Fr. Hym. Eur. p. 573. B. & Br. n. 1289. S. Mycol. Scot. n. 753. Ehrenb. Silv. Berol. p. 30.*

Polyporus.

\*\*\*\*\* Pores yellowish.

**94. P. nitidus** Fr.—*Yellow*, effused, thin, somewhat adnate, defined, *villosa-marginate at the circumference*. **Pores** minute, curt, *round*, equal, shining.

Remarkable for its bright, almost golden colour, and somewhat regular form.

On dead wood. Rare. March.

Name—*nitidus*, shining. From the pores. *Fr. Hym. Eur. p. 574. Berk. Out. p. 250. C. Hbk. n. 799. S. Mycol. Scot. n. 754—A. & S. p. 258.*

**95. P. Læstadii** Fr. & Berk.—Substance white, hymenium here and there tubercular. **Pores** bright lemon-yellow.

Colour bright persistent yellow, very beautiful. Forming confluent patches, many inches in length.

On the under side of a deal board in hothouse. Coed Coch.

Name—after C. P. Læstadius. *B. & Br. n. 2025.*

**96. P. bombycinus** Fr.—*Dingy-yellowish*, effused, *silky-membranaceous*, adhering laxly, *spider-web-velvety at the circumference*. **Pores** large, angular.

Very distinct and singular. Pores in their first stage forming only small pits, which are sunk in the mycelium, then developed, wholly formed from the tomentum, somewhat round, at length firmer, angular and here and there flexuous.

On dead wood. Rare.

Name—*bombyx*, silk. *Fr. Hym. Eur. p. 575. Berk. Out. p. 250. C. Hbk. n. 801.—Sow. t. 387. f. 5.*

**97. P. aneirinus** Sommerf.—*White*, effused, thin, somewhat innate, *flaxy at the circumference*. **Pores** large, *cell-like*, waxy, angular, smooth, tawny.

The growth is at first wholly flaxy; the formation of pores is the same as in *P. bombycinus*, but they are at length waxy and often exactly hexagonal.

On dead branches. Rare. Oct.

Name—*ἀν, εἰπίρεος*, woolly. Not woolly. From the pores becoming waxy as contrasted with those of *P. bombycinus*, which are persistently downy. *Sommerf. Lapp. p. 276.—Fr. Hym. Eur. p. 575. Berk. Out. p. 252. C. Hbk. n. 816. S. Mycol. Scot. n. 755.*

**98. P. ramentaceus** B. & Br.—Somewhat orbicular; subiculum white, tomentose, margin obsolete. **Pores** honey-colour, large,  $\frac{1}{32}$  in. across, somewhat hexagonal, dissepiments thin, slightly rigid, acute.

On Scotch fir. Glamis, 1874. Sept.

Polyporus.

Name—*ramenta*, shavings, chips. On chips. *B. & Br. n.* 1809. *S. Mycol. Scot. n.* 774.

\*\*\*\*\* *Pores white then changing colour.*

**99. P. cinctus** Berk.—White, turning pallid, forming *small erect scattered tufts*, each surrounded by radiating strigose fibres. **Pores** extremely minute, angular, dissepiments extremely thin, edge ragged.

Some of the patches are barren, and some at length become confluent. Under favourable circumstances a distinct hymenium, 3 mm. ( $1\frac{1}{2}$  lin.) thick, with a free, even, abrupt, vertical circumference is formed in the centre of each tuft.

On old deal boards. Rare. King's Cliffe.

Pores at length pallid ochraceous, darker when dry. Quite peculiar in its structure. *Fr.* Name—*cingo*, to encircle. From the fibres surrounding the tufts. *Berk. Out. p.* 250. *Mag. Zool. & Bot. i. t. 2. f.* 3. *C. Hbk. n.* 805. *Fr. Hym. Eur. p.* 575.

**100. P. armeniacus** Berk.—Suborbicular, confluent, extremely thin, circumference minutely downy. **Pores** shallow, minute, nearly round, at first white then bright buff, often confined to the centre, the marginal portion being of a byssoid structure under a lens.

Forming broadly effused patches, composed evidently of many confluent orbicular individuals.

On fir bark. Appin.

Name—of the colour of apricot (*Armeniacum*). *Berk. Eng. Fl. v. p.* 147. *Out. p.* 250. *C. Hbk. n.* 800.

**101. P. Rennyi** B. & Br.—Subiculum thick, pulvinate, pulverulent, forming a thick, at first somewhat frothy then pulverulent mass, white, turning to lemon-colour when dry. **Pores** sparingly produced, white, elongated, dissepiments thin.

On Scotch fir stump and running on the ground. Hereford, 1873. Glamis, 1873. Nov.

Name—after J. Renny. *B. & Br. n.* 1432. *S. Mycol. Scot. n.* 756.

**102. P. subfusco-flavidus** Rostk.—White then light yellow-fuscous, broadly effused by becoming confluent, thin, coriaceous, arid, adnate, margin thin, white-flaxy, determinate. **Pores** minute, irregular.

On oak planks. Roof of King's Cliffe Church.

The pores appear when viewed one way of a greyish-brown, and the other

Polyporus. white. *B. & Br.* Name—from the colours. *Rostk. 27. t. 11. B. & Br. n. 1022. Fr. Hym. Eur. p. 576. P. corticola C. Hbk. n. 817.*

**103. *P. viridans* Berk.**—At first white, then when dry *pallid green*, effused, crustaceo-adnate, thin, *margin pulverulentotomentose*. **Pores** minute, angular, dissepiments thin.

Habit that of *P. vulgaris*.

On rotten wood. Rare. Sept.

Name—*virido*, to be green. Greenish. *B. & Br. n. 347. C. Hbk. n. 810. Fr. Hym. Eur. p. 576.*

**104. *P. terrestris* Fr.**—White, effused, *spider-web-flaxy*, rather tender, *fugacious*. **Pores** central, extremely small, white then *rufescent*.

A doubtful and scarcely typical form.

On naked soil. Uncommon. Sept.—Oct.

Extremely delicate, vanishing when touched. Name—*terra*, earth. Growing on the soil. *Fr. Hym. Eur. p. 576. Berk. Out. p. 252. C. Hbk. n. 814. S. Mycol. Scot. n. 757.—Dec. Fl. fr. 6. p. 39.*

**105. *P. subgelatinosus* B. & Br.**—Orbicular, forming little pulvinate masses with an obtuse raised border, which is at first tomentose and pallid, becoming black, of a somewhat gelatinous consistence. **Pores** pale delicate grey, with an acute even edge, about  $\frac{1}{10}$  in. diameter.

On dead wood, parasitic on *P. amorphus*. Rannoch.

Name—*sub-gelatinosus*, somewhat gelatinous. *B. & Br. n. 1569. S. Mycol. Scot. n. 758.*

\*\*\*\*\* *Pores persistently white, &c.*

**106. *P. medulla-panis* Fr.**—White, effused, *determinate, somewhat undulated, firm*, smooth, *the naked circumference somewhat marginate*, almost wholly formed of the longish, medium-sized, entire pores.

Annual, moderately large, becoming hard when dry, separating. Softer and watery-soft when on the ground.

On decaying branches. Uncommon.

Becoming yellowish when old. Name—*medulla*, pith, &c.; *panis*, bread. From its likeness to soft crumb of bread. *Fr. Hym. Eur. p. 576. Icon. t. 190. f. 2. Berk. Out. p. 251. C. Hbk. n. 806. S. Mycol. Scot. n. 759.—Pers. Syn. p. 544.*

**107. P. mucidus** Fr.—White becoming pale, effused, *rather* Polyporus. *thick*, somewhat immersed, soft, *the indeterminate circumference flaxy*. **Pores** medium-sized (seated on the crust formed of the mycelium), unequal and torn.

It varies softer and firmer according to place, but is moderately persistent. Much larger than *P. molluscus*, and not allied to it.

On decayed fir. Penzance.

Name—*mucus*, mucous. Mucid. From its substance. *Fr. Hym. Eur. p. 577. B. & Br. n. 1967.—Pers. Syn. p. 546.*

**108. P. vitreus** Fr.—Whitish, somewhat hyaline, broadly and unequally effused, rather thick, somewhat undulated, indeterminate, the thin villous margin shining white; *mycelium forming a woody stroma, tough and separable*. **Pores** very small, round, long, obtuse, entire.

The pores are straight or oblique according to situation.

On dead wood, especially fir. Uncommon. Aug.—Nov.

Distinguished by its distinct xylostromatoid substratum, which separates easily from the matrix. *B. & Br. Name—vitrum*, glass. Glassy. From its shining appearance. *Fr. Hym. Eur. p. 577. Syst. Myc. i. p. 381. Berk. Out. p. 251. C. Hbk. n. 807. S. Mycol. Scot. n. 760.—Pers. Myc. Eur. 2. p. 112.*

**109. P. obducens** Pers.—White, effused, incrusting, innate, firm, wholly formed of very small, crowded, equal, *distinctly stratose pores*, the annual strata pallid-tan.

In the first year the plant resembles *P. vulgaris*, but in succeeding years it becomes thick from the numberless strata (2 mm., 1 lin. thick) placed one above another, altogether concrete with the mouldy wood which is pervaded with the mycelium. In vertical places it puts forth rudimentary pileoli. Allied to *P. connatus*.

On rotten trunks. Rare. Bristol.

Name—*ob*, and *duco*, to lead. Covering over. Probably from the strata covering each other. *Pers. Myc. Eur. 2. p. 104.—Fr. Hym. Eur. p. 577. Berk. Out. p. 251. C. Hbk. n. 808.*

**110. P. callosus** Fr.—White, circumference similar, widely effused, even, tough, *entire, separable* like soft leather. **Pores** *firm*, round, equal, quite entire, *obtuse*.

The pores are seated upon a skin, 2–4 mm. (1–2 lin.) thick, firm, but not hard, everywhere porous. Annual and not stratose.

On dead wood. Glamis, 1874. April.

Spores pruiniform, hyaline, 6 mk. *Q.* Name—*callum*, a hardened skin. From the skin on which the pores are seated. *Fr. Hym. Eur. p. 577. B. & Br. n. 1431. S. Mycol. Scot. n. 761.*

Polyporus.


**111. P. vulgaris** Fr.—White, broadly effused, thin, 1 m. ( $\frac{1}{2}$  lin.) thick, arid, *closely adnate*, even, *soon smooth at the circumference*, wholly consisting of firm, crowded, very small, round, somewhat equal pores.

The colour is sometimes light yellow. The pores vary oblique and gaping open. It cannot be separated from the wood except in fragments.

On dead wood. Common. April-Dec.

Effused sometimes to the breadth of 30 cent. (12 in.) *M. J. B.* Name—*vulgaris*, common. *Fr. Hym. Eur. p. 577. Syst. Myc. i. p. 381. Berk. Out. p. 251. t. 16. f. 6. C. Hbk. n. 809. S. Mycol. Scot. n. 762. Rostk. t. 60.*

**112. P. molluscus** Fr.—*White, effused, thin, soft, the flaxy circumference radiating in fibrils. Pores collected in the centre or now and then in patches, very small, thin, round, unequal and torn, pale.*



There is a variety with the circumference membranaceous sterile and continuous. *Sow. t. 387. f. 9.*

On dead wood. Uncommon. Sept.—Nov.

At first forming a mere fringed bysoid membrane. The partitions of the pores are so thin that they very generally become torn. *M. J. B.* Name—*molluscus*, soft. *Fr. Hym. Eur. p. 578. Berk. Out. p. 251. C. Hbk. n. 812. S. Mycol. Scot. n. 763. Fl. Dan. t. 1299.*

LXVII. *Polyporus molluscus*.  
Natural size.

**113. P. collabefactus** B. & Br.—Stratum quite smooth, resembling a *Corticium*. Pores seem at first to arise from the mere collapsing of the substance, short, margin obtuse.

The barren parts resemble exactly a very smooth *Corticium* like *C. calceum*.

On dead wood. Glamis, 1873. Oct.

Name—*collabefo*, to be brought to ruin. From the pores arising out of the collapsing of the substance. *B. & Br. n. 1432. S. Mycol. Scot. n. 764.*

\*\*\*\*\* Pores persistently white, unequal, &c.

**114. P. sanguinolentus** Fr.—*Whitish but bleeding when touched*, nodulose, soon confluent, effused, soft, the flaxy circumference vanishing. Pores small, somewhat round, unequal, at length torn.

Mycelium emerging and forming nodules, which are at first flaxy then Polyporus. smooth, in which the pores nestle, but it does not result in a pileus. Pores very changeable in form.

On dead branches. Uncommon. Sept.—Oct.

Name—*sanguis*, blood. Bleeding. *Fr. Hym. Eur. p. 578. B. & Br. n. 1290. C. Hbk. n. 811. S. Mycol. Scot. n. 765.—A. & S. p. 257.*

**115. P. radula** Fr.—*White*, effused, *made up of the naked tomentose mycelium* closely compacted, soft, villous beneath. **Pores** medium-sized, angular, toothed, pubescent when young.

Its growth is that of *P. sanguinolentus*, but it is drier, more lax, and separable from the matrix.

On fir. Strachan, Kincardineshire.

Name—*rado*, to shave off. Perhaps from the ease with which it is removed from the matrix. *Fr. Hym. Eur. p. 578. B. & Br. n. 1808. S. Mycol. Scot. n. 766.—Pers. Obs. 2. p. 14.*

**116. P. vaporarius** Fr.—Effused, innate, the white floccose mycelium creeping into the wood. **Pores** large, angular, *pale-white*, crowded into a continuous, firm, persistent stratum.

On dead branches. Very common. Jan.—Dec.

Extremely variable. Name—*vaporarium*, a stove. Perhaps first found in hothouse. *Fr. Hym. Eur. p. 579. Berk. Out. p. 252. C. Hbk. n. 815. S. Mycol. Scot. n. 767.—Pers. Myc. Eur. p. 106.*

**Var. secernibilis** B. & Br.—Shining white, honey-colour when dried, separable.

On fir-leaves under moss. Autumn. Forres. Name—from its being separable. *B. & Br. n. 1022. S. Mycol. Scot. n. 767.*

**117. P. Gordoniensis** B. & Br.—Persistently shining white, effused, superficial, *membranaceous*, very thin but *separable, margin* shortly *fringed*. **Pores** minute, unequal, angular, *dissepiments very thin, fimbriato-toothed*.

An extremely delicate species, and not in the slightest degree innate. The margin remains snow-white, and the pores themselves change colour only very slightly in drying.

On fir poles. Aboyne.

Name—after Marchioness of Huntly. *B. & Br. n. 1023. C. Hbk. n. 813. S. Mycol. Scot. n. 768. Fr. Hym. Eur. p. 579.*

**118. P. Vaillantii** Fr.—White, thin, *the free mycelium resulting in root-like ribs which are somewhat united by a membrane*. **Pores** now and then crowded together, curt, rather large, thin, unequal.

**Polyporus.** It occurs in innumerable more or less imperfect stages of development.  
 Var. Mycelium simply membranaceous.

On dead wood. Rare. April.

Name—after Vaillant. *Fr. Hym. Eur. p. 579. Berk. Out. p. 252. C. Hbk. n. 819. S. Mycol. Scot. n. 769.*—*Sow. t. 326.*

**119. P. farinellus** Fr.—White, widely effused, thin, *mycelium* naked, *flocculoso-pulverulent, not woven.* Pores thin, continuous, unequal, somewhat flexuous, intricate.

Fugacious to the touch. Of the consistence of *Thelaphora sambuci*.

On beech. Aboyne.

Name—*farina*, meal. From its pulverulent nature. *Fr. Hym. Eur. p. 579. B. & Br. n. 1292. S. Mycol. Scot. n. 770.*

**120. P. hymenocystis** B. & Br.—Snow-white, arachnoid beneath and at the margin. Pores large; the scarious dissepiments collapsing, at length pallid.

A very curious species.

On dead wood. Glamis, 1875.

Name—*μῆν*, membrane; *κύστις*, bladder. From the peculiar nature of the dissepiments. *B. & Br. n. 1810 bis. S. Mycol. Scot. n. 775.*

\*\*\*\*\* Pores white, superficial, &c.

**121. P. blepharistoma** B. & Br.—Wholly resupinate, snow-white, mycelium arachnoid, somewhat mealy. Pores small, dissepiments thin, margin ciliato-dentate.

Very thin and delicate; the ciliato-dentate margin of the pores very elegant.

On dead wood. Glamis, 1874. Nov.

Name—*βλέφαρον*, eyelid; *στόμα*, mouth. From the appearance of the pores. *B. & Br. n. 1434. S. Mycol. Scot. n. 771.*

**122. P. reticulatus** Fr.—White, orbicular, thin, *fugacious*, radiating like flax at the floccose margin. Pores distant, *cup-shaped.*

On rotten wood. Glamis, 1874. Forres. April–Sept.

Extremely delicate and beautiful. Name—*rete*, a net. From the network of pores. *Fr. Hym. Eur. p. 580. Icon. t. 190. f. 3. B. & Br. n. 1810. S. Mycol. Scot. n. 772.*—*Pers. Syn. p. 584.*

*Species of uncertain affinity.*

**123. P. bathyporus** Rostk.—“Effused, white, the thin circumference flaxy. Pores rather large, cyathiform, toothed.”



The description does not agree well with the figure. According to the fig. Polyporus. it wanders over beech-branches, and is stratose (the stratum of the second season fuscous-ferruginous), smooth at the margin, fuscous in the middle, livid at the circumference.

On dead oak-branches. Coed Coch, 1873, &c. Very like resupinate form of *Dædalea confragosa*. B. & Br. Name—*βαθύς*, deep; *πόρος*, a pore. From the cup-like pores. Rostk. iv. t. 59. Fr. Hym. Eur. p. 580. B. & Br. n. 1807.

**124. P. hybridus** B. & Br.—White; mycelium thick, forming a dense membrane or creeping branched strings; hymenium breaking up into areæ. Pores long, slender, minute.

The dry-rot of oak-built vessels. On oak, in ships, &c. According to Fries it is a monstrous form (perhaps of *P. destructor*), originating from its close situation. Name—*hybridus*, hybrid. Berk. Out. p. xvii. C. Hbk. n. 820. S. Mycol. Scot. n. 773. Fr. Hym. Eur. p. 581.—Sow. t. 289, t. 387. f. 6.

GENUS XXV.—**Trametes** (*trama*, the generic distinction depending on the trama).

Trametes.

Pores somewhat round, obtuse, entire, often unequal in depth and not forming a heterogeneous stratum, immersed at the base in the flesh of the pileus, and hence the trama is continuous with the flesh of the pileus and similar to it. *Growing on wood, woody or corky, becoming hard, but not stratose, commonly sweet-scented.*

A genus intermediate between the *Dædaleæ* and *Polypori*, the limits of which are not yet duly ascertained. Fr. Hym. Eur. p. 581.

I. APODES (*α*, *πούς*, a foot. Stemless). Pileus dimidiate, sessile.

\* Substance coloured.


\*\* Substance white.

II. RESUPINATI. Pileus resupinate.

I.—APODES. Pileus dimidiate, sessile.

\* Substance coloured.

**1. T. pini** Fr.—Pileus ferruginous-fuscous then blackish, internally tawny-ferruginous, corky-woody, *very hard, pulvinate*,



LXVIII. *Trametes gibbosa*. One-fourth natural size.

Trametes. concentrically sulcate, *rimoso-rugged*, rough. **Pores** large, somewhat round or oblong, yellow-brick-red.

Perennial, but without distinct strata. Odour weak.

On living pine. Rare.

Name—*pinus*, pine. *Fr. Hym. Eur. p. 582. Berk. Out. p. 253. C. Hbk. n. 821. S. Mycol. Scot. n. 776.—Brot. Kl. Bor. t. 380. Rostk. t. 50.*

**\*\* Substance white.**

**2. T. gibbosa** Fr.—Everywhere *whitish*. **Pileus** corky, compact, villous, obsoletely zoned, extended behind, *gibbous*. **Pores** *linear*, straight, equal.

Large, thick, margin obtuse.

On poplar, gate-posts, stumps, &c. Rare.

Zones convex and tuberculated; beautifully velvety; when old cinereous and green from minute *Algae*. *M. f. B.* Name—*gibbus*, gibbous. *Fr. Hym. Eur. p. 583. Berk. Out. p. 253. C. Hbk. n. 824. Fl. Dan. t. 1964. Hussey ii. t. 4.—Pers. Syn. p. 501. B. sinuosus Sow. t. 194 var.*

**3. T. Bulliardi** Fr.—**Pileus** *white becoming fuscous*, at length *zoned*, corky, *becoming plane*, even, *smooth*. **Pores** somewhat round, deep, unequal, pallid then *rufescent*.

It becomes at length fuscous internally. Odour fragrant.

On dead wood. Bathampton. Glamis. Oct.

Berkeley is inclined to regard it as a species of *Dædalea*. Name—after Bulliard. *Fr. Hym. Eur. p. 584. B. & Br. n. 1293. S. Mycol. Scot. p. 412.—Bull. t. 310.*

**4. T. suaveolens** Fr.—**Pileus** white, corky, *soft, pulvinate*, villous, *zoneless*. **Pores** round, rather large, obtuse, *white becoming fuscous*.

Odour fragrant, of anise.

On willow, lime, &c. Uncommon.

Spores white, oval or pip-shaped,  $9 \times 4$  mk. *W.G.S.* Name—*suavis*, sweet. Sweet-scented. *Fr. Hym. Eur. p. 584. Berk. Out. p. 253. C. Hbk. n. 822. S. Mycol. Scot. n. 777. Hussey i. t. 43. Krombh. t. 4. f. 25. Fl. Dan. t. 1849. Bot. Zeit. 1859. t. 11. f. 23. Harz. t. 49.—Linn. Sow. t. 228. Tratt. Austr. f. 4. Sterb. t. 27. D.*

**5. T. odora** Fr.—**Pileus** pallid, corky, elastic, uneven, becoming smooth, *zoneless*. **Pores** *minute*, round, equal, *whitish-ochraceous*.

Odour fragrant.

On willow. Rare.

Very nearly related to *T. suaveolens*. *M. J. B.* Spores white, oval or pip-shaped,  $3 \times 5$  mk. *W. G. S.* Name—odor, scent. *Fr. Hym. Eur. p. 584. Berk. Out. p. 253. C. Hbk. n. 823.—Sommerf. Bolt. t. 162.* Trametes.

**6. T. inodora** Fr.—Shining white. **Pileus** corky, *firm*, becoming smooth, zoneless. **Pores** minute, round, entire, *not changing colour*.

Very like *T. suaveolens*, &c., but smaller, firmer, always white (pores not becoming fuscous); distinguished by its want of odour.

On old mossy beech-stump. Stoke Poges.

Pores colourless, slightly angular, about  $\frac{1}{16}$  in. wide, nearly  $\frac{1}{4}$  in. long, not the least linear. Externally tomentose, white with a very slight tinge of pink at the base. As in *Dædalea latissima*, the texture radiates from a central point, and is of a pure white. *B. & Br.* Name—*inodorus*, scentless. *Fr. Hym. Eur. p. 584. Monogr. ii. p. 273. Icon. t. 191. f. 1. B. & Br. n. 1570.*

## II.—RESUPINATI. Pileus resupinate.

**7. T. mollis** Fr.—Resupinate, determinate, somewhat membranaceous, pallid-wood-colour, at length becoming fuscous, *umber at the margin, which is at length revolute* and pubescent *beneath*. **Pores** large, unequal and torn.

A very distinct species, at length becoming black, adnate behind in the centre. Pores irregular, but not flexuoso-labyrinthine.

On dead wood, especially beech. Common. Jan.—Dec.'

Dissepiments rigid. Name—*mollis*, soft. *Fr. Hym. Eur. p. 585. S. Mycol. Scot. n. 778. Dædalea Sommerf. Polyporus cervinus P. Berk. Out. p. 247. C. Hbk. n. 787.*

**8. T. Terrei** B. & Br.—Resupinate, broad, somewhat orbicular, pulvinate, substance corky and white. **Pores** angular, here and there sinuate, pallid.

About 3 in. across, 1 in. thick in the centre. Substance radiating from a central point, zoneless. Pores about  $\frac{1}{16}$  in. across, angular in the centre, sinuated towards the edge. Inodorous. Habit that of *Dædalea latissima*.

On beech. Stoke Poges.

Name—after Michael Terry. *B. & Br. n. 1571.*

**9. T. serpens** Fr.—White, arid, closely adnate, at first *erumpent, in the form of a tubercle*, orbicular, then confluent, the determinate circumference pubescent. **Pores** somewhat round and angular, unequal, obtuse.

Colour at length becoming pale. A very distinct species analogous with *Radulum orbiculare*.

On privet. Rare. Oct.

Pores rather large,  $\frac{1}{10}$  in. broad. *B. & Br.* Spores oval, white,  $14 \times 7$  mk. *W.G.S.* Name—*serpens*, creeping. *Fr. Hym. Eur. p.* 586. *Icon. t.* 182. *f.* 3. *Polyporus Stephensii B. & Br. n.* 356. *Berk. Out. p.* 252. *C. Hbk. n.* 818. *Saund. & Sm. t.* 45. *f.* 8-12.

## Dædalea.

GENUS XXVI.—*Dædalea* (δαῖδαλος, curiously wrought. From the elaborate hymenium). *Pers. Syn. p.* 449.

Pores firm, when fully grown sinuous, labyrinthine; in all other respects agreeing with *Trametes*. Growing on wood, becoming hard, the floccose substance descending unchanged into the trama. *Fr. Hym. Eur. p.* 586.



\* Corky-woody, dimidiate, sessile.

\*\* Coriaceous, dimidiate, sessile.

\*\*\* Resupinate.

\* Corky-woody, dimidiate, sessile.

LXIX. *Dædalea quercina*. One-fourth natural size.

1. *D. quercina Pers.*—Pale wood-colour. Pileus corky, rugulose, uneven, zoneless, rather smooth, of the same colour internally, at first porous; the pores then take the form of sinuses which are somewhat contorted or lamellose, becoming

at last labyrinthine, edge obtuse.

There are many varieties—effuso-reflexed: *Schæff. t.* 231; wholly resupinate: *Bull. t.* 352. *f. F.* *Pers. Myc. Eur. t.* 18. *f.* 1.

On dead oak. Frequent. June–Nov.

Perennial. Pileus 12.5–15 cent. (5–6 in.) broad. Pores 2 mm. (1 lin.) or more broad. Sometimes resupinate or decurrent and the partitions elongated into tooth-like processes. *M. f. B.* Name—*quercus*, oak. *Pers. Syn. p.* 500. *Fr. Hym. Eur. p.* 586. *Berk. Out. p.* 254. *t.* 19. *f.* 5. *C. Hbk. n.* 825. *S. Mycol. Scot. n.* 779. *Grev. t.* 238. *Krombh. t.* 5. *f.* 1, 2. *Bail. t.* 31.—*Sow. t.* 181. *Bolt. t.* 73. *Bull. t.* 352, 442. *f.* 1. *Vent. t.* 60. *f.* 1, 2 (var.)

2. *D. aurea Fr. Golden.*—Pileus 5 cent. (2 in.) broad, corky-coriaceous, gibbous, velvety, somewhat zoned, unicolorous, internally, as well as the hymenium which is porous then narrowly sinuato-labyrinthine, light yellow.

Pileus triangular, margin swollen.

On dead oak. Hereford.

Imbricated, the veins for the most part straight and radiating. *B. & Br.*

Name—*aurum*, gold. *Fr. Hym. Eur. p. 587. B. & Br. n. 1892.—Batt. t. Dædalea. 35. f. F.*

**3. D. confragosa** Pers.—*Pileus unicolorous, brick-red-fuscous, internally wood-colour, becoming ferruginous, corky, slightly convex, rough, somewhat zoned; sinuses porous then narrowly labyrinthine and torn, cinereous-pruinose then rufous-fuscous.*

On willow, service, &c. Rare.

Name—*confragus*, rough, rugged. *Pers. Syn. p. 501. Fr. Hym. Eur. p. 587. Berk. Out. p. 254. C. Hbk. n. 826.—Bolt. t. 160. Bull. t. 491. f. 1. Sow. t. 193.*

**4. D. cinerea** Fr.—*Pileus 4–7.5 cent. (1½–3 in.) broad, cinereous, corky-woody, thick, somewhat undulated, zoned, tomentose. Pores minute, obtuse, quite entire, some round, some very long, labyrinthiform, flexuous, intricate, white or cinereous.*

Perennial, stratose, allied however to *D. unicolor*.

On trunks. Penzance.

The thick substance separates this from every form of *D. unicolor*, also the inciso-strigose surface of the pileus. *B. & Br. Name—cinereus, ash-colour. Fr. Hym. Eur. p. 588. Syst. Myc. i. p. 336. Icon. t. 182. f. 2. B. & Br. n. 1968.*

**\*\* Coriaceous, dimidiate, sessile.**

**5. D. unicolor** Fr.—*Pileus cinereous, with zones of the same colour, coriaceous, villosa-strigose; sinuses labyrinthiform, flexuous, intricate, acute, at length torn into teeth.*

Commonly imbricated, fuliginous when damp, whitish-grey when dry. Hymenium whitish-cinereous, sometimes fuscous. There are two forms: one thicker, margin whitish, dissepiments firmer, distant, forming more entire sinuses; the other thinner, broader, the sinuses narrower, more torn. *Bull. t. 501. f. 3.* represents a remarkable variety (?) *rufescent* with very minute sinuses.

On stumps, especially birch. Common. July–Oct.

*Pileus 5–10 cent. (2–4 in.) broad. Often green from minute Algæ. M. J. B. Name—unus, one; color, colour. Unicolorous. Fr. Hym. Eur. p. 588. Berk. Out. p. 254. C. Hbk. n. 827. S. Mycol. Scot. n. 780.—Bull. t. 501. f. 3. Bolt. t. 163. Sow. t. 325.*

**\*\*\* Resupinate.**

**6. D. latissima** Fr.—*Pale wood-colour, effused, 5 in.–2 ft. broad, corky-woody, thick, undulated, zoned within; sinuses narrow, distant, somewhat round and very long, flexuous.*

The structure is very peculiar, radiating from the base to the surface, with parallel filaments.

Dædalea. On dead wood. Rare.

Name—*latus*, broad. Very broad. *Fr. Hym. Eur.* p. 589. *Syst. Myc.* i. p. 340. *Berk. Out.* p. 254. *C. Hbk. n.* 828. *S. Mycol. Scot. n.* 781.

**7. D. vermicularis** Pers. — *Flesh-colour-rufescent*, broadly effused, *thin*, becoming even; sinuses short, attenuated at both ends, flexuous, poriform at the margin.

It has much in common with *D. latissima*, but is much thinner, 10–15 cent. (4–6 in.) long, 5 cent. (2 in.) broad. Adhering to the soil by root-like fibres.

On the ground.

Berkeley refers the figure of Sowerby to *D. latissima*. Name—*vermis*, a worm. From the peculiar colour. *Pers. Myc. Eur.* 3. p. 2. *Fr. Hym. Eur.* p. 589. *Sow. t.* 424.

*Species of uncertain affinity.*

**8. D. ferruginea** Schum. — *Pileus light-yellow-ferruginous, white-villous when young and at the margin*, internally flesh-colour, coriaceous, effuso-reflexed, zoned; hymenium porous then narrowly labyrinthiform, tawny.

Imbricato-connate. It seems nearest to *D. unicolor*.

On dead wood. Strachan, Kincardineshire.

Name—*ferrum*, iron. Ferruginous. *Schum.*—*Fr. Hym. Eur.* p. 589. *S. Mycol. Scot. n.* 782. *Fl. Dan. t.* 2029.

Hexagona. GENUS XXVII.—**Hexagona** (*hexagonum*, a six-sided figure; from the form of the pores). *Poll. Pl. nov.* p. 35. *Epicr.* i. p. 496.

Hymenophore descending and forming a trama of the same colour as the pileus and similar to it. *Pores at the very first dilated into hexagonal*, regular, woody-hard *alveoli*, the dividing walls firm, never torn. *Dimidiate, sessile, growing on wood, corky-woody, persistent, but not growing again to form strata.*

A very natural genus, easily recognised. The essential feature consists in the depressions being equally wide in the earliest stage of growth, whereas in the Polypori the pores, which are minute when young, are gradually developed and dilated. *Fr. Hym. Eur.* p. 589.

No British species.

GENUS XXVIII.—**Favolus** (*favus*, honeycomb).

Favolus.

Fr. Elench. p. 44.


Hymenium reticulato-cellular, or alveolate. Alveoli radiating, formed of the densely anastomosing gills, elongated. Spores white (in pairs?). *Dimidiate, somewhat stipitate, fleshy, pliant, annual, growing on wood, differing in entire appearance and structure from the preceding genera*, wherefore they were formerly referred not to Polypori, but to Cantharelli or Merulii. *Fr. Hym. Eur. p. 590.*

No British species.

GENUS XXIX.—**Merulius**. Fr. Syst. Myc. i. p. 326.

Merulius.

Hymenophore formed from a woven mucedinous mycelium, covered with the waxy-soft, continuous hymenium which is reticulated on the surface with obtuse folds, imperfectly porous, at length gyrose and obsoletely toothed. *Growing on wood, soft, mucedinous, at the first at least resupinate. Fr. Hym. Eur. p. 591.*



I. LEPTOSPORI (λεπτός, fine, slight; πόρος, a pore. From the shallow pore-like folds). Hymenium naked or obsoletely pruinose with the white spores.

\* *Pileus effuso-reflexed, circumference determinate.*

\*\* *Resupinato-effused, flaxy-membranaceous, separating, flaxy beneath and at the circumference.*

\*\*\* *Crustoso-adnate, somewhat flaxy at the circumference.*

II. CONIOPHORI (κόνις, dust; φέρω, to bear). Hymenium pulverulent with the ferruginous spores. *All destructive.*

LXX. *Merulius lachrymans*. One-third natural size.

I.—LEPTOSPORI. Hymenium naked, &c.

\* *Pileus effuso-reflexed, circumference determinate.*

1. **M. tremellosus** Schrad.—White, resupinate then free or reflexed, *fleshy-tremellose*, tomentose, margin dentato-radiate. **Folds** porous, various, ruddy.

Sporting in form, becoming pale in hidden places, but easily recognised by its gelatinoso-cartilaginous flesh.

*Merulius*. On decayed trees, birch, &c. Rare. Sept.—Feb.

Sometimes the edge is beautifully tinged with pink. *M. f. B.* Spores cylindrical, curved, hyaline,  $4 \times 1$  mk. *K.* Name—*tremo*, to tremble. From the substance approaching *Tremella*. *Schrad.—Fr. Hym. Eur. p. 591. Berk. Out. p. 255. C. Hbk. n. 829. S. Mycol. Scot. n. 783. Fl. Dan. t. 1553, 776. f. 1. Hussey i. t. 10. Kl. Bor. t. 460.*

**2. *M. aurantiacus*** Klotsch.—**Pileus** 2.5 cent. (1 in.) broad, between yellow and dirty white, here and there cinereous, tough, fleshy-coriaceous, effuso-reflexed, *tomentose*, *zoned*. **Folds** minute, somewhat poriform, dull *orange*.

Zones obsolete, hirsuto-tomentose.

On dead beech. Rare. May.

Allied to *M. lachrymans*. *M. f. B.* Name—*aurum*, gold. Orange-yellow. *Kl. in Berk. Eng. Fl. v. p. 128. Berk. Out. p. 256. C. Hbk. n. 832. S. Mycol. Scot. n. 784. Fr. Hym. Eur. p. 591.*

**3. *M. corium*** Fr.—White, resupinato-effused, 5–7.5 cent. (2–3 in.) and more long, and often equally broad, *soft*, *somewhat of the consistence of paper*, reflexed at the circumference which is at length free, villous beneath; *hymenium* reticulato-porous, *flesh-colour* or pale tan.

Remarkable, variable in form and colour, sometimes short and reflexed, sometimes broadly expanded.

On dead wood. Common. March–Dec.

Spores oval, vivid orange,  $4 \times 5$  mk. *W.G.S.* Name—*corium*, skin, leather. *Fr. Hym. Eur. p. 591. Elench. p. 58. Berk. Out. p. 255. C. Hbk. n. 830. S. Mycol. Scot. n. 785. Grev. t. 147.—Bull. t. 402. Sow. t. 349.*

**\*\* *Resupinato-effused, flaxy-membranaceous, &c.***

**4. *M. læticolor*** B. & Br.—Bright-orange, wholly effused, adnate; margin tomentose, white; *hymenium* even then plicato-rugose; folds distant.

On sawdust and leaves. King's Lynn.

Name—*lætus*, bright; *color*, colour. *B. & Br. n. 1681.*

**5. *M. himantioides*** Fr.—*Lilac*, effused, silky, very soft, *fibrous-silky beneath*, circumference flaxy. **Folds** porous then gyrose, dingy yellow then somewhat olivaceous.

Like *M. lachrymans*, but thinner and not pulverulent.

On fir wood and on club-mosses. Rare. Sept.

Name—*ιμάς*, a strap or thong; *εἶδος*, appearance. From the thong-like



folds. *Fr. Hym. Eur. p. 592. Icon. t. 193. f. 1. Br. in Bath Trans. 1870, Merulius. p. 84. C. Hbk. n. 831. S. Mycol. Scot. n. 786. Pers. Myc. Eur. t. 14. f. 3, 4. B. arboreus Sow. t. 346?*

**6. M. molluscus** Fr.—*White*, effused, thin, soft, membranaceous, *margin flaxy. Folds* poroso-gyrose, *flesh-colour*.

It varies with the pileus shortly reflexed and then the hymenium is not reticulated, but at the first gyroso-dentate.

On fir-branches. Uncommon. Sept.—Feb.

Hymenium dark brown when old. *M. J. B. Name—molluscus, soft. Fr. Hym. Eur. p. 592. Icon. t. 193. f. 2. Berk. Out. p. 255. C. Hbk. n. 833. S. Mycol. Scot. n. 787. Pers. Myc. Eur. 2. t. 14. f. 1, 2.*

\*\*\* *Crustoso-adnate, circumference somewhat flaxy.*

**7. M. porinoides** Fr.—Crustaceo-adnate, thin, the flaxy circumference white. **Folds** poriform, distant, *light yellow-dingy*.

In respect of the hymenium it resembles the preceding rather than the following species.

On dead wood or on ground. Rare.

Name—*πόρος*, a pore; *εἶδος*, form. From the poriform folds. *Fr. Hym. Eur. p. 593. Berk. Out. p. 255. C. Hbk. n. 834. Pers. Myc. Eur. 2. t. 14. f. 7.*

**8. M. rufus** Pers.—*Red-flesh-colour*, crustaceo-adnate, waxy-soft, smooth, *circumference somewhat naked; hymenium equally porous*, of the same colour.

The habit is that of *Polyporus*, but it is distinct in its *waxy-soft* nature. Often immersed in the wood.

On dead oak. Uncommon. Oct.

It has a very polyporoid appearance. *M. J. B. Name—rufus, reddish. Pers. Myc. Eur. 2. t. 16. f. 1, 2. Fr. Hym. Eur. p. 593. Berk. Out. p. 255. C. Hbk. n. 835. S. Mycol. Scot. n. 788.*

**9. M. serpens** Tode.—*Pallid becoming red, crustaceo-adnate*, thin, becoming smooth, the flaxy circumference white. **Folds** at first in the form of wrinkles, then porous, angular, entire.

On dead fir. Rare. Oct.

Not separable like *M. corium*. *M. J. B. Name—serpo, to creep. Creeping. Tode.—Fr. Hym. Eur. p. 593. Icon. t. 193. f. 3. Berk. Out. p. 255. C. Hbk. n. 836. S. Mycol. Scot. n. 789.*

**10. M. pallens** Berk.—Pale-reddish, adnate, fleshy, somewhat gelatinous, thin, inseparable; margin indeterminate. **Folds** poriform.

Pores minute.

Merulius.

On fir and oak. Rare. Oct.—Dec.

Name—*palleo*, to be pale. *Berk. Out.* p. 256. *C. Hbk. n.* 837. *S. Mycol. Scot. n.* 790. *Fr. Hym. Eur.* p. 593.

**11. M. Carmichaelianus** Berk.—White, extremely thin, forming effused, entirely resupinate, irregular, interrupted, confluent patches. **Folds** forming regular, angular reticulations; dull brown when dry.

Forming a mere pellicle, with minute, often hexagonal, extremely shallow pores, like the cells of a honeycomb but quite superficial.

On bark. Appin.

Name—after Carmichael. *Berk. Out.* p. 256. *C. Hbk. n.* 838. *S. Mycol. Scot. n.* 791. *Fr. Hym. Eur.* p. 593.—*Grev. t.* 224.

## II.—CONIOPHORI. Hymenium pulverulent, &c.

**12. M. lachrymans** Fr.—*Light-yellow-ferruginous, spongy-fleshy, slightly moist*, spider-web-velvety beneath, margin swollen, tomentose, white. **Folds** large, porous and gyroso-toothed.

Exuding watery drops when in fullest vigour. Commonly in the form of a placenta, but also effuso-reflexed, nay, from a medial placenta stalked tubercles producing small pilei arise. Hymenium exceedingly changeable, sometimes like that of *Hydnum*.

On dressed wood, &amp;c. Very common.

The dry-rot of houses. Often covering a large surface of the wood. Spores oval, rich orange brown,  $9 \times 6$  mk. *W.G.S.*; ellipsoid, somewhat yellow,  $10-12 \times 5-6$  mk. *K.* Name—*lachryma*, a tear. From the watery drops. *Fr. Hym. Eur.* p. 594. *Sv. atl. Sv. t.* 70. *Berk. Out.* p. 256. *C. Hbk. n.* 839. *S. Mycol. Scot. n.* 792. *Harz. t.* 77. *Hussey i. t.* 3. *Fl. Dan. t.* 2026. *Krombh. t.* 46. *f.* 1-2. *Sow. t.* 113.—*Wulf in Jacqu. Coll. t.* 8. *f.* 2.

\* **M. pulverulentus** Fr.—Very broad, *membranaceous, becoming even, zoned, arid*, spider-web-velvety beneath, gradually decaying from the centre towards the circumference. **Folds** marginal, somewhat reticulated, tawny.

Of many colours. Singular from the peculiar way in which the hymenium recedes from the centre. Strachan, Kincardineshire. Name—*pulvis*, dust. Pulverulent. *Fr. Hym. Eur.* p. 594. *S. Mycol. Scot. n.* 792.

Porothe-  
lium.

GENUS XXX.—**Porothelium** (πόρος, a pore; θηλή, a teat. From the porous spines). *Fr. Obs.* 2. p. 272.


Whole fungus composed of mycelium, expanded resupinately, somewhat membranaceous, producing *papillæ which are at the*

*first distinct* (and closed), soon opened in the form of pores, at length elongated and tubular. *A genus very distinct from the preceding ones, in many respects analogous with* *Fistulina*. Fr. Hym. Eur. p. 595. Porothelium.

1. **P. Friesii** Mont.—White, somewhat tan-colour, effused, confluent, *flocculoso-membranaceous*, circumference simple. **Warts** containing the pores *immersed*, yellowish, at length opening in pitcher-shape.

On dead pine. Rare. Nov.—Feb.

Name—after Elias Fries. *Montagn. Ann. Sc. Nat.* 1836, *Jun. n.* 39. Fr. *Hym. Eur. p.* 595. *Berk. Out. p.* 257. *C. Hbk. n.* 840. *S. Mycol. Scot. n.* 793.



2. **P. Stevensoni** B. & Br.—Substance rather thick, gelatinous, margin rather coarsely hispid, at length more or less denuded. **Warts** of the hymenium distinct, interstices very smooth, bearing at the apex a limpid, diaphanous, yellow globule.

LXXI. *Porothelium Stevensoni*. Natural size. Section five times natural size.

On old pine rail. Glamis, 1877–1886. Spring, autumn.

Almost white when fresh, becoming yellowish when old. The limpid globule is at first pale. When full grown the papilla is from four to five times as long as the diameter of the globule. The mycelium imparts a sweet scent to the wood. Found in greatest perfection on the under side of very old pine rail lying on the ground. Name—after Rev. John Stevenson. *B. & Br. n.* 1683. *S. Mycol. Scot. n.* 794.

3. **P. Keithii** B. & Br.—Narrowly adnate, slightly umber-colour; circumference very thin, at first somewhat gelatinous. **Warts** short, at length collapsed, gelatinous in the centre.

On dead fir. Sanquhar. Dunphail. Glamis. April.

Name—after Rev. Dr Keith. *B. & Br. n.* 1684. *S. Mycol. Scot. n.* 795.

4. **P. confusum** B. & Br.—Narrowly adnate, pallid, margin very thin, arachnoid, substance at first floccoso-pulverulent. **Warts** very small.

On fir sticks. Glentanner. Leighwood. Glamis. Pitoulish, Inverness-shire. June–Sept.



Name—*confundo*, to confuse. From having been confounded with a species in another genus. *B. & Br. n.* 1685. *S. Mycol. Scot. n.* 796.

Solenia.

GENUS XXXI.—**Solenia** (σωλήν, a pipe). Hoffm. Deutschl. Fl.

Receptacle none. Tubes membranaceous, somewhat cylindrical, turbinate, distinct and free from one another, definitely facing the ground, *mouth narrowed*, in which respect they differ from the *Cyphellæ*. Fr. *Hym. Eur.* p. 595.

Most nearly allied to *Cyphellæ*.



LXXII. *Solenia ochracea*. Natural size. Section twenty times natural size.

1. **S. ochracea** Hoffm.—*Ochraceous*, white internally, scattered, *clavato-cylindrical*, tomentose.

On dead wood. Uncommon. Aug.—Dec.

Spores subglobose,  $6 \times 8$  mk. *W.G.S.* Name—colour of ochre. *Hoffm. D. Fl.* ii. t. 8. f. 2. Fr. *Hym. Eur.* p. 596. Br. *Bath Trans.* 1870, p. 90. C. *Hbk.* n. 957. *S. Mycol. Scot.* n. 797.

## ORDER III.—HYDNEI.

Hymenium inferior or amphigenous, from the first figurate, and definitely but variously protuberant in the form of persistent spines, teeth, tubercles, crests, papillæ. Sporophore sometimes single-spored. *The greater part are resupinate and effused, and hence they are inferior to the Agaricini, which are commonly circular on a central stem, and to the Polyporei, which are most frequently dimidiate.* Fr. Hym. Eur. p. 598.

GENUS XXXII.—**Hydnum** (ὕδνον, the Greek name for a kind of edible fungus). Linn. Gen. Plant. Hydnum.

Hymenium inferior, aculeate; spines awl-shaped, distinct at the base. The central genus of the order, various, analogous with the *Polyperi*. Fr. Hym. Eur. p. 598.

I. MESOPUS (μεσός, middle; πούς, a foot). Entire, simple, with a central stem. *All growing on the ground and chiefly in pine woods.*

† Carnosa. Pileus fleshy, somewhat fragile. *Most of them edible.*

\* Spines changing colour, pileus scaly or tomentose.

\*\* Spines changing colour, pileus even, smooth.

\*\*\* Spines unchangeable, unicolorous, whitish.

†† Lignosa. Pileus corky or coriaceous, tough.

\* Spines changing colour, and, as well as the spores, somewhat fuscous.

\*\* Spines changing colour, and, as well as the spores, ferruginous.


\*\*\* Spines unchangeable and spores whitish.

II. PLEUROPUS (πλευρόν, the side; πούς, a foot). Somewhat dimidiate, stem lateral.

III. MERISMA (μερίζω, to divide). Very much branched or tuberculiform and immarginate.

\* Very much branched.

\*\* Simple, tuberculiform, immarginate.



LXXIII. *Hydnum repandum*.  
One-third natural size. Section  
twice natural size.

Hydnum.

IV. APUS (ἄ, πούς, a foot. Stemless). Pileus sessile, dimidiate, marginate, often effuso-reflexed.

\* *Pileus fleshy, soft.*

\*\* *Cæspitose, pilei imbricated, commonly pliant, tough.* Very distinguished.

\*\*\* *Simple, pileus coriaceous or corky.*

\*\*\*\* *Pileus membranaceous.*

V. RESUPINATI. Pileus none. *Absolutely resupinate, with straight or oblique spines according to their situation.*

\* *Spines fuscous, ferruginous.*

\*\* *Spines yellowish, green.*

\*\*\* *Spines flesh-colour, lilac, rufescent.*

\*\*\*\* *Spines white or light-yellowish when old.*

I.—MESOPUS. Entire, simple, with a central stem.

† Pileus fleshy, somewhat fragile.

\* *Spines changing colour, pileus scaly or tomentose.*

1. **H. imbricatum** Linn.—**Pileus** about 10 cent. (4 in.) broad, *umber*, zoneless, fleshy, rather plane, somewhat umbilicate, *floccose*, *tessulato-scaly*; flesh dingy whitish. **Stem** curt, 2.5–7.5 cent. (1–3 in.) long, 2.5–5 cent. (1–2 in.) thick, even. **Spines** 8–12 mm. (4–6 lin.) long, decurrent, cinereous-white.

There are two forms: one with the pileus plane and with thick persistent scales, another with the pileus somewhat infundibuliform, and with thinner, at length separating scales.

In pine woods. Rare. Sept.–Nov.

Often abundant where it occurs. Edible but bitter. Spores spherical, subhyaline, 3 mk. *K.*; 6 × 5 mk. *W.G.S.* Name—*imbrex*, a tile. Imbricated. *Linn. Suec. n.* 1257. *Fr. Hym. Eur. p.* 598. *Sv. ätl. Sv. t.* 33. *Berk. Out. p.* 257. *C. Hbk. n.* 842. *S. Mycol. Scot. n.* 798. *Schæff. t.* 140. *Fl. Dan. t.* 176, 1500. *Sow. t.* 73. *Grev. t.* 71. *Barla t.* 38. *f.* 1–4. *Harz. t.* 36. *Kl. Bor. t.* 462. *Bail. t.* 29.—*Nees Syst. f.* 240.

2. **H. squamosum** Schæff.—**Pileus** rufous-fuscous, fleshy, irregular, depressed, *smooth*, breaking up into *irregular scales*; flesh whitish. **Stem** curt, attenuated downwards, white. **Spines** grey-fuscous, whitish at the apex.

Smaller than *H. imbricatum*, rigid. Pileus even when young.

On the ground. Fungus Show, South Kensington, 1873. Oct.

Name—*squama*, a scale. *Schæff. t.* 273. *Fr. Hym. Eur. p.* 598. *B. & Br. n.* 1435.

3. **H. scabrosum** Fr.—**Pileus** about 10 cent. (4 in.) broad, *umber-ferruginous*, compactly fleshy, *at first turbinate*, then plane above, very convex beneath, at first tomentose, then rough with flocci which are fasciculate in the form of minute crowded

squamules, slightly repand at the margin; flesh very thick, white, descending into the stem. **Stem** very curt, 2.5 cent. (1 in.) long, and equally thick, round or compressed, dotted with the rudiments of spines decurrent upon it, cinereous, attenuated downwards, rounded and blackish at the base. **Spines** 8 mm. (4 lin.) long, equal, awl-shaped, fuscous-ferruginous, whitish at the apex, at first sight grey-fuscous.

Very fleshy, compact, obese.

In pine wood. Forres, 1879. Sept.

Name—*scabrosus*, rough. *Fr. Hym. Eur.* p. 599. *Monogr.* ii. p. 273. *S. Mycol. Scot. Supp. Scot. Nat.* 1881, p. 37. *Schæff. t.* 271.

**\*\* Spines changing colour, pileus even, smooth.**

**4. H. lævigatum** Sw.—**Pileus** 10–15 cent. (4–6 in.) broad, *umber*, fleshy, *compact*, firm, regular, plane, *even, very smooth*, margin circinate (not repand); flesh whitish, compact, but by no means fibrous, soft when fresh, pliant when dry. **Stem** short, thick, even, pallid fuscous. **Spines** thin, pallid fuscous.

Its size is that of *H. imbricatum*, but it occurs twice as large, with the pileus minutely rimuloso-rivulose, by no means scaly. The stem varies curt and unequal or longer and equal. Quite distinct from *H. fragile*.

In pine woods. Rannoch. Aug.–Oct.

Spores globose, verrucose, pale lemon-yellow, 7 mk. *Q.* Name—*lævis*, even. *Swartz. Vet. Akad. Handl.* 1810, p. 243. *Fr. Hym. Eur.* p. 599. *Monogr.* ii. p. 275. *Sv. atl. Sv. t.* 81. *S. Mycol. Scot. n.* 799.

**5. H. fragile** Fr.—**Pileus** at first pallid, soon *cinereous* or brick-rufescent, somewhat zoned towards the margin, fleshy, fragile, unequal, surface *at first pubescent, then becoming smooth*, commonly even but here and there minutely squamulose and pitted (*slightly wrinkled*), margin undulated and lobed; flesh grey, somewhat zoned. **Stem** stout, unequal, smooth, cinereous. **Spines** scarcely decurrent, very much elongated, slender, fragile, whitish then grey.

Very changeable. Stem sometimes short and thick, sometimes elongated. Specimens occur with the stem 7.5 cent. (3 in.) long, 10 cent. (4 in.) thick, and the pileus more than 30 cent. (12 in.) broad. Quite distinct from *H. lævigatum* in its soft substance, in the pileus being at first pubescent, &c.

In pine woods and among heath. Rare. Sept.–Nov.

Name—*fragilis*, fragile. *Fr. Hym. Eur.* p. 599. *Monogr.* ii. p. 275. *Sv. atl. Sv. t.* 89. *B. & Br. n.* 1024, 1366. *C. Hbk. n.* 843. *S. Mycol. Scot. n.* 800. *Berg. Pyr. t.* 16. *Paul. t.* 34.

Hydnum.

\*\*\* *Spines unchangeable, unicolorous, whitish.*

**6. H. repandum** Linn.—**Pileus** 5–15 cent. (2–6 in.) broad, *pallid*, &c., fleshy, fragile, *somewhat repand*, rather smooth. **Stem** 5–12.5 cent. (2–5 in.) long, 1–4 cent. ( $\frac{1}{2}$ – $1\frac{1}{2}$  in.) thick, irregularly shaped, *pallid*. **Spines** 8 mm. (4 lin.) long, unequal, of the same colour.

The colour varies whitish, yellow, somewhat flesh-colour, unchangeable. Pileus sometimes floccoso-pruinose.

In woods. Common. Aug.–Nov.

Margin of pileus more or less arched. *M. J. B.* Spores white or with a suggestion of yellow, round with an obtuse apiculus at one end, 5 mk. *W. G. S.* Edible. One of the most delicious. Requires about four hours' slow cooking. Name—*repandus*, repand. *Linn. Suec. n. 1258. Fr. Hym. Eur. p. 601. Sv. atl. Sv. t. 15. Berk. Out. p. 258. t. 17. f. 2. C. Hbk. n. 844. S. Mycol. Scot. n. 801. Schæff. t. 318. Bull. t. 172. Sow. t. 176. Fl. Dan. t. 310. Vittad. t. 25. f. 2. Krombh. t. 50. f. 1–9. Vent. t. 27. f. 24. f. 4–6. Barla t. 32. f. 1–9. Grev. t. 44. Hussey i. t. 16. Harz. t. 23, &c.*

\* **H. rufescens** Pers. Rufescent.—**Pileus** 5–7.5 cent. (2–3 in.) broad, *thin*, somewhat regular, *pubescent*. **Stem** 2.5–7.5 cent. (1–3 in.) long, commonly thin, somewhat equal. **Spines** regular.

In woods. Frequent. Aug.–Nov. *Pers. Syn. p. 555. Berk. Out., C. Hbk., S. Mycol. Scot.* as quoted above. *Bolt. t. 89.*

†† Lignosa. Pileus corky or coriaceous, tough.

\* *Spines changing colour, and, as well as the spores, somewhat fuscous.*

**7. H. compactum** Pers.—**Pileus** 2.5–15 cent. (1–6 in.) broad, olivaceous-cinereous or fuscous, corky, compact, undulated and tuberculose in becoming confluent, *zoneless*, commonly wrapped round with whitish down, *internally variegated azure-blue*. **Stem** very short, 2.5 cent. (1 in.) and more long, irregularly shaped, tawny inclining to fuscous. **Spines** 2–4 mm. (1–2 lin.) long, fuscous, pallid at the apex.

Irregularly shaped, somewhat sessile, often forming a shapeless crust.

In fir woods and on heaths. Rare. Sept.–Nov.

Name—*compactus*, compact. *Pers. Syn. p. 556, partly. Fr. Hym. Eur. p. 603. Berk. Out. p. 258. B. & Br. n. 1367. C. Hbk. n. 845. S. Mycol. Scot. n. 802. Krombh. t. 50. f. 12. Strauss Sturm. 33. t. 6.—Schæff. t. 146 excluding f. 4, 7.*

**8. H. aurantiacum** Alb. & Schw.—**Pileus** 2.5–12.5 cent. (1–5 in.) broad, *orange*, corky, compact, turbinato-dilated, *with small*



*elevations, zoneless*, often covered over with whitish down, *zoned* Hydnum. *internally*. **Stem** short, 1–5 cent. ( $\frac{1}{2}$ –2 in.) long, 1–2.5 cent. ( $\frac{1}{2}$ –1 in.) thick, orange. **Spines** 2–4 mm. (1–2 lin.) long, whitish becoming fuscous.

Inodorous, firm.

In pine woods. Grantown. Sept.–Nov.

Name—*aurum*, gold. Orange-yellow. *A. & S. p.* 265. *Fr. Hym. Eur. p.* 603. *B. & Br. n.* 1368. *S. Mycol. Scot. n.* 803. *Fl. Dan. t.* 1439. *Batsch f.* 222 (bad).

**9. H. ferrugineum** Fr.—**Pileus** 2.5–10 cent. (1–4 in.) broad, obconic or when more fully grown flattened, even depressed, spongy-corky, soft, *at first* covered over with whitish down *which exudes blood-red drops* in the depressions, *then* more even and *internally ferruginous*. **Stem** firm, 4–7.5 cent. (1½–3 in.) long, unequal, fuscous-ferruginous. **Spines** awl-shaped, equal, fuscous-ferruginous.

Gregarious but not confluent. Spongy-soft when young, corky and juiceless when full grown.

In fir wood. Rothiemurchus. Aug.–Oct.

Name—*ferrum*, iron. Ferruginous. *Fr. Hym. Eur. p.* 603. *Icon. t.* 4. *B. & Br. n.* 1369. *S. Mycol. Scot. n.* 804. *Krombh. t.* 50. *f.* 10, 11. *Bull. t.* 409 (var.)

**\*\* Spines changing colour, and, as well as the spores, ferruginous.**

**10. H. scrobiculatum** Fr.—Ferruginous. **Pileus** 2.5–5 cent. (1–2 in.) and more broad, corky-coriaceous, *clavate then plano-infundibuliform*, pubescent, *disc slightly pitted, scaly, zoned within*. **Stem** very short, 12 mm. ( $\frac{1}{2}$  in.) or a little more long, 6–8 mm. (3–4 lin.) thick, equal, smooth, often rooted. **Spines** very short, slightly decurrent, thin, fragile, of the same colour as the pileus, and at length becoming fuscous.

Gregarious, often growing into each other. Pileus becoming pale when dry.

In fir woods. Minstead, Lyndhurst. Rothiemurchus, &c. Aug.–Nov.

Name—*scrobis*, a trench. From the slightly pitted disc. *Fr. Hym. Eur. p.* 604. *Icon. t.* 5. *f.* 1. *B. & Br. n.* 1294. *S. Mycol. Scot. n.* 805. *Mich. Gen. t.* 72. *f.* 7.—*Bull. t.* 156.

**11. H. zonatum** Batsch.—Ferruginous. **Pileus** 2.5–5 cent. (1–2 in.) broad, *equally coriaceous*, thin, expanded, somewhat infundibuliform, *zoned, becoming smooth, radiato-rugose*, the paler

Hydnum. margin sterile beneath. **Stem** 12-18 mm. ( $\frac{1}{2}$ - $\frac{3}{4}$  in.) long, 4-6 mm. (2-3 lin.) thick, slender, somewhat equal, floccose, base tuberous. **Spines** 2-3 mm. (1-1  $\frac{1}{2}$  lin.) long, slender, pallid then ferruginous.

Very like *H. scrobiculatum*.

In fir woods. Rare. Aug.-Oct.

Name—*zona*, a zone. Zoned. *Batsch f.* 224. *Fr. Hym. Eur. p.* 605. *Berk. Out. p.* 258. *C. Hbk. n.* 846. *S. Mycol. Scot. n.* 806. *Nees Syst. f.* 242.

\*\*\* *Spines unchangeable and spores whitish.*

**12. H. nigrum** Fr.—**Pileus** azure-blue-black, *zoneless*, but with a white margin when in fullest vigour, *corky-rigid*, club-shaped when young, then turbinate, at length flattened, plano-depressed, tubercular, tomentose, black internally. **Stem** 2.5 cent. (1 in.) long, stout, often rooting, unequal, black, tomentose at the base, internally of the same colour. **Spines** awl-shaped, thin, rather short, white.

Cæspitose, rarely scattered, growing into each other in various forms. *Inodorous*. It differs from all others in being internally black.

In pine woods. Street, Somerset.

Sometimes zoned, as in the fig. of Fries. Spores round, papillose, 4 mk. *W.G.S.* Name—*niger*, black. *Fr. Hym. Eur. p.* 605. *Icon. t.* 5. *f.* 2. *B. & Br. n.* 1296. *Seem. Journ.* 1868, *p.* 334. *Bath Trans.* 1870, *p.* 86. *C. Hbk. n.* 847. *B. pileus* olivaceous-cinereous. *Batsch f.* 223.

**13. H. graveolens** Delast.—**Pileus** 2.5 cent. (1 in.) and more broad, black-fuscos, internally fuscous, margin whitish, coriaceous, thin, *soft*, zoneless, rugose, smooth; flesh of the same colour but paler. **Stem** 2.5 cent. (1 in.) or a little more long, scarcely 2 mm. (1 lin.) thick, tough, very slender, equal, smooth, even polished, fuscous-black. **Spines** curt, decurrent, *grey*.

The pileus when dry becomes cinereous. It differs remarkably from neighbouring species in its softer, thin substance, and in its *odour* which is almost that of Melilot. Commonly densely gregarious or cæspitose, but not growing into each other.

In fir woods. Rare. Sept.-Nov.

Retaining its scent for years. Name—*graveolens* (*gravis-oleo*), strong-smelling. *Delast.—Fr. Hym. Eur. p.* 605. *Icon. t.* 6. *f.* 1. *Berk. Out. p.* 258. *C. Hbk. n.* 848. *S. Mycol. Scot. n.* 807.

**14. H. melaleucum** Fr.—**Pileus** black, coriaceous, thin, rigid, irregular, striate, *with little elevations at the disc*, margin white. **Stem** slender, *smooth, black*. **Spines** white.

*Inodorous*. Pileus plane.

In fir woods. Ascot. Forres. Sept.

Hydnum.

Name—μέλας, black; λευκός, white. From the black pileus and white spines. *Fr. Hym. Eur. p. 606. B. & Br. n. 1295. S. Mycol. Scot. Supp. Scot. Nat. 1881, p. 38.—Schæff. t. 272.*

**15. H. cyathiforme** Schæff.—**Pileus** *pale cinereous*, margin white, coriaceous, *thin*, plano-infundibuliform, zoned, disc somewhat tomentose. **Stem** slender, smooth, *pale cinereous*. **Spines** white.

Small, commonly growing into each other.

On fir wood. Ascot. Rothiemurchus. Sept.–Oct.

Spores globular, rough or papillated, 3 mk. *W.G.S.* Name—*cyathus*, a cup; *forma*, form. *Schæff. t. 139. Fr. Hym. Eur. p. 606. S. Mycol. Scot. n. 808. Fl. Dan. t. 1020. f. 2. H. tomentosum B. & Br. n. 1025. C. Hbk. n. 849. Harz. t. 3. a. Seem. Journ. 1868, t. 76.*

II.—PLEUROPUS. Somewhat dimidiate, stem lateral.

**16. H. auriscalpium** Linn.—**Pileus** 12–18 mm. ( $\frac{1}{2}$ – $\frac{3}{4}$  in.) broad, date-brown then blackish, *dimidiate*, coriaceous, *reniform*, *hairy*. **Stem** slender, 5–7.5 cent. (2–3 in.) long, 1–3 mm. ( $\frac{1}{2}$ –1  $\frac{1}{2}$  lin.) thick, vertical, rooted, hairy, of the same colour as the pileus. **Spines** 4–6 mm. (2–3 lin.) long, tough, date-brown.

The pileus sometimes appears entire with lobes all round.

On fir-cones and among fir-leaves. Frequent. Aug.–Nov.

Name—*auriscalpium*, an ear-pick. From its shape. *Linn. Suec. n. 1260. Fr. Hym. Eur. p. 607. Berk. Out. p. 258. C. Hbk. n. 850. S. Mycol. Scot. n. 809. Schæff. t. 143. Bull. t. 481. f. 3. Fl. Dan. t. 1020. f. 1. Curt. Lond. t. 190. Bolt. t. 90. Grev. t. 196. Sow. t. 267. Krombh. t. 50. f. 15-17.*

III.—MERISMA. Very much branched or tuberculiform, &c.

\* *Very much branched.*

**17. H. coralloides** Scop.—Very much branched, shining white, at length yellowish, *wholly broken up into* attenuated intricate branches; primary branches 12 mm. ( $\frac{1}{2}$  in.) and more thick, ultimate ones 2 mm. (1 lin.) and less. **Spines** 6–8 mm. (3–4 lin.) long, *unilateral*, awl-shaped, entire.

On decayed fir, beech, ash, &c. Rare.

Young plant resembling a cauliflower. *Pers.* Edible. Name—κοράλλιον, coral; εἶδος, appearance. Coral-like. *Scop. Carn. 2. p. 472. Fr. Hym. Eur. p. 667. Sv. atl. Sv. t. 34. Berk. Out. p. 259. C. Hbk. n. 851. Schæff. t. 142. Sow. t. 252. Krombh. t. 51. f. 4-7.—Bull. t. 390.*

Hydnum.

\*\* *Simple, tuberculiform, immarginate.*

18. **H. erinaceum** Bull.—**Pileus** 20 cent. (8 in.) and more broad, often much less, *white, becoming yellow*, fleshy, elastic-tough, pendulous, tubercular, immarginate, *torn into fibrils above*. **Spines** very long, 4–6 cent. ( $1\frac{1}{2}$ – $2\frac{1}{2}$  in.), straight, equal, pendulous.

Striking, changeable in form, in the form of a heart when most perfect, often with the rudiment of a lateral stem. Internally as if latticed in a branched manner.

On trunks of oak, beech, &c. Rare.

Spores white, plain,  $5 \times 6$  mk. *W.G.S.* Name—*erinaceus*, a hedgehog. From its appearance. *Bull. t.* 34. *Fr. Hym. Eur. p.* 608. *Berk. Out. p.* 259. *C. Hbk. n.* 852. *Vittad. t.* 26. *Krombh. t.* 51. *f.* 1–3.

19. **H. caput-Medusæ** Bull.—*White, then fuliginous-cinereous*, fleshy, tuberculiform, somewhat stipitate. **Spines** on upper surface distorted, the lower ones long, straight.

Large. The trunk is dilated into a pileus, which is covered over with slender spines on all sides.

On trunks. Rare.

Name—Medusa's head. *Bull. t.* 412. *Fr. Hym. Eur. p.* 608. *Berk. Out. p.* 259. *C. Hbk. n.* 853.

#### IV.—APUS. Pileus sessile, dimidiate, &c.

\* *Pileus fleshy, soft.*

20. **H. cirrhatum** Pers.—**Pileus** pallid, fleshy, expanded, *fibrillose-curved above* with scattered decumbent abortive spines, *the fimbriate margin incurved*. **Spines** very long, 10–16 mm. (5–8 lin.), slightly tough, equal.

Simple or imbricated. Varying white, light yellowish, rufescent.

On a beech-tree. Epping Forest.

Name—*cirrus*, a curl, lock of hair. *Pileus curled.* *Pers. Syn. p.* 558. *Fr. Hym. Eur. p.* 609. *Sv. atl. Sv. t.* 71. *f.* 1. *B. & Br. n.* 1370. *Fl. Dan. t.* 1789. *f.* 2.

21. **H. diversidens** Fr.—**Pileus** 5 cent. (2 in.) broad, white fleshy, thick, *substipitate*, very irregularly shaped, densely beset *above with erect, variously shaped, incised teeth*, clothed at the entire margin with club-shaped spines, and beneath with entire awl-shaped regular spines.

Colour almost that of *repandum*. Singular from having spines of three kinds. Allied to *H. cirrhatum*, but not *cæspitose*, more irregularly shaped, here and there lobed.

On beech and hornbeam. Guildford. Epping Forest. Oct. Hydnum.

Name—*diversus*; *dens*, a tooth. With teeth and spines of different shapes. *Fr. Hym. Eur.* p. 609. *Sv. atl. Sv. t.* 71. f. 2. *Grevillea*, vol. xiii. p. 49. *Krombh. t.* 51. f. 8-12.

**\*\* *Cæspitose, pilei imbricated, &c.***

No British species.

**\*\*\* *Simple, pileus coriaceous  
or corky.***

**22. *H. ochraceum* Pers.—*Pilei***  
2.5-7.5 cent. (1-3 in.) broad, *ochra-*  
*ceous*, effuso-reflexed, coriaceous,  
thin, *zoned*. **Spines** *very small*,  
ochraceous-flesh-colour.

Very small, sometimes wholly resupinate.

On dead branches. Common.  
Nov.

Easily removed from the matrix. *M. F. B.*  
Name—*ochraceus*, ochrey-yellow. *Pers.*  
*Syn.* p. 559. *Fr. Hym. Eur.* p. 612. *Syst.*  
*Myc.* i. p. 414. *Berk. Out.* p. 259. *C. Hbk.*  
n. 855. *S. Mycol. Scot.* n. 810.—*Sow. t.* 15.



LXXIV. *Hydnum ochraceum*.  
Natural size. Section twice  
natural size.

**\*\*\*\* *Pileus membranaceous.***

No British species.

V.—RESUPINATI. *Pileus* none, &c.

**\* *Spines fuscous, ferruginous.***

**23. *H. squalinum* Fr.—*Subiculum*** pale wood-colour, *coria-*  
*ceous*, *firm*, adnate. **Spines** crowded, 4-6 mm. (2-3 lin.) and more  
long, *stout*, *compressed*, entire, at length becoming fuscous.

Rather large, firmer than the rest, villous becoming smooth, variable in  
stature and colour, abnormal, here and there with the connate teeth of *Irpex*,  
and at the same time allied to *Radulum*.

On trunks, especially beech.

Admitted on the authority of Ray and Bolton. Name—*squalus*, a shark.  
From the strong spines like shark's teeth. *Fr. Hym. Eur.* p. 612. *Syst.*  
*Myc.* i. p. 420. *Berk. Out.* p. 259. *C. Hbk.* n. 856. *Ray Syn. t.* 1. f. 5.—  
*Bolt. t.* 74.

Hydnum.

**24. H. membranaceum** Bull.—**Subiculum** *tawny-ferruginous*, effused, *waxy-membranaceous*, agglutinated, *smooth*. **Spines** awl-shaped, crowded, equal, acute, of the same colour as the subiculum.

It varies yellowish and thinner.

On fallen sticks. Frequent.

Name—*membrana*, a membrane. *Bull. t.* 481. *f.* 1. *Fr. Hym. Eur. p.* 613. *Berk. Out. p.* 260. *C. Hbk. n.* 857. *Sow. t.* 327.

**25. H. Weinmanni** Fr.—**Subiculum** *fawn-grey*, effused, *waxy-membranaceous*, *agglutinated*, *smooth*. **Spines** minute, acute, somewhat distant, equal.

Becoming cinereous when dry.

On fallen branches. Bristol.

Name—after Weinmann. *Fr. Hym. Eur. p.* 613. *Berk. Out. p.* 260. *C. Hbk. n.* 858. *Pers. Myc. Eur. t.* 22. *f.* 2?

**26. H. crinale** Fr.—**Subiculum** *umber*, effused, *villos-interwoven*, thin. **Spines** long, crowded, equal, *very slender*, *umber*.

Unicolorous. The spines are almost like hairs.

On dead wood. Rare. Essex.

Remarkable for the long dark rufous-brown spines which shine as if varnished. *M. J. B.* Name—*crinis*, hair. From the hair-like spines. *Fr. Hym. Eur. p.* 613. *H. fuscum Berk. Out. p.* 260. *C. Hbk. n.* 859.—*Pers. Myc. Eur. t.* 17. *f.* 3.

**27. H. ferruginosum** Fr.—**Subiculum** *tawny-ferruginous*, effused, 5–7.5 cent. (2–3 in.) long, tomentose. **Spines** crowded, conico-awl-shaped, acute, *tawny-ferruginous*.

The spines are straight and equal or oblique and somewhat compressed according to situation.

On decayed wood. Uncommon.

Separable from the matrix, almost mucedinous when fully developed. *M. J. B.* Name—*ferrum*, iron. Ferruginous. *Fr. Hym. Eur. p.* 613. *Syst. Myc. i. p.* 416. *Berk. Out. p.* 260. *C. Hbk. n.* 860. *S. Mycol. Scot. n.* 811.—*Schrad. Spic. t.* 4. *f.* 2. *Nees Syst. f.* 248.

**28. H. variegolor** Fr.—**Subiculum** *white*, effused, adnate, *forming a furfuraceous crust*. **Spines** crowded, conical, short, unequal, minute, somewhat fuscous.

Like *H. farinaceum*. Spines commonly adpressed.

On dead stumps. Rare.

Name—*varius, color*. From the colours. *Fr. Hym. Eur. p. 613. Berk. Hydnum. Out. p. 260. C. Hbk. n. 861.*

**\*\* Spines yellowish, green.**

**29. H. aureum** Fr.—**Subiculum** *golden*, determinate, somewhat cartilaginous, continuous, smooth, radiate in the form of teeth at the circumference. **Spines** awl-shaped, equal, of the same colour as the subiculum.

Distinguished, analogous with *Phlebia radiata*, remarkable moreover for the mycelium permeating the wood and giving rise to a distinctly defined yellow-flesh-coloured spot.

On dead branches. Penzance.

With a merulioid aspect. *B. & Br.* Name—*aurum*, gold. *Fr. Hym. Eur. p. 613. B. & Br. n. 1970.*

**30. H. denticulatum** Pers.—*Light-yellow-ochraceous, shining*, longitudinally effused, somewhat fleshy, subiculum thin, smooth, somewhat mealy. **Spines** crowded, equal, slightly toothed at the margin.

Colour bright yellowish almost between light-yellow and egg-yellow.

On rotten wood. Penzance.

Name—*dens*, a tooth. From the *slightly toothed* spines. *Pers. Myc. Eur. 2. p. 181. Fr. Hym. Eur. p. 614. B. & Br. n. 1971.*

**31. H. alutaceum** Fr.—**Subiculum** *pallid-ochraceous*, longitudinally effused, 7.5–12.5 cent. (3–5 in.) long, *crustose*, adnate, smooth, *circumference naked*. **Spines** minute, crowded, equal, acute.

In colour like *Grandinia granulosa*, but wholly distinct.

On dead wood. Rare.

Name—*aluta*, tanned leather. From the colour. *Fr. Hym. Eur. p. 614. Syst. Myc. 1. p. 417. Berk. Out. p. 260. C. Hbk. n. 862.*

**32. H. sordidum** Weinm.—*Dingy yellowish*, broadly effused, thin, easily separating, mucid, *at the circumference gyroso-porous, sulphur-yellow*. **Spines** very crowded, compressed, incised, rather acute.

In the form of *Merulius* at the circumference, spines somewhat gelatinous, appearing as if composed of several growing together, smooth.

On dead wood. Stoke Poges. Nov.

Name—*sordidus*, dirty, dingy. Of colour. *Weinm. Ross. p. 370. Fr. Hym. Eur. p. 614. B. & Br. n. 1688.*

**33. H. limonicolor** B. & Br.—Adnaté, bright lemon-yellow.

Hydnum.

**Spines** crowded, acute, short; mycelium shining white, scanty or obsolete.

On stone buried among pine-leaves. Glamis, 1876. Oct.

Name—lemon-colour. *Citrus Limonum*. B. & Br. n. 1686. *S. Mycol. Scot.* n. 812.

**34. H. spathulatum** Fr.—**Subiculum** whitish-yellowish, effused, membranaceous, separable, circumference fimbriated. **Spines** spathulate, oblique, orange.

Membrane villous beneath.

On dead wood. Rare. Jan.

Name—*spatha*, a spatula. Spoon-shaped. *Fr. Hym. Eur.* p. 614. *Berk. Out.* p. 261. *C. Hbk.* n. 863. *S. Mycol. Scot.* n. 813. *Schwein. Car.* n. 993.

**35. H. multiforme** B. & Br.—Ochrey-white, at first even, resembling a *Corticium*, at length here and there fertile. **Spines** very acute, crowded, becoming pallid and fimbriate; texture floccoso-mealy.

Very variable, sometimes almost towy, with the margin inflexed.

On dead wood. Glamis, 1877.

Name—*multus*, many; *forma*, form. Of many forms. B. & Br. n. 1687. *S. Mycol. Scot.* n. 814.

**36. H. anomalum** B. & Br.—Pallid light yellow; stratum thin, gelatinous; teeth in the form of granules, then stipitate and obtusely divided upwards.

Substance of teeth tough, with large ovate or globose vesicles immersed in it. Near to Fries's genus *Mucronella*.

In the inside of very rotten ash-tree. Langridge, Somersetshire. March.

Spores globose, shortly pedicellate. B. & Br. Name—*ἀνόμελος*, irregular. B. & Br. n. 1438, with fig.

**37. H. melleum** B. & Br.—Honey-colour, effused, thin, margin delicately flaxy; subiculum and teeth, which are acute and sometimes divided at the apex, pulverulent downwards, naked at the middle.

On broken rails on the ground. Coed Coch, 1873.

Name—*mel*, honey. Honey-colour. B. & Br. n. 1436.

**38. H. sepultum** B. & Br.—Wholly resupinate, golden, margin white. **Spines** acute, medium-sized.



Forming little scattered patches.

On stones buried among pine-leaves. Glamis, 1876. Forbes. Sept.

Name—*sepelio*, to bury. Buried. *B. & Br. n.* 1813. *S. Mycol. Scot. n.* 820.

\*\*\* *Spines flesh-colour, lilac, rufescent.*

**39. H. udum** Fr.—**Subiculum** *flesh-colour then watery-yellowish*, effused, thin, *somewhat gelatinous*, agglutinated, *smooth*. **Spines** crowded, unequal, 2-3-forked and fimbriated, of the same colour as the subiculum.

Very long and broad. Not separable from the wood. Allied to *H. membranaceum*.

On dead branches. Frequent. Oct.

Forming elongated patches 10-12.5 cent. (4-5 in.) long. *M. J. B.* Name—*udus*, moist. *Fr. Hym. Eur. p.* 615. *Syst. Myc. i. p.* 422. *Berk. Out. p.* 261. *t. 17. f. 3.* *C. Hbk. n.* 864. *S. Mycol. Scot. n.* 815.

**40. H. bicolor** Alb. & Schw.—**Subiculum** *white*, widely effused, 20 cent. (8 in.) and more, *tomentose, adpressed*, thin. **Spines** minute, curt, 1 m. ( $\frac{1}{2}$  lin.) long, *white-villous*, naked at the apex, acute, rufous-date-brown.

On bramble. Batheaston. March.

Name—*bis, color*. Of two colours. *A. & S. p.* 270. *Fr. Hym. Eur. p.* 615. *Syst. Myc. i. p.* 417. *B. & Br. n.* 1297.

\*\*\*\* *Spines white, or light yellowish when old.*

**41. H. nodulosum** Fr.—Whitish, very broadly effused, crustaceous, closely adnate, smooth, even and at the same time *nodulose*. **Spines** very long, on the even part depressed and adnate to the wood, on the lower side of the nodules free and pendulous.

The nodules which vary in size are sterile above, even and smooth, pallid yellowish, but furnished with spines on the side facing the ground.

On fir stumps. Glamis, 1876. Oct.

Name—*nodus*, a knot. Nodulose. *Fr. Hym. Eur. p.* 616. *S. Mycol. Scot. n.* 816.—*H. macrodon* *Fr. Monogr. ii. p.* 279, not Pers. *Pers. Myc. Eur. 2. t. 2, 22, f. 1.*

**42. H. Stevensoni** B. & Br.—White, effused, mealy beneath, here and there flaxy. **Spines** cylindrical obtuse or truncate, sometimes compressed, pulverulent at the apex.

Hydnum. On dead wood. Glamis, 1874. March.

Name—after Rev. John Stevenson. *B. & Br. n.* 1437. *S. Mycol. Scot. n.* 817.

**43. H. niveum** Pers.—Shining white. **Subiculum** effused, 5–7.5 cent. (2–3 in.) and more long, thin, *membranaceous*, *adnate*, *circumference flaxy*. **Spines** crowded, curt, short, equal, smooth.

*B. fimbriatum* circumference fibrilloso-fimbriate.

On dead wood. Uncommon. April–Nov.

Name—*nix*, snow. Snow-white. *Pers. Disp. t. 4. f. 6, 7.* *Fr. Hym. Eur. p.* 616. *Syst. Myc. i. p.* 419. *Berk. Out. p.* 261. *C. Hbk. n.* 865. *S. Mycol. Scot. n.* 818. *Nees Syst. f.* 246.

**44. H. farinaceum** Pers.—White. **Subiculum** effused, indeterminate, closely *adnate*, *mealy crustose*, circumference somewhat flocculose. **Spines** thin, somewhat distant, very acute, quite entire.

It varies yellowish.

On dead wood. Frequent. Oct.–Nov.

Forming thin effused patches, resembling scattered meal. *M. J. B.* Name—*farina*, meal. *Pers. Syn. p.* 562. *Fr. Hym. Eur. p.* 616. *Syst. Myc. i. p.* 419. *Berk. Out. p.* 261. *C. Hbk. n.* 866. *S. Mycol. Scot. n.* 819.—*Fl. Dan. t.* 1375.

**45. H. argutum** Fr.—White. **Subiculum** effused, in scattered patches, *tomentoso-interwoven*, *adnate*. **Spines** acute, awl-shaped, unequal, under a lens serrated or slightly toothed at the sides.

On wood and bark. Bodelwyddan. Twycross. Sept.

Distinguished by its spongy subiculum, which consists of rather strong perpendicular threads. *B. & Br.* Name—*argutus*, acute. Of the spines. *Fr. Hym. Eur. p.* 616. *B. & Br. n.* 1026. *C. Hbk. n.* 867.—*Roth. Ust. Ann. 1. t. 1. f.* 5 s.

**46. H. stipatum** Fr.—Whitish. **Subiculum** very widely effused, *flocculoso-furfuraceous*, forming an *adnate* crust. **Spines** crowded, in the form of granules, rather obtuse and slightly toothed.

The circumference is sterile or covered with spines. It varies in colour, isabelline or light yellowish.

On rotten wood. Carlisle.

Name—*stipatus*, crowded. Of the spines. *Fr. Hym. Eur. p.* 617. *Gre-villea, vol. xii. p.* 98.

GENUS XXXIII.—**Hericium** Pers. Comm. Clav. p. 23. partly. *Hericium*.  
Fr. Syst. Orb. Veget. p. 88.

Fleshy, club-shaped, in place of a pileus divided at the apex into spines, which are extended upwards, not inferior and facing the ground. *Growing on trunks.*

An anomalous genus, resembling *Clavariæ*, hymenium amphigenous, but undoubtedly nearest to *Hydna*. *Fr. Hym. Eur. p. 617.*

No British species.

GENUS XXXIV.—**Tremellodon** (*Tremella*, a gelatinous genus of fungi; *ὀδὺς*, a tooth). Pers. Myc. Eur. p. 172. *Tremello-*  
*don.*


Gelatinous, pileate, aculeate below; spines awl-shaped, equal. *Fr. Hym. Eur. p. 618.*

**1. T. gelatinosum** Pers.—**Pileus** glaucous-fuscous, *gelatinous*, tremulous, dimidiate, somewhat stipitate, covered with small pimples. **Spines** soft, glaucous.

Extremely variable in form, stipitate (on the ground) and sessile.

On fir, trunks and sawdust. Rare. Sept.—Oct.


Of a soft gelatinous consistence approaching that of a *Tremella*. Pileus fan-shaped, or rounded in front, attenuated behind so as to make a short spurious stem. In an early stage it is decurrent at the point of attachment. The surface is of a bistre-brown, uneven with frequent depressions, and rough at first with little points, but at length nearly smooth. The hymenium is either pure white or shaded with a delicate blue tint, the spines straight, of moderate length, and very delicate. The plant shrinks much in drying. *M. f. B.* Of singular beauty, almost translucent with steel-blue tints shading into violet, while the spines are of a pure soft white. Spores round, somewhat irregular, white, 2 mk. *W. G. S.* Name—*gelatina*, jelly. Gelatinous. *Pers. Scop. Carn. 2. p. 472. Fr. Hym. Eur. p. 618. S. Mycol. Scot. Supp. Scot. Nat. 1885, p. 75. Hydnum gelatinosum Berk. Out. p. 259. C. Hbk. n. 854. Schaff. t. 144. Jacqu. Austr. t. 239. Krombh. t. 50. f. 18-22.—Fl. Dan. t. 717. Var. with pines throughout Jacqu. Misc. i. t. 9.*



LXXV. *Tremellodon gelatinosum*. One-third natural size.  
Section twice natural size.

Sistotrema. GENUS XXXV.—**Sistotrema** (*sisto*, to place; *τρήμα*, a pore. From the arrangement of the pores). Pers.—Fr. Syst. Myc. i. p. 426.

Fleshy, with an inferior hymenium spread over teeth resembling broken gills. Gills somewhat waxy, *irregularly arranged* (not radiating), *distinct*, bent, easily rubbed off from the pileus. Sporophores 4-spored, spores oval. *Irregular in form or dimidiate*.



LXXXVI. *Sistotrema confuens*.  
Natural size. Section twice  
natural size.

A genus nearest to *Irpex*, but clearly distinct by reason of the gill-like, irregularly arranged, distinct, non-concatenate teeth, and by the fleshy substance. *Fr. Hym. Eur. p. 618*.

1. **S. confuens** Pers. — *Entire, shining white*. Pileus 1–2.5 cent. ( $\frac{1}{2}$ –1 in.) broad, fleshy, irregular, horizontal, villous. Stem 2.5 cent. (1 in.) and less long, *somewhat excentric*. Gills (tooth-like plates) flexuous, changeable in form.

Simple, but cæspitose and growing into each other.

On the ground and sticks. Uncommon. Autumn.

At length yellowish or tinged with brown. Tooth-like plates of the hymenium entire or jagged. *Grev.* Name—*confuo*, to flow together. Confluent. Pers. Syn. p. 551. *Fr. Hym. Eur. p. 619*. *Syst. Myc. i. p. 426*. *Berk. Out. p. 262*. *C. Hbk. n. 869*. *S. Mycol. Scot. n. 821*. *Grev. t. 248*. —*Bull. t. 453. f. 1*. *Sow. t. 112*.

*Irpex*. GENUS XXXVI.—**Irpex** (*irpex*, a harrow). Fr. Elench. p. 142.

Hymenium inferior, toothed from the first. Teeth firm, somewhat coriaceous, acute, concrete with the pileus, arranged in rows or like network, connected at the base by folds, which are gill-like (in sessile species) or resemble honeycomb (in resupinate ones). Sporophores 4-spored. *Growing on wood, somewhat sessile or resupinate, approaching Lenzites and Dædaleæ*. *Fr. Hym. Eur. p. 619*.

\* *Stipitate.*

\*\* *Pendulous with the pileus extended behind.*

\*\*\* *Sessile or effuso-reflexed, marginate.*

\*\*\*\* *Resupinate.*

\* *Stipitate.*

No British species.


\*\* *Pendulous with the pileus extended behind.*

1. **I. pendulus** Fr.—**Pilei** light yellow, *extended behind, pendulous*, membranaceous, plicate, adpressedly squamuloso-pilose. **Teeth** in rows, large, incised, shining white.

Membrane of the pileus elastic, but when extended it contracts again. Its characters are wholly those of the genus, but it is not allied to any other plant among the *Hydne*. It is nearly allied to *Cyphella*, more remotely to *Trogia*, and might almost be regarded as the type of a separate genus.

On pine. Rare. Castle Semple (Edin. Herbarium). Menmuir.

Name—*pendo*, to hang. Pendulous. *Fr. Hym. Eur. p. 620. Berk. Out. p. 262. C. Hbk. n. 870. S. Mycol. Scot. n. 822. —Alb. & Schw. p. 261. t. 6. f. 7.*



LXXVII. *Irpex obliquus*. Natural size. Section twice natural size.

\*\*\* *Sessile or effuso-reflexed, marginate.*

2. **I. fusco-violaceus** Fr.—**Pileus** 5 cent. (2 in.) long, more than 2.5 cent. (1 in.) broad, *white inclining to hoary*, effuso-reflexed, coriaceous, silky, *zoned*. **Teeth** in rows in the form of plates, *fuscous-violaceous*, incised at the apex.

It differs certainly from *Polyporus abietinus*. It is marked with a few depressed, concentrically sulcate zones of the same colour. Margin deflexed, often repand. Often wholly effused.

On pine trunks. Rare. Leigh Woods.

Name—From the colour of the teeth. *Fr. Hym. Eur. p. 620. Elench. p. 144. Br. Bath. Trans. 1870, p. 87. C. Hbk. n. 873. Kl. Bor. t. 536. —Willd. Bot. Mag. iv. t. 2. f. 5.*

\*\*\*\* *Resupinate.*

3. **I. Johnstoni** Berk.—Resupinate, 5 cent. (2 in.) long, pure white, coriaceous-membranaceous, separable from the matrix, cir-

Irpex.

cumference naked. **Teeth** compressed, unequal, *arranged in rows*.

Effused, with the margin reflexed all round, and the teeth exactly resembling those of many true species of *Hydnum*, but upon minute inspection they will be found to be seated upon fine folds.

On dead beech. Berwick.

Name—after Dr George Johnstone. *Berk. Out.* p. 262. *C. Hbk.* n. 871. *S. Mycol. Scot.* n. 823. *Fr. Hym. Eur.* p. 621. *Irpex lacteus Eng. Fl.* v. p. 161.

**4. I. spathulatus** Fr.—Shining white, effused, membranaceous, at first flaxy, at length becoming smooth. **Teeth** *spathulate, equal*, entire, reticulato-connected with obsolete veins.

Very distinct, never porous. Here and there yellowish when dry.

On larch. Wallington, Northumberland. Nov.

Name—*spatha*, a spatula. From the spoon-shaped teeth. *Fr. Hym. Eur.* p. 622. *B. & Br.* n. 1815.—*Schrad. Spic. t.* 4. f. 3.

**5. I. obliquus** Fr.—White inclining to pale, effused, forming an adnate crust, circumference flaxy. **Teeth** *extended from a base resembling honeycomb, compressed, unequal, incised*, oblique.

At first abundantly porous, but toothed from the first, at length quite as in *Hydnum*.

On stumps and dead branches. Uncommon. Nov.—Feb.

Name—*obliquus*, oblique. From the teeth. *Fr. Hym. Eur.* p. 622. *Berk. Out.* p. 262. *C. Hbk.* n. 872. *S. Mycol. Scot.* n. 824.—*Bolt.* t. 167. f. 1.

**6. I. deformis** Fr.—White, effused, crustaceous, thin, circumference pubescent, somewhat flaxy. **Teeth** *extended in awl-shape from a minutely porous base, thin*, somewhat digitato-incised.

It approaches the *Polypori*.

On wood. Strachan, Kincardineshire.

Name—*deformis*, misshapen. *Fr. Hym. Eur.* p. 622. *S. Mycol. Scot.* n. 825.

**7. I. carneus** Fr.—*Reddish*, effused, 2.5–7.5 cent. (1–3 in.) long, *cartilaginous-gelatinous*, membranaceous, adnate. **Teeth** obtuse and awl-shaped, entire, united at the base.

It inclines to *Radula* and *Phlebia*.

On wood. Penzance.

When perfect it is a true *Irpex*. *B. & Br.* Name—*caro*, flesh. Flesh-colour. *Fr. Hym. Eur.* p. 622. *B. & Br.* n. 1972.

GENUS XXXVII.—**Radulum** (from the *root-like* appearance of the processes of the hymenium). Fr. Elench. p. 148. Radulum.

Hymenium amphigenous, tubercular. Tubercles rude, irregularly shaped, commonly elongate, obtuse, waxy (except in *R. aterrimum*), irregularly arranged, distinct. Sporophores 4-spored, occurring not only on the tubercles, but also on the even parts of the hymenium. *Growing on wood, resupinato-effused, often bursting through the bark.* Fr. Hym. Eur. p. 622.

\* *Denuded, very variable.*

\*\* *Innate, removing the bark.*

\* *Denuded, very variable.*

1. **R. orbiculare** Fr.—A. *First season's growth, white then yellowish, orbicular, confluent, circumference flaxy. Tubercles elongated, somewhat round, scattered or fasciculate.* B. *Second season's growth, vernal, flesh-colour, waxy-fleshy, smooth. Tubercles softer, shorter, as if rubbed away.*



LXXVIII. *Radulum quercinum.*  
Natural size.

Entirely changeable in form, but very distinct. The tubercles vary flexuous.

On dead bark, birch, cherry, &c. Frequent. Oct.

5-7.5 cent. (2-3 in.) broad, quite membranaceous or above 4 mm. (2 lin.) thick. *M. f. B.* Name—*orbiculus*, a small disc. Orbicular, round. Fr. Hym. Eur. p. 623. Berk. Out. p. 263. C. Hbk. n. 874. S. Mycol. Scot. n. 826. Grev. t. 278.

2. **R. quercinum** Fr.—White then pallid, 5-7.5 cent. (2-3 in.), somewhat round then broadly confluent, innate, crustaceous, *becoming smooth. Tubercles round, stout, obtuse, scattered or fasciculate, irregularly shaped, floccoso-villous at the apex.*

Rude, changeable in form, often confounded with *Hydnum squalinum* and *H. varicolor*.

On dead oak. Rare. Dec.

Name—*quercus*, oak. Growing on oak. Fr. Hym. Eur. p. 623. Syst. Myc. i. p. 423. Berk. Out. p. 263. C. Hbk. n. 875. S. Mycol. Scot. n. 827. —Ray Syn. t. 1. f. 4.

Radulum.

**3. R. tomentosum** Fr.—White inclining to pale, effused, innate, rather thick, *circumference swollen, erect, tomentose*. **Tubercles** short, crowded together, angular, obtuse, smooth, confluent.

When dry pale wood-colour, the tomentum sometimes becoming ferruginous.

On *Pyrus aucuparia*. Menmuir, Forfarsh. New Pitsligo, Aberdeensh.

Name—*tomentum*, down. Downy. *Fr. Hym. Eur. p. 624. B. & Br. n. 1439. S. Mycol. Scot. n. 828.*

**4. R. deglubens** B. & Br.—Orbicular, 8 mm. ( $\frac{1}{3}$  in.) across, ferruginous, somewhat diaphanous. **Tubercles** erect, somewhat cylindrical, irregular, scattered; interstices even, pulverulent with the white spores.

On ash. Sanquhar, Moraysh. New Pitsligo, Aberdeensh. Jan.

Name—*deglubo*, to peel off. Peeling off. *B. & Br. n. 1440. S. Mycol. Scot. n. 829.*

**5. R. corallinum** B. & Br.—Effused, white; subiculum shining, very thin, pelliculose. **Tubercles** fasciculate (fascicles 6 mm.,  $\frac{1}{4}$  in. or more across), divided downwards, obtuse, coralloid.

Effused for 7.5 cent. (3 in.) over branches partially covered with lichens.

On oak branches. Glamis, 1874. Sept.

Name—*κοράλλιον*, coral; *εἶδος*, appearance. Coral-like. *B. & Br. n. 1441. S. Mycol. Scot. n. 830.*

**6. R. epileucum** B. & Br.—Effused for several inches, ochrey-white, wholly resupinate; subiculum snow-white, covered by a waxy stratum. **Tubercles** scattered, cylindrical, under a lens fimbriated at the apex, deciduous.

Tubercles falling out and showing the white mealy subiculum, round which is an annular depression.

On decorticated wood. Glamis, 1874. Aug.

Name—*ἐπί, λευκός*, white. Whitish. *B. & Br. n. 1442. S. Mycol. Scot. n. 831.*

**\*\* Innate, removing the bark.**

**7. R. fagineum** Fr.—*White*, becoming pale, innate, removing the bark, smooth. **Tubercles** elongated, round, various, obtuse, *entire*.

Surrounding decorticated branches like stalactite.



On dead beech. Epping Forest. Feb.—March.

Radulum.

Name—*fagus*, beech. Growing on beech. *Fr. Hym. Eur. p. 624. C. Hbk. n. 876.—Pers.*

**8. R. aterrimum** Fr.—*Black*, innate, removing the bark, *strigose*. **Tubercles** elongated, distant, large, irregularly shaped, somewhat compressed, black.

Hymenium not continuous.

On branches. Kensington Gardens.

Berkeley now regards *Hydnum erectum* Sow. *Herb.* as identical. Name—*ater*, black. Very black. *Fr. Hym. Eur. p. 624. B. & Br. n. 1814, 1573.*


GENUS XXXVIII.—**Phlebia** (φλέψ, a vein, from the veiny appearance of the hymenium). *Fr. Syst. Myc. i. p. 426.*

Hymenium amphigenous, waxy-soft, smooth, continuous, *from the first wrinkled into crests*; covered with wrinkles which are crowded, interrupted, persistent, quite entire at the edge, everywhere bearing spores. *Resupinato-effused, somewhat gelatinous - waxy when moist, cartilaginous when dry.* *Fr. Hym. Eur. p. 624.*

**1. P. merismoides** Fr. — *Flesh-colour then livid, villous and white beneath, orange-strigose at the circumference*, effused, even or slightly branched in becoming incrustated. **Wrinkles** simple, straight, crowded.

The wrinkles are never porous or reticulate.

On stumps and branches. Rare. Sept.—Feb.



LXXIX. *Phlebia merismoides*.  
Natural size.

Often running over mosses. Thin, tremelloid when fresh. *M. F. B. 5-7.5 cent. (2-3 in.) broad.* Name—*Merisma*, εἶδος, appearance. Merisma-like. *Fr. Hym. Eur. p. 624. Berk. Out. p. 263. C. Hbk. n. 877. S. Mycol. Scot. p. 156. Grev. t. 280. Huss. ii. t. 44.*

**2. P. radiata** Fr.—*Red-flesh-colour*, 2.5-7.5 cent. (1-3 in.) across, somewhat round, equal, *smooth* on both sides, *circumference radiate in the form of teeth*. **Folds** straight, radiating in rows.

Phlebia.

Thinner and brighter-coloured than the rest.

On dead wood. Rare. Nov.

Almost orange. *M. F. B.* Spores curved, hyaline,  $4.5 \times 1-1.5$  mk. *K.* Name—*radiatus*, rayed. From the folds. *Fr. Hym. Eur. p. 625. Syst. Myc. i. p. 427. Berk. Out. p. 263. C. Hbk. n. 878. S. Mycol. Scot. p. 156. —Sow. t. 291.*

**3. P. contorta** Fr.—*Rufous then fuscous*, effused, slightly firm, smooth on both sides, circumference indeterminate. **Folds** *here and there conglomerated*, or slightly branched, somewhat flexuous, irregularly arranged.

Very much allied to *P. radiata*.

On decayed wood. Rare.

Name—*contorqueo*, to twist. Contorted. Of the more intricate folds. *Fr. Hym. Eur. p. 625. Syst. Myc. i. p. 427. Berk. Out. p. 264. C. Hbk. n. 879. —Pers. Myc. Eur. 2. t. 18. f. 5.*

**4. P. vaga** Fr.—*Dingy yellowish*, effused, 5 cent. (2 in.) and more, adnate, *circumference flaxy-fibrillose*. **Hymenium** yellowish-grey, gradually formed by the union of creeping intricate wrinkles, which when perfect terminate in a granulose papilla.

The hymenium in perfect specimens is wholly that of the genus, waxy, but the wrinkles are at length densely granulose with papillæ.

On dead wood. Common.

Arachnoid at first, then traversed with intricate, fructifying veins rather than wrinkles, which multiply rapidly and form an intricate mass. *M. F. B.* Name—*vagus*, wandering. From the peculiar mode of formation of the hymenium. *Fr. Hym. Eur. p. 625. Syst. Myc. i. p. 428. Berk. Out. p. 264. C. Hbk. n. 880. S. Mycol. Scot. p. 156.*

**5. P. lirellosa** Pers. — Umber-grey, resupinate, free at the margin. **Sinuses** very small, linear, intermixed with pores.


The sinuses at first sight assume the appearance of small cracks, but, when examined under a lens, many of them appear forked, others simple, some poriform.

Habitat not recorded. Penzance.

Found by Persoon on dried branch of black poplar. Name—*lira*, a ridge or furrow. From the appearance of the small sinuses. *Dædalea Pers. Myc. Eur. iii. p. 2. t. 17. f. 2, 3. Phlebia B. & Br. n. 1973.*

GENUS XXXIX.—**Grandinia** (*grando*, hail. From the granular *Grandinia*. character of the hymenium). Fr. Gen. Hymenom.

Hymenium amphigenous, continuous, waxy, warty with papillæ or rather granulose; granules globose or hemispherical, entire, obtuse or hollowed out at the apex, crowded, regular, smooth, persistent. *Crustaceous, effused, soft, improperly referred on account of their equal and persistent papillæ to the Thelephoræ, from which they are distinguished by their polished granules.* Fr. Hym. Eur. p. 625.



LXXX. *Grandinia granulosa*.  
Natural size. Section four  
times natural size.

1. **G. granulosa** Fr.—*Tan-colour*, waxy, broadly effused, agglutinated, circumference determinate, smooth; hymenium equal, continuous. **Granules** hemispherical, equal, crowded.

On dead branches. Common. Oct.—Feb.

Forming a thin adnate stratum. *M. F. B.* Name—*granum*, a small seed. Granular. Fr. Hym. Eur. p. 626. Berk. Out. p. 264. *C. Hbk. n. 882. S. Mycol. Scot. n. 832.—Pers. Syn. p. 576.*

2. **G. mucida** Fr.—*Pale yellowish*, waxy-mucid, effused, somewhat innate, circumference determinate, somewhat radiate; hymenium continuous. **Granules** crowded, *large, unequal*, hemispherical, soft.

Somewhat gelatinous when moist, corrugated when dry.

On rotting pine. Glamis, 1877. May—Nov.

Name—*mucidus*, mucid. Fr. Hym. Eur. p. 626. *B. & Br. n. 1691. S. Mycol. Scot. n. 833.*

3. **G. ocellata** Fr.—*Livid*, waxy, broadly effused, agglutinated, the indeterminate circumference sterile; hymenium unequal, continuous. **Granules** crowded, *somewhat conical, obtuse*, equal, of the same colour.

Unchanged when dry, not cracked, sterile in some places. The granules terminate in a very delicate whitish apiculus, and on this sinking in the apex is hollowed out forming as it were a small eye.

On dead trunks. Bodolwyddan. Coed Coch. Aug.—Sept.

Grandinia.

Name—*ocellus*, a little eye. From the eyelet at the apex of the granules *Fr. Hym. Eur. p. 626. B. & Br. n. 1027. C. Hbk. n. 883.*

4. *G. papillosa* Fr.—*Milk-white*, yellowish beneath, membranaceous, somewhat round, *separating when entire*, smooth beneath, circumference furfuraceous; hymenium very much cracked. **Granules** minute, crowded, somewhat spherical, equal.

On sticks. Rare.

Name—*papilla*, a teat. From the granules. *Fr. Hym. Eur. p. 626. B. & Br. n. 282. C. Hbk. n. 881. S. Mycol. Scot. n. 834.*

5. *G. crustosa* Fr.—White, *floccoso-mealy*, irregularly effused, crustaceous, adnate. **Granules** somewhat round, crowded, obtuse, of the same colour at the apex.


On *Polyporus versicolor*. Dun, 1874. Feb.

Name—*crusta*, a crust. Crustaceous. *Fr. Hym. Eur. p. 627. B. & Br. n. 1443. S. Mycol. Scot. n. 835.—Pers. Syn. p. 562. Nees Syst. f. 247.* Var. *lignorum* A. & S. Often yellowish, granules unequal, often hollowed out at the apex. On rotting pine. Glamis, 1877. May. *B. & Br. n. 1690. S. Mycol. Scot. n. 835. Fr. Hym. Eur. p. 627.*

Odontia.

GENUS XL.—*Odontia* (ὀδούς, a tooth). Fr. Gen. Hymen.

Hymenium composed of interwoven fibres, which form papillose, rarely spine-shaped warts, with a multifid crest or pencilled at the apex. *Resupinato-effused*, arid and not waxy like the preceding, tending more towards *Hydna*. *Fr. Hym. Eur. p. 627.*



LXXXI. *Odontia fimbriata*.  
Natural size.

1. *O. barba-Jovis* Fr.—White then yellowish, flaxy-membranaceous, effused, equal. **Warts** at first papillose then conical, almost 4 mm. (2 lin.) long, with an orange fringe at the apex.

The only species in which the warts resemble spines. Subiculum expanded to 20 cent. (12 in.) and more.

On decayed wood. Epping Forest.

Sowerby's figure seems to represent the true plant, but the specimens in his herbarium are *Radulum quercinum*. *B. & Br.* Name—Jove's beard. *Fr.*

*Hym. Eur. p. 627. Syst. Myc. i. p. 421. B. & Br. n. 1299.*—*Hydnum Odontia. With.—Sow. t. 328. Bull. t. 481. f. 2.*

**2. *O. fimbriata* Pers.**—Pallid, effused, membranaceous, separating, *traversed by root-like ribs*, circumference fibrilloso-fringed. **Warts** minute, *in the form of granules*, multifid at the apex, *rufescent*.


Elegant. The type of the genus.

On dead wood. Uncommon. Sept.–Oct.

Frequently tinged with lilac. *M. F. B.* Name—*fimbriæ*, a fringe. From the fringed circumference. *Pers. Obs. i. p. 88. Fr. Hym. Eur. p. 627. Berk. Out. p. 264. C. Hbk. n. 884. S. Mycol. Scot. n. 836.*

GENUS XLI.—***Kneiffia*** (after Kneif, a promising mycologist of Baden, who died young). *Fr. Gen. Hymen.* *Kneiffia.*

*Hymenium* amphigenous, *continuous, united*, but incomplete (imperfect), similar (to the hymenophore), destitute of granules and warts, *strigose and roughened with rigid, scattered or fasciculate bristles*. Sporophores single-spored; spores elliptic. Laxly fleshy, soft, when dry collapsing and becoming flocculose. *Swelling out of chinks in the bark of trees into various forms, hemispherical in its most perfect state.* *Fr. Hym. Eur. p. 628.*



LXXXII. *Kneiffia setigera.*  
Natural size.

**1. *K. setigera* Fr.**—Milk-white, irregularly effused, thick, fleshy, undulated, flocculose within, silky beneath, beset with scattered hyaline bristles.

Of no constant form, but certainly a typical plant, not a degenerate condition of any other, and *the type of a most distinct genus*, in its substance and white colour comparable only with *Corticium serum*, but differing in the fructification. The bristles are rigid, equal, not to be compared with those of *Auricularini*. The largest specimens are of the size of an apple, but commonly thin, effused, often incrusting *Sphæria*.

On dead wood. Uncommon. April.

Somewhat resembling *Grandinia granulosa* in general appearance. *M. F. B.* Spores elliptic, 11–14 mk. *B. & Br.* Name—*seta*, a bristle; *gero*, to carry.

Kneiffia.

Bristly. *Fr. Hym. Eur. p. 628. Elench. p. 208. Berk. Out. p. 265. B. & Br. n. 1299\*. t. 18. f. 1. C. Hbk. n. 885. S. Mycol. Scot. n. 837.*

**2. K. subgelatinosa** B. & Br.—Yellowish then cream-colour, thin. **Granules** minute, somewhat gelatinous, fringed at the apex.

Accompanied by a green *Alga*, which penetrates the tissue of the fungus.

On stumps of fir. Glamis, 1874. April.

Name—*sub, gelatinosus*, somewhat gelatinous. *B. & Br. n. 1444. S. Mycol. Scot. n. 838.*

Mucronella.

GENUS XLII.—**Mucronella** (*mucro*, a sharp point ; dim.)

Receptacle none. Spines awl-shaped, simple, acute, smooth, definitely facing the ground. *Small*, growing on wood. *Fr. Hym. Eur. p. 629.*

No British species.

## ORDER IV.—THELEPHOREI.

Hymenium inferior or amphigenous, *coriaceous or waxy, even, rarely ribbed or papillose*. Sporophores 4-spored. *The beginning of a new series with the hymenium not figurate.*

The hymenium is commonly said to be papillose, but these (so-called) papillæ are for the most part spurious or accidental, generally arising from incrustated *Sphæriæ*, and disappearing when the fungus grows on an even surface. Thus also the occasional papillæ in *Stereum rubiginosum*, *hirsutum*, &c., arise from the rough surface on which they grow. In a few species *papillæ* are present when they are in full vigour, but these collapse when the plant dries. *Fr. Hym. Eur. p. 629.*

GENUS XLIII.—**Craterellus** (*crater*, a bowl).

Craterellus.

Fr. Gen. Hymen.

Hymenium waxy-membranaceous, distinct, but adnate to the hymenophore, definitely inferior, continuous, smooth, even or rugose. Spores white. *Growing on the ground, fleshy or membranaceous, furnished with an entire pileus, stipitate, autumnal, allied to Cantharelli.* *Fr. Hym. Eur. p. 630.*

\* *Tubæform, pervious to the base of the stem.*


\*\* *Infundibuliform, stem stuffed.*

\*\*\* *Irregularly shaped, pileus and stem fleshy.*

\* *Tubæform, pervious to the base of the stem.*

1. **C. lutescens** Fr.—**Pileus** 2.5–10 cent. (1–4 in.) broad, *fuscous*, somewhat membranaceous, tubæform, soon pervious, undulated, *flocculose*.

**Stem** about 5 cent. (2 in.) long, hollow, smooth, *yellow*. **Hymenium** yellow, &c., *remotely ribbed*, even then rugose with interwoven veins.



LXXXIII. *Craterellus lutescens*.  
One-half natural size.

Craterellus. Odour strong, spirituous. Hymenium yellow, inclining to reddish, orange, and bluish-grey. Formerly confounded with *Cantharellus tubæformis* (*lutescens*).

In moist woods. Rare.

Name—yellowish. *Fr. Hym. Eur. p. 630. Berk. Out. p. 265. C. Hbk. n. 886. S. Mycol. Scot. n. 839.—Schæff. t. 157. Bolt. t. 105. f. 2. Pers. Myc. Eur. 2. t. 13. f. 1.*

**2. C. cornucopioides Pers.**—*Pileus blackish-fuliginous*, somewhat membranaceous, tubæform, pervious, *squamulose*. **Stem** hollow, smooth, black. **Hymenium even**, at length slightly wrinkled, cinereous.

Very variable in size. Very like *Canth. cinereus*, but differing in the structure of the hymenium.

In woods. Uncommon. Sept.

Distinguished from *Canth. cinereus* by the spores. In this moreover the sporophores are forked above, and the spicules long and often less than four in number; in *Canth. cinereus* the sporophores are obtuse, and the spicules four. *B. & Br.*

*Pileus 7.5 cent. (3 in.) or more broad. More or less tufted. M. & B. Spores 10 × 4 mk. B. & Br.; ellipsoid, hyaline, 12-14 × 7-8 mk. K. Name—cornucopiæ, horn of plenty. From its shape. Pers. Myc. Eur. 2. p. 5. Fr. Hym. Eur. p. 631. Berk. Out. p. 266. t. 19. f. 6. B. & Br. n. 1139\*. C. Hbk. n. 887. S. Mycol. Scot. n. 840. Schnizl. Sturm. 31. t. 5. Hussey ii. t. 37.—Krombh. t. 46. f. 10-13. Fl. Dan. t. 588, 1260. Bolt. t. 103. Sow. t. 74. Schæff. t. 165, 166.*

**\*\* Infundibuliform, stem stuffed.**

**3. C. sinuosus Fr.**—*Pileus 2.5 cent. (1 in.) broad, fuscous-grey*, slightly fleshy, infundibuliform, undulated and floccose. **Stem** 2.5 cent. (1 in.) and more long, *stuffed*, pallid cinereous. **Hymenium at length with interwoven wrinkles**, pallid cinereous.

Strong-smelling, "moschatus" (Secr.) Hymenium like that of *Cr. lutescens*.

In mixed woods. Frequent. Sept.-Nov.

Name—*sinus*, a curve. Sinuous, undulated. *Fr. Hym. Eur. p. 631. Berk. Out. p. 266. C. Hbk. n. 888. S. Mycol. Scot. n. 841.—Vaillant Par. t. 11. f. 11-13.*

**4. C. crispus Fr.**—*Pileus fuliginous becoming fuscous*, fleshy-membranaceous, somewhat pervious, *crisped*. **Stem** stuffed at the base, *pallid*. **Hymenium even, pallid**.

Commonly confounded with *Thelephora undulata*.

In mixed woods. Frequent. Oct.-Nov.

This and *C. sinuosus* are so near that probably they are one species. Name



—*crispa*, a curl. Crisped. *Fr. Hym. Eur. p. 631. Berk. Out. p. 266. C. Craterellus. Hbk. n. 889. S. Mycol. Scot. n. 842.—Sow. t. 75. Hussey ii. t. 18.*

\*\*\* *Irregularly shaped, pileus and stem fleshy.*

**5. C. clavatus** Fr.—**Pileus** 5 cent. (2 in.) broad, somewhat light yellowish, fleshy, *turbinate, truncate* or depressed, flexuous, unpolished, *attenuated into the solid stem*; flesh thick, white. **Hymenium** even, then corrugated, purplish then changing colour.

Solitary or cæspitose, occasionally branched. The hymenium is white-pruinose with the spores, not separate, violaceous-flesh-colour passing into fuliginous or umber; hence the following forms are distinct: A. violaceous then date-brown. B. flesh-colour.—*Schæff. t. 164. Schmid. Ic. 2. t. 60. C. purplish. Schæff. t. 276. D. umber.—Wulf. in Jacq. Coll. ii. t. 12. f. 3.*

In beech wood. Bisham, Berks.

Name—*clavis*, a club. Club-shaped. *Fr. Hym. Eur. p. 632. Sv. atl. Sv. t. 91. B. & Br. n. 1445.—Krombh. t. 45. f. 13-17.*

#### GENUS XLIV.—**Thelephora** (θηλή, a teat; φέρω, to bear).

Thelephora.

Ehrh. Fr. Gen. Hymen.

Hymenium inferior or amphigenous, continuous with the hymenophore and similar to it, even or ribbed, without an intermediate stratum. Sporophores 4-spored. *Coriaceous, destitute of a cuticle, very varied in form (pileate, clavate, resupinate), growing on the ground. Fr. Hym. Eur. p. 633.*


\* *Erect, pileus entire or divided into branches.*

\*\* *Pileate, dimidiate, horizontal, somewhat sessile or effuso-reflexed.*

\*\*\* *Resupinate, for the most part incrusting, wherefore the forms are various.*

\* *Erect, pileus entire or divided into branches.*

**1. T. Sowerbeii** B. & Br.—Snow-white, at length changing colour, here and there dingy yellow. **Pileus** coriaceous, entire, *infundibuliform, rough with radiating processes* projecting from the surface. **Hymenium** smooth (not bristly).



LXXXIV. *Thelephora laciniata*.  
One-third natural size.

*Thelephora*. The true plant of Sowerby, whose figure is very faithful though evidently taken from discoloured specimens.

On the ground. Rare.

Name—after Sowerby. *B. & Br. n.* 1027. *Berk. Out. p.* 266. *C. Hbk. n.* 890. *Fr. Hym. Eur. p.* 633.—*Sow. t.* 155.

**2. *T. multizonata* B. & Br.—****Pileus** bright rufous-flesh-colour above, *many-zoned, multiplex*, infundibuliform, formed from various confluent lobes and stems, margin lobed and crenulate. **Hymenium** slightly ribbed, smooth, paler.

Forming a dense mass, of a beautiful reddish tint; flesh and stem zoned within. Quite distinct from Sowerby's plant.

On the ground. Rare.

Name—*multus*, many; *zona*, a zone. *B. & Br. n.* 1028. *t.* 13. *f.* 4. *C. Hbk. n.* 891. *Fr. Hym. Eur. p.* 633.

**3. *T. undulata* Fr.—***Pallid.* **Pileus** 1–2.5 cent. ( $\frac{1}{2}$ –1 in.) broad, coriaceo-membranaceous, depressed, even, *slightly smooth*, margin entire and undulated. **Stem** curt, 12 mm. ( $\frac{1}{2}$  in.) long, villous. **Hymenium** *ribbed, slightly bristly*.

**Pileus** central, entire, very thin, almost diaphanous and plano-infundibuliform.

On the ground. Coed Coch, 1878. Oct.

Spores ellipsoid,  $4-5 \times 2-2\frac{1}{2}$  mk. *K.* Name—*unda*, a wave. Undulated. *Fr. Hym. Eur. p.* 633. *B. & Br. n.* 1816.—*Schæff. t.* 278 near it but perhaps distinct.

**4. *T. caryophyllea* Pers.—***Fuscous-purple.* **Pileus** somewhat coriaceous, depressed, *fibrous-torn*, margin sometimes incised, sometimes divided into a few linear branches. **Stem** short. **Hymenium** *rather even, smooth*.

Inodorous.

On the ground about roots, especially fir. Rare.

It assumes every form, from that of a perfect cup with a central stem to a much and irregularly branched frond. *M. J. B.* Spores  $6 \times 5$  mk. *W. G. S.*; sphaeroid, tuberculate, fuscous, about 8 mk. *K.* Name—from its likeness to a carnation (*Dianthus caryophyllus*). *Pers. Syn. p.* 565. *Fr. Hym. Eur. p.* 634. *Berk. Out. p.* 267. *C. Hbk. n.* 894. *S. Mycol. Scot. n.* 843. *Saund. & Sm. t.* 41. *f.* 2.—*Schæff. t.* 325. *Schnizl. t.* 6.

**5. *T. tuberosa* Fr.—***Pallid then rufescent.* **Pileus** somewhat coriaceous, *divided to the bulbous stem into compressed branches which are arranged in an infundibuliform manner.* **Hymenium** inferior, smooth.

On the ground. Rare. Foxhall, Edinburgh. Moncreiffe. Thelephora.

Scattered, subcoriaceous, about 2.5 cent. (1 in.) high; pileus about two-thirds of the height of the entire plant, variously divided into compressed, acute or obtuse branches. The main branches are given off from the same point, and are disposed in a circular manner, leaving the centre free, and somewhat infundibuliform; stem nearly cylindrical, obscurely furrowed, or lacunose, bulbous at the base. Hymenium covering the whole plant except the stem. Spores oval, numerous. *Grev.* Name—*tuber*, a tuber. Tuberous. *Fr. Hym. Eur. p. 634. Berk. Out. p. 267. C. Hbk. n. 892. S. Mycol. Scot. n. 844.—Grev. t. 178.*

**6. T. anthocephala** Fr. — Somewhat ferruginous, becoming fuscous. **Pileus** coriaceous-soft, pubescent, *divided to the equal villous stem into flaps which are dilated upwards* and fringed and whitish at the apex, or divided into irregular branched erect branches. **Hymenium** even.

Extremely variable, inodorous.

On the ground in woods. Uncommon.

Branches spreading upwards into greyish or purplish-brown, strongly streaked branchlets, disposed frequently like the petals of a pink, their apices dilated, pale and generally fimbriated. Very variable as to the form of the bleached laciniae (flaps). Sometimes regular as in Bulliard's fig. *M. f. B.* Name—*άνθος*, a flower; *κεφαλή*, the head. From its resemblance to a flower. *Fr. Hym. Eur. p. 634. Berk. Out. p. 267. t. 17. f. 4. C. Hbk. n. 893. S. Mycol. Scot. n. 845.—Bull. t. 452. f. 1.—B. Th. digitata Fr. Sow. t. 156. Berk. Out. t. 17. f. 4.*

**7. T. clavularis** Fr.—Rufous-fuscous. Coriaceous, soft, irregularly branched, branches *round, attenuated, even*, smooth, delicately pruinose; apices acute, pubescent, whitish. **Stem somewhat tuberous.**

Smaller and thinner than neighbouring species, inodorous.

On the ground. Wallington, Northumberland.

Name—*clavula*, a small nail. From the shape of its branches. *Fr. Hym. Eur. p. 634. B. & Br. n. 1575.*

**8. T. palmata** Fr.—The whole 2.5–5 cent. (1–2 in.) high, *fuscous-purple*, coriaceous-soft, erect, very much branched, *pubescent*, the simple base stem-like; branches palmate, flattened, even, somewhat fastigiata, fringed and whitish at the apex.

Odour foetid. Becoming ferruginous when dry. The lobes are wedge-shaped, occasionally arranged in the form of small infundibuliform pilei.

On the ground in woods. Frequent. Aug.–Nov.

Forming a dense cluster, 2.5–7.5 cent. (1–3 in.) thick. Smell shortly after being gathered corpse-like. Spores subsphaeroid, slightly fuscous, tuberculate, 8–12 mk. *K.* Name—*palma*, the palm. From the branches resembling

Thelephora. a palmate leaf. *Fr. Hym. Eur. p. 634. Berk. Out. p. 267. C. Hbk. n. 895. S. Mycol. Scot. n. 846. Holmsk. Ot. i. p. 106. t. 10. Nees f. 151.—Grev. t. 46. Krombh. t. 54. f. 24-25.*

**\*\* Pileate, dimidiate, horizontal, &c.**

**9. T. intybacea** Pers. — Cæspitose, *whitish then rufous-ferruginous*, at length fuliginous, soft. **Stems** somewhat lateral, growing into each other. **Pileoli** imbricated, fibrous, dilated *at the margin, at first white-fringed*, then entire and of the same colour. **Hymenium** inferior, papillose, somewhat floccose.

Tufts sometimes central and obconic, sometimes lateral with somewhat triangular pileoli.

On the ground in Scotch fir wood. Glamis, 1875.

Name—from its likeness to succory (*Cichorium intybus*). *Pers. Syn. p. 567. Fr. Hym. Eur. p. 635. B. & Br. n. 1576. S. Mycol. Scot. n. 847. Bull. t. 483. f. 6, 7, t. 278.*

**10. T. terrestris** Ehrb. — Cæspitose, 2.5-5 cent. (1-2 in.) broad, *fuscous, becoming black*, soft. **Pileoli** imbricated, becoming plane, fibrous-strigose, zoneless, extended into a somewhat lateral stem, *margin similar*. **Hymenium** inferior, radiato-rugose.

Smaller than *T. intybacea*. The stems when confluent become also central; sometimes the pilei are lacinate, scaly, merismoid.

In fir woods. Uncommon.

Name—*terra*, the earth. Growing on the ground. *Ehrb. Cr. n. 179. Fr. Hym. Eur. p. 635. Berk. Out. p. 267. C. Hbk. n. 896. S. Mycol. Scot. n. 848. Nees Syst. f. 251. Batsch f. 121. Kl. Bor. t. 473.*

**11. T. laciniata** Pers. — *Ferruginous-fuscous*, coriaceous-soft, incrusting. **Pilei** somewhat imbricated, *effuso-reflexed, fibrous-scaly; margin fibrous-fringed*, at first whitish. **Hymenium** inferior, papillose, flocculose.

Perennial, growing again in strata.

On stumps, chiefly Scotch fir. Common. Aug.-Nov.

Larger than *T. terrestris*. The fibres being adnate form little ridges rather than scales. *M. J. B.* Spores angulato-sphaeroid, tuberculate, somewhat fuscous, 6-9 mk. *K.* Name—*lacinia*, a flap. From the shape of the pilei. *Pers. Syn. p. 567. Fr. Hym. Eur. p. 636. Berk. Out. p. 268. C. Hbk. n. 899. S. Mycol. Scot. n. 849.—Bolt. t. 173. Sow. t. 213. Fl. Dan. t. 1198, 949.*

**12. T. biennis** Fr. — *Cinereous-fuscous*, coriaceous-soft, broadly incrusting. **Pilei** at length narrowly reflexed, *tomentose*, circumference fringed. **Hymenium** somewhat resupinate, *smooth*, very slightly bristly, plicate at the base.

It corresponds with *T. terrestris*, but is more vigorous in growth, thinner, *Thelephora*. broader, not striate beneath. Biennial, not growing again.

On the ground, incrusting stones, &c. Rare.

Name—*biennis*, lasting two years. *Fr. Hym. Eur. p. 636. Berk. Out. p. 268. C. Hbk. n. 900. S. Mycol. Scot. n. 850.—Bull. t. 436. f. 2.*

**13. *T. mollissima*** Pers.—Scarcely 2.5 cent. (1 in.) long, fleshy-soft, incrusting. *Pilei whitish*, effuso-reflexed, *forming flaps, somewhat tomentose. Hymenium* inferior, smooth, even, fuscous-purple.

Soft to the touch.

On the ground in woods. Frequent. Oct.

Sometimes quite effused, sometimes assuming the form of *T. palmata*. *M. f. B. Name—mollis*, soft. *Very soft. Pers. Syn. p. 572. Berk. Out. p. 268. t. 17. f. 5. C. Hbk. n. 898. S. Mycol. Scot. n. 851.*

\*\*\* *Resupinate, for the most part incrusting, &c.*

**14. *T. cristata*** Fr.—*Pallid*, incrusting, slightly tough, passing into branches or ascending flaps, *the apices somewhat awl-shaped or fringed. Hymenium* papillose on even patches and on the sides of the branches.

Not flaxy when young. No form constant. The primary and typical condition, on the under surface of beech-leaves, is even, hyaline-fuscous, adpressed at the circumference, fibrillose. It varies with few, obconic, penicillate branches.

On the ground in woods among mosses. Frequent.

Name—*crista*, a crest. Crested. *Fr. Hym. Eur. p. 637. Berk. Out. p. 268. C. Hbk. n. 897. S. Mycol. Scot. n. 852. Fl. Dan. t. 2272.—Bull. t. 415. f. 1. Sow. t. 158.—Jungh. Linn. v. t. 7. f. 2.*

**15. *T. fastidiosa*** Fr.—*Shining white*, effused, soft, shapeless, growing into each other by incrusting, *passing into plate-like branches. Hymenium* inferior, papillose, rufescent.

In its foetid odour it agrees with *T. palmata*. The tufts are at first without distinct form, confluent mutually and with neighbouring bodies in a circle of as much as 30 cent. (12 in.) wide.

On the ground. Uncommon.

Looks at first sight like some mycelium. *M. f. B. Name—fastidiosus*, disgusting. From the odour. *Fr. Hym. Eur. p. 637. Berk. Out. p. 268. C. Hbk. n. 901. Saund. & Sm. t. 41. f. 1.*

**16. *T. sebacea*** Pers.—Whitish, effused, fleshy-waxy, becoming hard, changeable in form by incrusting, tubercular or resembling stalactite, *circumference similar. Hymenium* collapsing, *flocculoso-pruinose*.

Thelephora. Circumference not penicillate.

On stumps, logs, grass, &c. Common. Jan.—Oct.

Name—*seba*, tallow. From its consistence. *Pers. Syn. p. 577. Fr. Hym. Eur. p. 637. Berk. Out. p. xvii. t. 17. f. 6. C. Hbk. n. 904. S. Mycol. Scot. n. 853. Letell. t. 607. f. 3.—Pers. Clavæf. t. 4. f. 4. Fl. Dan. t. 1302. f. 2.*

**17. T. crustacea** Schum.—*Fuscous-umber, broadly effused, somewhat fleshy, undulated and tubercular, the similar circumference white-floccose. Hymenium papillose, slightly bristly.*

On the ground. Burnham.

Name—*crusta*, a crust. Crustaceous. *Schum. Sæll. p. 396. Fr. Hym. Eur. p. 637. B. & Br. n. 1577. Fl. Dan. t. 1851. f. 2.*


**18. T. cæsia** Pers.—*Cinereous-bluish-grey, effused, determinate, soft, continuous. Hymenium even, slightly bristly.*

A very thin, even, grey-cinereous form approaching this has occurred on sterile heathy ground.

On stumps and the ground. Frequent. Nov.

The surface is sometimes quite smooth. With its quaternate spores it is a pretty opaque object under the microscope. *M. J. B.* Name—*cæsius*, bluish-grey. *Pers. Obs. Myc. i. t. 3. f. 6. Fr. Hym. Eur. p. 638. Berk. Out. p. 268. C. Hbk. n. 903. S. Mycol. Scot. n. 854. Nees Syst. f. 254.*

Cladoderris. GENUS XLV.—**Cladoderris** (κλάδος, a young branch; δέρρις, a leathern coat. From its coriaceous and branched structure). Pers.



Coriaceous; hymenium woody, radiated on branched ribs, persistent, at length roughened with warts. *Fr. Sum. Veg. Scan. p. 332.*

**1. C. minima** B. & Br.—White, 4–6 mm. (2–3 lin.) across, flabelliform, springing from a stem-like or obsolete base, resupinate. **Pileus** tomentose. **Hymenium** radiated on branched ribs.

Though small it has exactly the structure of the exotic species.

On birch. Glamis, 1876. Dec.

Name—*minimus*, least. Very small. *B. & Br. n. 1692. S. Mycol. Scot. n. 855.*

LXXXV. *Cladoderris minima*.  
Natural size. Section three times  
natural size.

GENUS XLVI.—**Stereum** (στερεός, hard). (Pers.) Fr. Epicr. Stereum.  
p. 545.

Hymenium definitely inferior, coriaceous, separated from the fibrous-coated pileus by an intermediate fibrillose stratum, even, remaining unchanged. Sporophores 4-spored; spores minute, roundish-ovate. *Coriaceous or woody, mostly perennial, somewhat zoned, entire, definite in form. Fr. Hym. Eur. p. 638.*

I. MERISMA (μερίζω, to divide). Pilei very numerous, ascending, imbricated in a free tuft.

II. APUS (ἄ, πούς, a foot). Pileus sessile, at first resupinate, then commonly pileato-reflexed or adnate behind, coriaceous, marginate.

\* *Hymenium smooth.*

\*\* *Hymenium simple, slightly bristly or velvety.*

\*\*\* *Woody, perennial, scarcely reflexed, but marginate, the hymenium strato-se by confluent pilei, at the first pruinose.*

\*\*\*\* *Rigid, the simple hymenium pruinose.*

III. RESUPINATA. Crustaceo-adnate, *indeterminate*, without a distinct pileus, but at length admitting of being detached entire like a piece of leather. *Circumference not flaxy. Incrusting wood.* (Species of Subdivision II. (Apus) occur in a resupinate state, but being limited (marginate) are from the first distinct.)

\* *True species, coriaceous.*

\*\* *Imperfect, crustaceo-adnate, sterile.*



LXXXVI. *Stereum hirsutum.*  
Natural size.

# I.—MERISMA.

No British species.

## II.—APUS. Pileus sessile, at first resupinate, &c.

\* *Hymenium smooth.*

1. **S. ochroleucum** Fr.—Pileus hoary, coriaceous-membranaceous, free, expanded, *flaccid, silky, zoned*. **Hymenium** even, smooth, yellowish (or in var. grey).

Commonly large, with zones of the same colour or fuscous. Coating of the pileus here and there separating.

On wood. Glamis, 1874. Jan.

Name—ὤχρως, pale yellow; λευκός, white. *Fr. Hym. Eur. p. 639. B. & Br. n. 1818. S. Mycol. Scot. n. 856.*

Stereum.

**2. S. purpureum** Pers.—*Coriaceous-soft*. **Pileus** pallid or whitish, effuso-reflexed, somewhat imbricated, zoned, *villosotomentose*. **Hymenium** naked, even, smooth, *purplish*.

On dead wood, especially fir and poplar. Common. May-Dec.

Varying greatly in colour, whitish, yellowish, pallid-lilac, &c., with frequently a black zone near the margin. *M. f. B.* Name—*purpureus*, purple. *Pers. Obs. Myc.* 2. p. 92. *Fr. Hym. Eur.* p. 639. *Berk. Out.* p. 270. *C. Hbk.* n. 910. *S. Mycol. Scot.* n. 857. *Hussey i. t.* 20.—*Bull. t.* 483. *Sow. t.* 388. f. 1.

**3. S. vorticosum** Fr.—**Pileus** pallid, margin of the same colour, coriaceous, effuso-reflexed, obscurely zoned, *strigoso-hirsute*. **Hymenium** somewhat ribbed, smooth, *purplish*.

Exactly intermediate between *S. purpureum* and *S. hirsutum*; the colour is that of the former, the coating and entire structure as in the latter. It resembles *Auricularia* in the widely ribbed hymenium.

On beech. Menmuir, Forfarshire.

Name—*vorticosus*, full of eddies. From the ribbed hymenium. *Fr. Hym. Eur.* p. 639. *B. & Br.* n. 1578. *S. Mycol. Scot.* n. 858. *Bull. t.* 483. f. 1-5. *Bolt. t.* 82. f. D?

**4. S. hirsutum** Fr.—Coriaceous, stiff. **Pileus** somewhat zoned, inclining to pale, effused and reflexed, *strigoso-hirsute*, margin rather obtuse, *yellow*. **Hymenium** even, smooth, naked, juiceless, *yellowish*, varying in colour.

There are many varieties, even persistently resupinate. The down which covers it is yellowish, pallid or grey.

On stumps, &c. Very common. Jan.—Dec.

Name—*hirtus*, hairy. *Fr. Hym. Eur.* p. 639. *Berk. Out.* p. 270. t. 17. f. 7. *C. Hbk.* n. 911. *S. Mycol. Scot.* n. 859. *Hussey i. t.* 58. *Theleph. Fr. Syst. Myc.* i. p. 439.—*Bull. t.* 274. *Sow. t.* 27. *Fl. Dan.* t. 1199. *Michel. t.* 66. f. 2.

**5. S. spadiceum** Fr.—Coriaceous. **Pilei** somewhat ferruginous, effuso-reflexed, *villos*, the rather obtuse margin white. **Hymenium** even, smooth, inclining to *fuscous*, *bleeding if bruised* when fresh.

The colour varies in intensity. *Intermediate stratum white*. Very distinct from *S. sanguinolentum*.

On stumps and branches, chiefly ash and oak. Common. Oct.—Dec.

Name—*spadix*, date-brown. *Fr. Hym. Eur.* p. 640. *Berk. Out.* p. 270. *C. Hbk.* n. 912. *S. Mycol. Scot.* n. 860.—*Fl. Dan.* t. 1619. f. 1. *Bull. t.* 483. f. 5.



**6. S. sanguinolentum** Fr.—Coriaceous, thin. **Pileus** pallid, *Stereum*. effused and reflexed, *adpressedly silky*, somewhat striate, *the acute margin* white. **Hymenium** even, smooth, *cinereous inclining to fuscous, bleeding when touched*, pruinose when old.

It must not be confounded with *S. rugosum*. It is distinguished by its thin substance, by the pileus being adpressedly silky and somewhat striate, by the hymenium never being yellowish, &c.

On wood, chiefly pine. Common. June–Nov.

Name—*sanguis*, blood. Bleeding. *Fr. Hym. Eur. p. 640. Berk. Out. p. 271. C. Hbk. n. 913. S. Mycol. Scot. n. 861. Theleph. Fr. Elench. n. 178. Grev. t. 225. Kl. Bor. t. 381.*

**\*\* Hymenium simple, slightly bristly or velvety.**

**7. S. rubiginosum** Fr.—Coriaceous-rigid. **Pileus** rubiginous, effuso-reflexed, somewhat fasciate, velvety, becoming smooth and date-brown, *the intermediate stratum tawny-ferruginous*. **Hymenium** ferruginous, velvety with small bristles.

Margin paler when young.

On stumps, &c. Common. July–Nov.

Spores cylindrical, curved. *Fr. Name—rubigo, rust. Fr. Hym. Eur. p. 641. Hymenochæte Berk. Out. p. 271. C. Hbk. n. 916. S. Mycol. Scot. n. 866. Theleph. Fr. Syst. Myc. i. p. 436. Fl. Dan. t. 1619. f. 2. Sow. t. 26 (the under side downy).*

**8. S. tabacinum** Fr.—Coriaceous, thin, flaccid. **Pileus** somewhat ferruginous, *the margin, and intermediate stratum which is composed of filaments, golden*, effused, reflexed, silky, at length becoming smooth. **Hymenium** paler, pubescent with small bristles.

The hymenium varies in colour. Very widely effused.

On fallen branches. Rare.

Spores elongated, slightly curved, hyaline,  $3\text{--}5 \times 1$  mk. *K.* Name—*tabacum*, tobacco. From the colour. *Fr. Hym. Eur. p. 641. Hymenochæte Berk. Out. p. 271. C. Hbk. n. 917. Theleph. Fr. Syst. Myc. i. p. 437.—Sow. t. 25. Bolt. t. 174.*

**9. S. rufo-hispidum** Stev.—Pallid fawn-colour, rigid, margin obtuse, raised, hispid with rufous bristles.

A very distinct species.

On bark of yew. Craig-y-barns, Dunkeld. Sept.

Name—from the rufous bristles. *Hymenochæte Stevensoni B. & Br. n. 1817. S. Mycol. Scot. n. 868.*

Stereum.

**10. S. avellanum** Fr.—Coriaceous, *hard*. **Pileus** dingy, effused and, as well as the *obtuse*, free, *narrowly reflexed*, *date-brown margin*, *villous*. **Hymenium** even, velvety then pruinose and becoming smooth, pale-ferruginous (here and there bleeding).

In many respects it approaches *S. rugosum*, but is certainly very different. Colour not yellowish.

On bark and wood, especially hazel. Appin.

Name—*Corylus avellana*, hazel. *Fr. Hym. Eur.* p. 642. *S. Mycol. Scot.* n. 862. *Theleph. Fr. Syst. Myc.* i. p. 442. *Berk. Eng. Fl.* v. p. 167.

\*\*\* *Woody, perennial, scarcely reflexed, but marginate, &c.*

**11. S. frustulosum** Fr.—Date-brown-blackish, woody, resupinate, tubercular, crowded and as if confluent, then *broken up into fragments*, smooth beneath and at the obsoletely marginate circumference. **Hymenium** *convex*, *cinnamon becoming pale*, pruinose.

Easily distinguished by the colours, and by its thick, very hard, woody substance. Hymenium when perfect facing the ground, at first pruinose, when fertile pulverulent with the cinnamon spores; when turned to the light it becomes sterile, pale, denuded.

On hard oak wood. South of England.

Spores ellipsoid-sphaeroid, hyaline  $4-5 \times 3-4$  mk. *K.* Name—*frustum*, a piece. From its being broken up into fragments. *Fr. Hym. Eur.* p. 643. *B. & Br. n.* 1300. *Theleph. Fr. Syst. Myc.* i. p. 445. *Elench. p.* 191.

\*\*\*\* *Rigid, the simple hymenium pruinose.*

**12. S. rugosum** Fr.—Corky, rigid. **Pileus** *date-brown*, effused and shortly reflexed, *obtusely marginate*, *at length smooth*. **Hymenium** unpolished, pruinose, somewhat bleeding when bruised.

A. Hymenium yellowish. B. Hymenium cinereous-livid. Varying very much, even cup-shaped.

On stumps, &c. Very common. Jan.—Dec.

Name—*ruga*, a wrinkle. *Fr. Hym. Eur.* p. 643. *Berk. Out. p.* 271. *C. Hbk. n.* 914. *S. Mycol. Scot. n.* 863.

**Var. lauro-cerasi** Berk. On laurel. *S. Mycol. Scot. n.* 863. *Theleph. Berk. Eng. Fl.* v. p. 173.

**13. S. pini** Fr. — *Pallid*, coriaceous-cartilaginous, resupinate, adnate in the form of a shield, somewhat marginate, *smooth beneath*, *at length bullate*. **Hymenium** purplish-flesh-colour, becoming fuscous, pruinose.

The primary shields are minute, crowded, but already free, never agglutinated or waxy.

On bark of Scotch fir. Glamis. Aviemore.

Stereum.

Name—*pinus*, pine. *Fr. Hym. Eur. p. 643. B. & Br. n. 1579. S. Mycol. Scot. n. 864.*

**14. S. rufum** Fr.—Rufous, becoming fuscous, coriaceous-cartilaginous, erumpent, *tuberculiform then somewhat round, marginate*, smooth beneath. **Hymenium** grey-pruinose, at length bullate-tubercular.

Wholly *immarginate* when young.

On dead pine and ash. Glamis, 1874. Sept.—March.

Name—*rufus*, rufous. *Fr. Hym. Eur. p. 644. Elench. p. 187. B. & Br. n. 1580. S. Mycol. Scot. n. 865.—Sow. t. 338. f. 2.*

III.—RESUPINATA. Crustaceo-adnate, *indeterminate*, &c.

\* *True species, coriaceous.*

**15. S. stratosum** B. & Br.—Effused, bright ochraceous-white, smooth, becoming yellow, here and there wrinkled.

Substance pallid, stratose, the strata at length broken up.

Habitat not recorded. Penzance.

Name—*stratum*. Stratose. *B. & Br. n. 2027.*

\*\* *Imperfect, crustaceo-adnate, sterile.*

**16. S. acerinum** (Theleph. *acerina* Pers.)—Crustaceo-adnate, even, smooth, white.

Effused like the crust of a lichen. It must be sought for in a perfect condition on fallen trunks. When young it is continuous and somewhat pruinose. Name—*acer*, maple. *Pers. Syn. p. 581. Fr. Hym. Eur. p. 645. Syst. Myc. i. p. 453. Berk. Out. p. 271. C. Hbk. n. 915.*

GENUS XLVII.—**Auricularia** (*auricula*, the ear). Bull.

Auricularia.

Champ. p. 277. sp. 1.—*Fr. Epicr. p. 555.*

Hymenium definitely inferior, remotely and irregularly costatoplicate, swelling and becoming gelatinous and tremulous when moist, collapsing (and coriaceous *Epicr.*) when dry. *Habit exactly that of Stereum.*

Intermediate between *Thelephorei* and *Tremellini*; nearest in affinity to the former, when dry scarcely to be distinguished from *Stereum*; departing from the *Tremellæ* in the coriaceous pileus. *Fr. Hym. Eur. p. 645.*

Auricularia.

1. **A. mesenterica** Fr.—*Pilei fuscous-cinereous*, resupinate then reflexed, *entire*, villous, fasciato-zoned. **Hymenium** costato-plicate, fuscous-violaceous.



LXXXVII. *Auricularia mesenterica*. One-third natural size.

On stems of trees. Frequent.

5-7.5 cent. (2-3 in.) broad, gelatinous in wet weather, hard and cartilaginous when dry, reviving with wet. Hymenium powdered with a beautiful bloom. *M. f. B.* Spores white,  $7 \times 4$  mk. *W. G. S.* Name—*μεσέντερον*, the mesentery. *Fr. Hym. Eur. p. 646. Berk. Out. p. 272. C. Hbk. n. 919. —Sow. t. 290. Hussey ii. t. 6. Bolt. t. 172. Bull. t. 290. Michel. t. 66. f. 4.*

2. **A. lobata** Sommerf.—As much as 17.5 cent. (7 in.) long, 5-12.5 cent. (2-5 in.) broad, 2-4 mm. (1-2 lin.) thick. **Pileus** fuscous-whitish, effuso-reflexed, lobed, variegated with strigoso-tomentose, velvety and smooth zones. **Hymenium** fuscous-livid, reticulato-ribbed with distant folds.

Very much allied to *A. mesenterica*.

On bark of trees. Uncommon.

Name—*lobatus*, lobed. *Sommerf. in Mag. Nat. Vidensk. 1827*, with fig. *Fr. Hym. Eur. p. 646. Elench. ii. p. 34. Berk. Out. p. 272. t. 18. f. 1. C. Hbk. n. 920.*

Corticium.

GENUS XLVIII.—**Corticium** (*cortex*, bark). *Fr. Epicr. p. 556.*

Hymenium amphigenous, even or tubercular, springing immediately from the mycelium without an intermediate stratum; in typical species fertile and swelling when moist, fleshy-soft, contracted and hence commonly cracked when dry, or the whole breaking up. There are other species added to the genus with the hymenium arid and incrusting. Sporophores 4-spored, rarely single-spored. *Resupinate, growing on wood, often sterile. Fr. Hym. Eur. p. 646.*

I. **LOMANTIA** (λωμα, a fringe). Resupinate, but free at the circumference, determinate, marginate, commonly cup-shaped then expanded.

II. **HIMANTIÆ** (ιμάς, a strap). Resupinate, effused, immarginate, clothed with flaxy fibrils or strigosely hairy at the circumference and beneath. *Often sterile, fibrillose (mycelia). In their perfect state they are furnished with a remarkably well developed, waxy-soft hymenium.*

\* *Mycelium and circumference white.*

\*\* *Mycelium and circumference coloured.*

III. LEIOSTROMA (λεῖος, smooth; στρώμα, a stratum). Agglutinated (rarely when mature becoming free), without a strigose or fibrillose circumference. For the mycelium passes at once into the hymenium which is agglutinated to the substratum, but this (the hymenium) occurs of two kinds, one thicker, waxy-soft, fertile; the other thin and sterile. *The most of the species are undistinguished in appearance.*

\* *Waxy, when fertile facing the ground, uniform, rimose when dry.*

\*\* *Amphigenous, turning rigid, circumference at first very thinly floccose and radiating, soon naked.*

\*\*\* *Amphigenous, at the very first hardened, somewhat grumous, circumference always naked.*

\*\*\*\* *Amphigenous, very thin, innate, removing the bark.*

\*\*\*\*\* *Less known species, doubtful.*


Subgenus I. CONIOPHORA (κόνις, dust; φέρω, to bear). Pers. Myc. Eur. i. p. 153. Fleshy, undulate and tubercular, rarely membranaceous, becoming even; hymenium smooth, pulverulent with the spores; in other respects agreeing with *Corticium*.

Growing on wood, analogous to the group *Coniophori* of *Merulius*.

Subgenus II. HYPOCHNUS (ὑπό and χνός, the first down on the chin). Fr. Obs. ii. p. 278. Breaking up and becoming floccose or furnished with a tomentose, somewhat pulverulent hymenium.

\* *Breaking up and becoming floccose.*

\*\* *Hymenium tomentose, somewhat pulverulent.*



LXXXVIII. *Corticium sambuci*.  
Natural size.

I.—LOMANTIA. Resupinate, but free at the circumference, &c.

1. *C. evolvens* Fr.—Resupinate, marginate or effuso-reflexed, soft, flocculose with whitish down beneath, zoneless. **Hymenium naked, smooth**, somewhat wrinkled, *fuscous becoming pale fuscous*, divided in cracks when dry.

So polymorphous in stature and form, that at first sight the different forms seem scarcely allied. It varies when smaller cup-shaped, marginato-reflexed; wholly resupinate, broadly effused, immarginate. It is distinguished by the hymenium being soft, somewhat tubercular, very much cracked when dry, fuscous or becoming pale, and by the floccose receptacle.

On dead wood, especially cherry. Frequent. Oct.—Feb.

Name—*evolve*, to unroll. Fr. Hym. Eur. p. 646. B. & Br. n. 1029. C. Hbk. n. 921. S. Mycol. Scot. n. 872.

2. *C. salicinum* Fr.—White-villous externally, coriaceous, soft, rigid when dry, persistently *cup-shaped, adfixed at the centre*.

Corticium. **Hymenium** rather even, naked, *persistently red-blood-colour, continuous when dry.*

Very elegant, entirely like a *Peziza*. It varies becoming smooth.

On *Salix aurata*. Kinrara, Morayshire. Aug.-Sept.

Berkeley considers this the same with *Exidia cinnabarina* B. & C. Spores oblong, curved, hyaline, 16-18 × 4-6 mk. *K.* Name—*salix*, willow. *Fr. Hym. Eur. p. 647. B. & Br. n. 1581. S. Mycol. Scot. n. 873.*

**3. C. amorphum** Fr.—4-6 mm. (2-3 lin.) broad, white-tomentose externally, waxy-pliant, somewhat coriaceous, *cup-shaped then flattened*, confluent, marginate. **Hymenium** even, continuous, pale.

On silver-fir. Perth. Dunkeld. Keir. Aug.-Oct.

The substance is white and fleshy, consisting of rather coarse threads, which at the base form a close sclerotoid network. The hymenium consists of colourless threads, and orange-coloured clavate bodies filled with pigment. *B. & Br.* There is considerable doubt whether this is a true *Corticium*. Fries says the habit is exactly that of *Corticium* of this section, but the structure of the disc is ambiguous. Berkeley, who compares it in appearance with *Peziza calycina*, considers it may perhaps be the type of a new genus. Name—*ἀμορφή*, form. Without regular form. *Fr. Hym. Eur. p. 648. B. & Br. n. 1582. S. Mycol. Scot. n. 874. Theleph. Fr. Elench. p. 183.*

II.—HIMANTIÆ. Resupinate, effused, immarginate, &c.

\* *Mycelium and circumference white.*

**4. C. giganteum** Fr.—*Hyaline-white*, very broadly effused, swelling when moist, waxy, *when dry cartilaginous, of the consistence of paper, free*, milk-white, circumference strigoso-radiate. **Hymenium** even, continuous.

At the first agglutinated, but in contracting with dryness it breaks up, and does not become cracked. Smaller and less perfect forms occur.

On pine, stumps, &c. Common. Oct.-April.

Spores ellipsoid or sphaeroid-ellipsoid, hyaline, 4-5 × 3 mk. *K.* Name—*giganteus*. Of very large size. *Fr. Hym. Eur. p. 648. Berk. Out. p. 272. C. Hbk. n. 922. S. Mycol. Scot. n. 875.*

**5. C. lacteum** Fr.—*Milk-white*, effused, membranaceous, *laxly fibrillose* beneath and at the circumference. **Hymenium** (when perfect) waxy, deeper in colour, *divided by cracks* when dry.

Much smaller and thinner than *C. giganteus*.

On trunks, &c. Frequent. Nov.-Feb.

Name—*lac*, milk. Milk-white. *Fr. Hym. Eur. p. 649. Berk. Out. p. 273. C. Hbk. n. 923. S. Mycol. Scot. n. 876.—Bonord. f. 259.*

**6. C. porosum** B. & Curt.—Resupinate, milk-white, here and Corticium. there porous, margin free, reflexed.

The pores look as if little dew-drops had settled on the hymenium, which had in consequence contracted or rather retracted.

On wood. Aboyne.

Name—*porosus*, porous. *B. & Br. n. 1821. S. Mycol. Scot. n. 886.*

**7. C. arachnoideum** Berk.—Effused, delicately flaxy as is the circumference. **Hymenium** white, very thin, patchy.

The mycelium is as delicate as a spider's web.

On wood, lichens, &c. Frequent. Oct.—Feb.

Name—*ἀράχνη*, a spider. From the spider-web mycelium. *Berk. Out. p. 273. C. Hbk. n. 924. S. Mycol. Scot. n. 877. Fr. Hym. Eur. p. 649.*

**8. C. foetidum** B. & Br.—Effused, resupinate, arachnoid beneath, white then ochraceous and smooth.

Not apparently a resupinate form of *Thelephora fastidiosa*.

On sawdust. Coed Coch.

Name—*foetidus*, foetid. *B. & Br. n. 1824.*

**9. C. læve** Pers.—Effused, membranaceous, separating, villous beneath, *circumference flaxy* (not fibrilloso-radiate). **Hymenium** even, smooth, *somewhat flesh-colour* or livid.

On dead wood, sticks, &c. Very common. Nov.—March.

Hymenium sometimes pure white. *M. J. B.* Spores ellipsoid or ovoideo-sphaeroid, subchlorino-hyaline,  $12 \times 8-10$  mk. *K.* Name—*lævis*, even. *Pers. Disp. p. 30. Fr. Hym. Eur. p. 649. Berk. Out. p. 273. C. Hbk. n. 925. S. Mycol. Scot. n. 878.—Bonord. f. 251.*

**10. C. roseum** Pers.—*Rose-colour*, effused, adnate, circumference fringed and whitish. **Hymenium** *pruinose, becoming pale*, at length rimoso-corrugated, hardened.

Very pretty, cognate with *Hyphelia rosea*.

On dead wood, birch, poplar, &c. Frequent. Oct.—Feb.

Name—*rosa*, a rose. Rose-coloured. *Pers. Disp. p. 31. Fr. Hym. Eur. p. 650. Berk. Out. p. 273. C. Hbk. n. 926. S. Mycol. Scot. n. 879.*

**\*\* Mycelium and circumference coloured.**

**11. C. velutinum** Fr.—*Flesh-colour*, effused, adnate, *circumference strigose with straight, diverging fibres of the same colour*. **Hymenium** thick, fleshy, even, *velvety with dense small bristles*.

Corticium. Among the most distinguished species of the genus, large, firm, never opening in flocculose clefts, rarely cracked.

On dead wood. Frequent. Aug.—Feb.

Spores ellipsoid, elongated, hyaline, 8 mk. *Q.* Name—*vellus*, fleece. Velvety. *Fr. Hym. Eur. p. 650. Berk. Out. p. 273. C. Hbk. n. 927. S. Mycol. Scot. n. 880.*

**12. *C. subdealbatum* B. & Br.**—Shining white, effused. **Hymenium** pallid fawn-colour with the bristles.

On fir. Badminton, 1866. Dec.

Name—*sub*, and *dealbatus*, whitened. *B. & Br. n. 1823.*

**13. *C. lactescens* Berk.**—*Flesh-colour, milky*, agglutinated, soft, waxy, undulated, margin shortly byssoid, at length cracked, interstices silky.

Smell like that of *Lactarius quietus*. Milk-white, watery. Hymenium flesh-colour or pale salmon-colour.

On dead wood, willow, &c. Frequent. Oct.—Feb.

Name—*lac*, milk. Milky. *Berk. Out. p. 274. C. Hbk. n. 932. S. Mycol. Scot. n. 881. Fr. Hym. Eur. p. 650.*

**14. *C. sanguineum* Fr.**—*Blood-red*, effused, adhering laxly, *like spider-web beneath*, circumference laxly fibrillose, *flesh-colour*. **Hymenium** even, smooth, *flesh-colour*.

The mycelium tinges the wood blood-red. Rarely gathered in a fertile state.

On dead pine and larch. Uncommon. Nov.—Feb.

Name—*sanguis*, blood. Of the colour. *Fr. Hym. Eur. p. 650. Berk. Out. p. 273. C. Hbk. n. 928. S. Mycol. Scot. n. 882.*

**15. *C. sulphureum* Fr.**—*Bright sulphur-yellow, effused*, fibriloso-flaxy. **Hymenium** (when perfect) thick, waxy-soft, cracked when dry.

The very variable mycelium is frequent, but it is rarely perfectly developed. It occurs with the hymenium tawny.

On dead wood. Frequent. Nov.—Dec.

Fries has gathered it in greatest perfection on rotting beech-leaves. Name—*sulphur*, brimstone. Of the colour. *Fr. Hym. Eur. p. 650. Berk. Out. p. 274. C. Hbk. n. 929. S. Mycol. Scot. n. 883.*

**16. *C. cinnamomeum* Fr.**—*Cinnamon*, effused, irregular by becoming confluent, adpressed, fibrilloso-strigose beneath and



at the circumference. **Hymenium** fleshy, soft, *smooth, naked*, Corticium. of the same colour, cracked when dry.

It varies *dark ochraceous*. When fertile it approaches *C. velutinus*.

On birch log. Glamis, 1874. Nov.

Name—*cinnamomum*, cinnamon. Of the colour. *Fr. Hym. Eur. p. 650. B. & Br. n. 1584. S. Mycol. Scot. n. 884.*

**17. C. cæruleum** Fr.—*Beautiful azure-blue*, somewhat round, then effused, adnate, at first tomentose, circumference flaxy and of the same colour inclining to be whitish. **Hymenium** waxy-soft, papillose, slightly bristly becoming smooth.

On rails, dead wood, fir, &c. Common.

Name—*cæruleum*, azure-blue. *Fr. Hym. Eur. p. 651. Berk. Out. p. 274. C. Hbk. n. 930. S. Mycol. Scot. n. 885.—Letell. Suppl. t. 630. Roth. Catal. 2. t. 9. f. 2. Sow. t. 350.*

**18. C. atro-virens** Fr.—*Black-green*, irregularly effused, tomentose beneath and at the circumference, of the same colour. **Hymenium** (found only as a barren cinereous pellicle).

On sticks in woods. Rare.

Name—*ater*, black; *vireo*, to be green. *Fr. Hym. Eur. p. 651. Berk. Out. p. 274. C. Hbk. n. 931.*

III.—LEIOSTROMA. Agglutinated (rarely when mature becoming free), &c.

\* *Waxy, when fertile facing the ground, &c.*

**19. C. calceum** Fr.—*White*, effused, agglutinated, waxy, *very smooth*, circumference similar. **Hymenium** even, smooth, when dry cracked, rigid.

Very polymorphous. The colour varies clay and whitish inclining to fuscous. Hymenium sometimes broken up into minute lumps.

On pine wood. Common. Oct.–Feb.

Name—*calx*, lime. Chalk-white. *Fr. Hym. Eur. p. 652. Berk. Out. p. 274. C. Hbk. n. 933. S. Mycol. Scot. n. 887.*

**20. C. puberum** Fr.—*White or clay-colour*, broadly effused, waxy, closely adnate, indeterminate. **Hymenium** even, *velvety with short bristles*, cracked when dry.

Like *C. calceum*, differing only in the hymenium being velvety with bristles. Very thin, separable from the matrix only in small fragments.

On dead wood. Aboyne. Forres. Glamis.

Corticium. Name—*puber*, adult. As applied to plants *downy*. From the hymenium. *Fr. Hym. Eur. p. 652. B. & Br. n. 1822. S. Mycol. Scot. n. 888. Theleph. Fr. Elench. p. 215.—Bonord. f. 256 (?)*

**21. C. scutellare** B. & Curt.—Whitish, then slightly tan-colour, resupinate, effused, immarginate. **Hymenium** broken up into minute areolæ.

On *Ulex*. Strachan, Kincardineshire. Glamis.

Name—*scutella*, a salver. From its form. *B. & Br. n. 1825. S. Mycol. Scot. n. 889. Grevillea, vol. ii. p. 4.*

**22. C. lividum** Pers.—*Changeable in colour*, effused, agglutinated, waxy-soft, becoming smooth, circumference similar. **Hymenium** even, *naked, somewhat viscid when moist*, cracked when dry.

In the same specimen the colour has been observed livid-azure-blue, and fuscous-purplish.

On dead birch. Rare.

Name—*lividus*, livid. *Pers. Obs. i. p. 38. Fr. Hym. Eur. p. 652. Berk. Out. p. 275. C. Hbk. n. 934. S. Mycol. Scot. n. 890.*

**23. C. ochraceum** Fr.—Broadly effused, 30 cent. (12 in.) and more, agglutinated, waxy-soft, smooth, circumference white, somewhat radiating, soon vanishing. **Hymenium** pallid then *ochraceous, sprinkled with golden-glistening atoms*, at length naked, papillose or tubercular, cracked when collapsed.

There is a remarkable variety with concentric papillæ.

On pine stumps. Uncommon. Nov.—Dec.

Name—*ochraceus*, ochrey-yellow. *Fr. Hym. Eur. p. 652. Berk. Out. p. 275. C. Hbk. n. 935. S. Mycol. Scot. n. 891. Theleph. Fr. Syst. Myc. i. p. 446.*

**\*\* *Amphigenous, becoming rigid, &c.***

**24. C. quercinum** Fr.—Cartilaginous-membranaceous, at first agglutinated, then adfixed in the centre, loosened on all sides, and at length involute, rigid, *smooth and becoming black beneath*. **Hymenium** continuous, flesh-colour.

It does not become cracked because it is set free in dying.

On oak branches. Very common. Aug.—April.

Name—*quercus*, oak. *Fr. Hym. Eur. p. 653. Berk. Out. p. 275. C. Hbk. n. 936. S. Mycol. Scot. n. 892.—Nees f. 253. Grev. t. 142. Bull. t. 436. f. 1.*

**25. C. cinereum** Fr.—Lurid, waxy becoming rigid, confluent, Corticium. agglutinated, circumference similar. **Hymenium** sprinkled with very thin *cinereous pruina*.

There are two forms, the thicker on wood, the thinner on bark.

On dead branches. Very common. Nov.—May.

Spores oblong or elongated, cylindrical, curved,  $3-5 \times 1$  mk. *K.* Name—*cinis*, ashes. Ash-colour. *Fr. Hym. Eur. p.* 654. *Berk. Out. p.* 275. *C. Hbk. n.* 937. *S. Mycol. Scot. n.* 893.

**26. C. incarnatum** Fr.—Waxy, becoming rigid, agglutinated, indeterminate, *circumference radiate*. **Hymenium** persistently *bright-coloured* (red, orange), sprinkled with very thin, *somewhat flesh-coloured pruina*.

The form on wood is rather thick, undulato-papillose, when collapsed even, cracked; that on bark thinner, persistently spuriously papillose from *Sphaeria* being incrusting by it.

On dead pine, Ulex, broom, &c. Very common. Aug.—March.

Spores ellipsoid or oblong-ellipsoid, often obsoletely curved,  $9-12 \times 5$  mk. *K.* Name—*incarnatus*, flesh-coloured. *Fr. Hym. Eur. p.* 654. *Berk. Out. p.* 275. *C. Hbk. n.* 938. *S. Mycol. Scot. n.* 894. *Fl. Dan. t.* 2035. *f.* 2.

**27. C. nudum** Fr.—*Flesh-colour becoming pale*, waxy, becoming rigid, agglutinated, *circumference determinate*, smooth. **Hymenium** even, cracked when dry, sprinkled (under a lens) with *whitish, fugacious pruina*.

On dead wood. Frequent. Nov.—Feb.

Name—*nudus*, naked. *Fr. Hym. Eur. p.* 655. *Berk. Out. p.* 276. *C. Hbk. n.* 939. *S. Mycol. Scot. n.* 895.

\* **C. citrinum** Pers.—Agreeing entirely with *C. nudum*, except in the hymenium being papillose, and in the colour being bright light yellow.

On dead wood. Moncreiffe.

Name—*citrus*, citron. Lemon-yellow. *Pers. Myc. Eur. i. p.* 136 (Theleph.) *Fr. Hym. Eur. p.* 655. *B. & Br. n.* 1695.

**28. C. confluens** Fr.—Somewhat membranaceous, *indeterminate*, agglutinated, *circumference radiate* (not fibrillose). **Hymenium** even, naked, hyaline, shining whitish when dry.

Circumference adnate, very delicate.

On dead branches, ash, &c. Frequent.

Name—*confluens*, to flow together. Confluent. *Fr. Hym. Eur. p.* 655. *Berk. Out. p.* 276. *C. Hbk. n.* 940. *S. Mycol. Scot. n.* 896.

Corticium

\*\*\* *Amphigenous, from the very first hardened, &c.*

**29. C. polygonium** Pers.—*Flesh-colour*, adnate, defined, soon hardened, somewhat grumous, circumference similar. **Hymenium** red *under the dense pruina*.

On dead branches, poplar, &c. Frequent.

Growing in little round detached patches from the ostiola of *Sphæria*. *M. J. B.* Name—*polygonius*, polygonal. From the shape of the tubercles which are owing to covered *Sphæria*. Pers. *Disp.* 30. *Fr. Hym. Eur.* p. 655. *Berk. Out.* p. 276. *C. Hbk.* n. 941. *S. Mycol. Scot.* n. 897.—*Hoffm. D. Fl.* 2. t. 6.

**30. C. violaceo-lividum** Fr.—*Violaceous-livid*, somewhat effused, adnate, hardened. **Hymenium** spuriously corrugated, tubercular, sprinkled with *thin, whitish pruina*.

Like *C. cinereum*, but allied to *C. polygonium*. Often growing on *Sphæria*.

On dead wood. Glamis, 1877.

Name—from the colours. *Fr. Hym. Eur.* p. 655. *B. & Br.* n. 1696. *S. Mycol. Scot.* n. 898.

**31. C. maculæforme** Fr.—*Somewhat rose-colour*, orbicular, then confluent, hardened, thin, circumference similar, smooth. **Hymenium** spuriously papillose, *bluish-grey pruinose*.

Commonly small, 2-8 mm. (1-4 lin.) broad.

On dead wood. Penzance.

Name—*macula*, a spot; *forma*, form. Spot-like. *Fr. Hym. Eur.* p. 656. *B. & Br.* n. 1975. *Theleph. Fr. Syst. Myc.* i. p. 454. *Fl. Dan.* t. 1738. f. 2.

**32. C. limitatum** Fr.—Lurid becoming pale, roundish, closely adnate, grumous-hardened, smooth, *circumference with a defined black edge*. **Hymenium** naked.

Easily distinguished by its essential character in having a black edge.

On *Cytisus*. Perth.

Name—*limitatus*, marked off. From the defined black circumference. *Fr. Hym. Eur.* p. 656. *B. & Br.* n. 1697. *S. Mycol. Scot.* n. 899.

**33. C. corrugatum** Fr.—*Pallid cinnamon*, somewhat effused, 5-7.5 cent. (2-3 in.) long, closely adnate, soon grumous. **Hymenium** beset with *ferruginous bristles*, very much cracked when dry.

The bristles are wholly like those of *Stereum rubiginosum*, to which it is otherwise scarcely allied. It has occurred somewhat waxy.

On dead wood. Very common. Aug.—Nov.

Name—*con*, and *ruga*, a wrinkle. Corrugated. *Fr. Hym. Eur. p. 656.* Corticium, Theleph. *Elench. p. 224.* Hymenochæte *Berk. Out. p. 272.* *C. Hbk. n. 918.* *S. Mycol. Scot. n. 867.*—*Grev. t. 234.*

\*\*\* *Amphigenous, very thin, innate, removing the bark.*

**34. C. comedens** Fr.—Flesh-colour becoming pale, effused, innate, growing under the bark, *exposed on the epidermis splitting asunder.* **Hymenium** even, *smooth*, cracked when dry.

Slightly viscid when moist. Erumpent specimens of *C. incarnatum* must not be confounded with it. Distinguished from all others by its manner of growth, never formed on decorticated wood or above the bark.

On dead branches. Very common. Nov.—March.

Hymenium sometimes white. The bark forms a margin round it. *M. J. B.* Name—*comedo*, to consume. From its destroying the bark under which it grows. *Fr. Hym. Eur. p. 656.* *Berk. Out. p. 276.* *C. Hbk. n. 942.* *S. Mycol. Scot. n. 900.* Theleph. *Nees Syst. f. 255.* *Fr. Elench. p. 219.*

\*\*\*\*\* *Species less known, doubtful.*

**35. C. aurora** Berk.—*Rose-colour*, turning pallid, very thin, effused, agglutinated, circumference indeterminate.

On dead leaves of *Carices*. Batheaston.

Name—*Aurora*, the dawn. From the colours. *Berk. Out. p. 276.* *C. Hbk. n. 944.* *Fr. Hym. Eur. p. 657.*

**36. C. typhæ** Fckl.—Longitudinally effused, thin, at first orbicular, white and flaxy, wholly smooth, then *somewhat mealy, tan-colour.*

On *Typha latifolia*. North Wooton, 1874.

Spores oblongo-lanceolate, multiguttate, hyaline. *Fckl.* Name—from the host plant. *Fuckel Symb. p. 27.* *Fr. Hym. Eur. p. 657.* *Grevillea, vol. iv. p. 119.*

*Subgenus* I.—CONIOPHORA. Fleshy, undulated and tubercular, &c.

**37. C. puteanum** Fr.—Light-yellowish-pallid, at length fuscous-olivaceous, the mucedinous circumference white, roundish or effused, *fleshy*, fragile. **Hymenium** somewhat undulated, pulverulent with the *fuscous-olivaceous spores.*

On dead wood in cellars, &c. Frequent. Nov.

Very fragile, 2 mm. (1 lin.) thick. It loosens itself from the trunk with age. It decays and grows black in winter, but in the following year a new plant crusts over the destroyed one. *Sommerf.* Sometimes dripping with moisture.

Corticium. *M. J. B.* Spores sphaeroid-ellipsoid, slightly fuscous,  $12-10 \times 8-9$  mk. *K.* Name—*puteum*, pertaining to a well. From its sometimes dripping. *Fr. Hym. Eur. p. 657. S. Mycol. Scot. n. 901. Theleph. Schum.—Berk. Out. p. 269. C. Hbk. n. 905. Fl. Dan. t. 2035.*

**38. C. stabulare** Fr.—Effused, flaxy when young, white, then fleshy-soft, *vinous-fuscous*, circumference flaxy, white. **Hymenium** tubercular, *white-pruinose*, the tubercles collapsing.

Odour foetid from its habitat.

On fir wood. Perth.

Name—*stabulum*, a stall. From its being found in an ox-stall. *Fr. Hym. Eur. p. 658. S. Mycol. Scot. n. 902.*

**39. C. umbrinum** Fr.—*Umber*, effused, fleshy-soft, *villous beneath*, circumference short, radiating, of the same colour. **Hymenium** tubercular then collapsing, *ferruginous-pulverulent*.

A distinguished species, not very broad but confluent, not easily separated from the matrix.

On dead wood. Penzance. Hothorpe. Feb.

Spores sphaeroid, punctate, 8 mk. *Q.* Name—from the *umber* colour, *Fr. Hym. Eur. p. 658. B. & Br. n. 1974. Grevillea, vol. xi. p. 14. Theleph. Alb. & Schw. p. 281. β. Fr. Elench. p. 199.*

**40. C. laxum** Fr.—About 5–7.5 cent. (2–3 in.) long, white, membranaceous, soft, adhering laxly, spider-web-tomentose beneath, circumference flaxy. **Hymenium** papillose, pallid then *ochraceous-ferruginous*, *ferruginous-pulverulent*.

Papillæ rather large.

On dead fir, lichens, moss, &c. Rare. Nov.

Spores ovoid-sphaeroid, punctate, spotted, 10 mk. *Q.* Name—*laxus*, loose. From its adhering laxly. *Fr. Hym. Eur. p. 659. S. Mycol. Scot. n. 903. Theleph. Fr. Elench. p. 190. Berk. Out. p. 269. C. Hbk. n. 906.*

**41. C. byssoideum** Fr.—*Ochrey-white*, irregularly effused, at first flaxy, *at length compact* at the disc, fleshy, pulverulent, *light yellow*, circumference flaxy, rather whitish.

Among fir-leaves, mosses, &c. Frequent.

Name—*bysusus*, flax. Flaxy. *Fr. Hym. Eur. p. 659. S. Mycol. Scot. n. 904. Theleph. Pers. Syn. p. 577. Berk. Out. p. 269. C. Hbk. n. 902.*

**42. C. aridum** Fr.—Membranaceous, effused, adnate, continuous, circumference whitish. **Hymenium** *even*, *sulphur-yellow*, then *setuloso-pulveraceous*, *umber inclining to ferruginous*.

On dead and decayed pine in woods and cellars. Common. Corticium. Nov.

Name—*aridus*, dry. *Fr. Hym. Eur.* p. 659. *S. Mycol. Scot. n.* 905. Theleph. *Berk. Out.* p. 269. *C. Hbk. n.* 907.

*Subgenus* II.—HYPOCHNUS. Breaking up and becoming floccose, &c.

\* *Breaking up and becoming floccose.*

**43. C. serum** Fr.—*White*, broadly effused, incrusting, thin, fleshy when fresh, *smooth, pruinose*, then and when dried *splitting open into flocci*, papillæ round, crowded together, equal.

A very singular species. Papillæ regular.

On trunks. Epping. Glamis. Menmuir. Winter.

Spores sphaeroid or ovoideo-ellipsoid, smooth, hyaline or chlorino-hyaline,  $9-11 \times 7-8$  mk. *K.* Name—*serus*, late. Growing in winter. *Fr. Hym. Eur.* p. 659. *B. & Br. n.* 1583, 1697. *S. Mycol. Scot. n.* 906. Theleph. *Pers. Syn. p.* 580. *Fr. Elench. p.* 211.

**44. C. sambuci** Fr.—*White*, effused, somewhat innate, *incrusting, surrounding branches, indeterminate*, continuous when in vigour, breaking into floccules when dry.

Easily recognised from its habitat.

On dried branches of elder. Very common. Aug.—Dec.

Very difficult to separate from *C. calceum* by a strict definition. *M. J. B.* Name—*sambucus*, elder. *Fr. Hym. Eur.* p. 660. *Berk. Out. p.* 276. *C. Hbk. n.* 943. *S. Mycol. Scot. n.* 907.—*Grev. t.* 242. *Letell. t.* 607. f. 2.

\*\* *Hymenium tomentose, somewhat pulverulent.*

**45. C. olivaceum** Fr.—Membranaceous, adnate, circumference fringed and *whitish*. **Hymenium** thin, *dark olivaceous*, tomentose with small bristles.

Broadly effused, orbicular or elongated. The colour is distinguishing and remarkably constant.

On pine wood. Uncommon.

Spores sphaeroid or sphaeroid-ellipsoid, very pale yellowish or slightly fuscous, 4 mk. *K.*; ovoideo-ellipsoid, nucleate, yellow, 8 mk. *Q.* Name—from the colour. *Fr. Hym. Eur.* p. 660. Theleph. *Fr. Elench. p.* 197. *Berk. Out. p.* 269. *C. Hbk. n.* 908.

**46. C. ferrugineum** Fr.—*Ferruginous*, effused, adnate, everywhere *tomentose*. **Hymenium** papillose, pulveraceous.

It may be compared with *Hydnum ferruginosum* and *Polyporus floccosus*.

Corticium. On branches. Altyre, &c. Oct.

Name—*ferrugo*, iron-rust. From the rusty colour. *Fr. Hym. Eur. p. 661. B. & Br. n. 1585. S. Mycol. Scot. n. 908.*

**47. C. lacunosum** B. & Br.—Broadly effused, soft, mycelium woolly-tawny, lacunose. **Hymenium pulverulent.**

On branches. Aboyne, 1870. Sept.

Name—*lacuno*, to hollow out. Full of hollows, lacunose. *B. & Br. n. 1371. S. Mycol. Scot. n. 909. Fr. Hym. Eur. p. 661.*

**48. C. anthochroum** Fr.—*Brick-red-rose-colour*, becoming pale, effused, even above, wholly broken up, *floccoso-velvety*, circumference flaxy, paler.

Very pretty. Without a hymeneal membrane.

On sticks, sycamore, birch. Rare. Feb.

Name—*ἄνθος*, a flower; *χρῶς*, colour. *Fr. Hym. Eur. p. 661. S. Mycol. Scot. n. 910. Theleph. Pers. Syn. p. 576. Berk. Out. p. 270. C. Hbk. n. 909.*

Cyphella.

# GENUS XLIX.—**Cyphella** (κύφος, a goblet).

Fr. Syst. Myc. 2. p. 201.

Somewhat membranaceous, cup-shaped, rarely plane, adnate behind, commonly extended in stem-like form, pendulous. Hymenium definitely inferior, similar, without the heterogeneous disc and asci of *Pezizæ*, even or at length slightly wrinkled. Sporophores 4-spored, spores subovate.

Without doubt very near to *Soleniæ*, and removed from them by artificial characters. *Fr. Hym. Eur. p. 661.*



LXXXIX. *Cyphella catilla*.  
Natural size.

- \* *Lignicolæ* (growing on wood).
- \*\* *Muscicolæ* (growing on mosses).
- \*\*\* *Stipiticolæ* (growing on stems).
- \*\*\*\* *Caulicolæ* and *epiphyllæ* (growing on stalks and leaves).

\* *Lignicolæ* (growing on wood).

**1. C. Bloxami** B. & Phill.—White, floccoso-membranaceous, disc becoming light yellow, crenato-lobed; flocci even.



Sporophores turbinate. Spores 8-11 mk. Spores terminating slightly Cyphella. branched threads.

On *Ulex*. Twycross.

Name—after Rev. A. Bloxam. *B. & Br. n.* 1894.

**2. C. griseo-pallida** Weinm.—*Pallid-grey*, somewhat membranaceous, globose then campanulate, sessile, *floccose externally*. **Hymenium** even, smooth.

On dead *Carex paniculata*. Spye Park, Wiltshire.

Whole plant  $\frac{1}{2}$  lin. diam., at first granuliform. Hymenium pale reddish-grey, border slightly undulated. *M. J. B.* Name—from the colour. Weinm. *Ros. p.* 522. *Fr. Hym. Eur. p.* 662. *Berk. Out. p.* 277. *C. Hbk. n.* 945.

**3. C. ochroleuca** B. & Br.—Ochrey-white and villous above, membranaceous, cup-shaped, *margin at length split*. **Hymenium** even, pale ochre, brighter than the pileus.

2 mm. (1 lin.) and more broad.

On bramble sticks. Rare.

Name—*ὀχρὸς*, pale yellow; *λευκός*, white. *B. & Br. n.* 719. *Berk. Out. p.* 277. *C. Hbk. n.* 948. *S. Mycol. Scot. n.* 911. *Fr. Hym. Eur. p.* 662.

**4. C. stuppea** B. & Br.—Erumpent, sessile, pezizæform, externally coarsely hispid, brownish then becoming white. **Hymenium** fuscous.

On broom. Menmuir, Forfarshire. March.

Name—*stuppeus*, made of tow. From being coarsely hispid. *B. & Br. n.* 1698. *S. Mycol. Scot. n.* 912.

**5. C. fulva** B. & Rav.—*Brown-tawny*, membranaceous, cup-shaped, deflexed, externally *tomentose*.

On dead bark.

Remarkable for its deflexed mouth. Spores ovate, 17 mk. *B. & Br.* Name—*fulvus*, tawny. *B. & Br. n.* 936. *C. Hbk. n.* 954. *Fr. Hym. Eur. p.* 662.

**6. C. brunnea** Phill.—Dirty-brown, sessile, cupulate, clothed near the margin with a grey pruina, margin incurved, lacerated, mouth oblique. **Hymenium** smooth, discoloured-brown; flesh paler, subgelatinous.

Scattered or crowded. Cups .5 mm. across, .8 mm. high. Basidia clavate, two to four spicules; spores colourless, globose (5-6 mk.)

On bark and wood of old elder-trees. Shrewsbury.

Cyphella.

Name—*brunneus*, brown. *Phill. Grevillea*, vol. xiii. p. 49.

7. **C. fraxinicola** B. & Br.—Minute, orbicular, externally snow-white, shortly villous; disc light yellow, becoming fuscous with the spores, proliferous.

On ash. Batheaston, 1873. Dec.

Name—*fraxinus*, ash-tree; *colo*, to inhabit. *B. & Br. n.* 1446.

\*\* *Muscicolæ* (growing on mosses).

8. **C. muscigena** Fr.—Very small, 6–10 mm. (3–5 lin.) broad, *shining white*, membranaceous-soft, *becoming plane*, somewhat dimidiate, externally *silky under a lens*. **Hymenium** slightly wrinkled.

A variable form, also stipitate, spathulate, &c.

On moss. Uncommon. Sept.–Dec.

At first flabelliform. *M. f. B.* Name—*muscus*, moss; *gigno*, to bear. Growing on moss. *Fr. Hym. Eur. p.* 663. *Berk. Out. p.* 277. *C. Hbk. n.* 946. *S. Mycol. Scot. n.* 914. *Theleph. Pers. Myc. Eur. t.* 7. f. 6. *Canth. lævis Fr. Syst. Myc. i. p.* 324.

9. **C. galeata** Fr.—Whitish, membranaceous-soft, somewhat sessile, *obversely cup-shaped, then dimidiate, helmet-shaped*, even, margin quite entire. **Hymenium** at length rufescent, slightly wrinkled.

When young entire, cup-shaped, grey when moist, snow-white when dry, then rufescent.

On mosses. Uncommon.

Name—*galea*, a helmet. Helmet-shaped. *Fr. Hym. Eur. p.* 663. *Berk. Out. p.* 277. *C. Hbk. n.* 947.—*Fl. Dan. t.* 2027. f. 1.

10. **C. catilla** Smith.—Somewhat membranaceous, expanded, margin crisped and undulated. **Hymenium** veined, 18 mm. ( $\frac{3}{4}$  in.) broad, grey.

Often imbricated. Allied to *C. galeata*.

On moss and dead leaves. King's Lynn.

Name—*catillus*, a small bowl. From its shape. *Smith Seem. Journ. Bot.* 1873, p. 337.

11. **C. muscicola** Fr.—About 4 mm. (2 lin.) broad, pale-whitish, membranaceous, somewhat sessile, persistently *cup-shaped*, nodding, externally *slightly fibrilloso-striate, margin slightly downy, repand, torn*. **Hymenium** even.

B. cup irregular, externally cinereous,

On mosses. Rare.

Cyphella.

Name—*muscus*, moss; *colo*, to inhabit. *Fr. Hym. Eur. p.* 663. *Syst. Myc.* ii. *p.* 202. *Berk. Out. p.* 277. *C. Hbk. n.* 949.—*B. Fl. Dan. t.* 2083. *f.* 2.

\*\*\* *Stipiticolæ* (growing on stems).

**12. C. lacera** Fr.—When perfect 4–6 mm. (2–3 lin.) in stature, whitish or yellow, membranaceous, cup-shaped, stipitate from the vertex being extended, pendulous, *then torn into many clefts, slightly striate above with dense black fibrils.* **Hymenium** slightly wrinkled, whitish.

On dead twigs. Rare.

Name—*lacerus*, torn. *Fr. Hym. Eur. p.* 664. *Syst. Myc.* ii. *p.* 202. *Berk. Out. p.* 277. *C. Hbk. n.* 950.—*Alb. & Schw. t. i. f.* 5.

**13. C. capula** Fr.—*Whitish*, membranaceous, obliquely campanulate, extended into an oblique stem, *smooth, margin sinuate*, irregularly shaped. **Hymenium** even.

It varies yellowish.

On dead stems of herbaceous plants. Frequent.

Looks like *Peziza*. *M. J. B.* Spores ovoid, 7 mk. *Q.* Name—*capulæ*, small bowls with handles. *Fr. Hym. Eur. p.* 664. *Berk. Out. p.* 278. *C. Hbk. n.* 951. *S. Mycol. Scot. n.* 915. *Holmsk. ii. t.* 22. *Peziza Nov. Act. Hafn. i. p.* 286. *f.* 7.—*Fl. Dan. t.* 1970. *f.* 3 seems a var.

**14. C. Pimii** Phill.—White or very pale yellow, cup-shaped, erect or pendent, membranaceous, pubescent, margin of cup somewhat incised. **Stem** rather slender, crooked, enlarged upwards.

Fasciculate. About 4 mm. (2 lin.) high and cup 2 mm. (1 lin.) wide. Basidia cylindraceo-clavate, with sometimes only two spicules, generally four; spores subpyriform, colourless (7–10 × 4 mk.)

On dead herbaceous stems in water. Dublin.

Name—after Greenwood Pim. *Phill. Grevillea, vol. xiii. p.* 49.

\*\*\*\* *Caulicolæ and epiphyllæ* (growing on stalks and on leaves).

**15. C. pallida** B. & Br.—Cups at first orbicular, *at length irregularly lobed*, plane, tomentose or slightly hispid, *sessile.* **Hymenium** at length wrinkled, pallid ochraceous.

Cups  $\frac{1}{4}$ –1 line across, sometimes proliferous. Differs from *C. Curreyi* in the colour of the hymenium, which is rugose, like that of *Cantharellus muscigenus*, and its more irregular form. It appears also not to be erumpent, as that species often is, but is seated on the bark or wood.

Cyphella.

On old stems of *Clematis vitalba*.

Spores elliptic, 6-9 mk. *B. & Br.* Name—*pallidus*, pallid. *B. & Br. n.* 1372. *Rabenh. exs. n.* 1415. *Fr. Hym. Eur. p.* 664.

**16. *C. cuticulosa*** Berk.—*White*, diaphanous, membranaceous, at first oblong, then cup-shaped, elongated into a stem, *smooth* externally.

On dead grass stems. Very rare. Oct.—Feb.

Not recorded since found by Dickson. Name—*cuticula*, the thin external skin. From its membranaceous character. *Berk. Out. p.* 278. *C. Hbk. n.* 955. *Fr. Hym. Eur. p.* 665.—*Dicks. Crypt. iii. t. 9. f.* 11.

**17. *C. Goldbachii*** Weinm.—*White*, membranaceous, cup-shaped, sessile, *pitcher-shaped-concave, lobed*, externally *villous*. **Hymenium** even, pallid.

Minute, irregular and lobed, in which it differs from the *Soleniæ*.

On dead leaves of *Aira cæspitosa*. Spye Park, Wilts.

Near to *C. cuticulosa*, from which it differs in its villous coat. *B. & Br.* Name—after Goldbach. *Weinm. Ross. p.* 522. *Fr. Hym. Eur. p.* 665. *Berk. Out. p.* 278. *C. Hbk. n.* 952.—*Corda Sturm t.* 63.

**18. *C. dochmiospora*** B. & Br.—Snow-white, minute, pezizæ-form.

Resembles externally *Peziza villosa*; but the hairs are not granulated.

Habitat not recorded. Batheaston, 1864. Oct.

Spores oblique, ovate, rather acute, 14-17 mk. *B. & Br.* Name—δόχμιος, oblique; σπορά, seed. *B. & Br. n.* 1373. *S. Mycol. Scot. n.* 916.

## ORDER V.—CLAVARIEI.

Hymenium not distinct from the hymenophore, amphigenous. Somewhat fleshy fungi, vertical, simple or branched. *Never coriaceous or incrusting as the Merismoid Thelephoræ which are similar in form.* *Fr. Hym. Eur. p. 666.*

GENUS L.—**Sparassis** (σπαρσσω, to tear in pieces).  
Fr. Syst. Myc. p. 464.

Sparassis.


Fleshy, branched, with flat, leaf-like branches, composed of two plates, fertile on both sides, with 4-spored sporophores. *Very handsome.* *Fr. Hym. Eur. p. 666.*

**1. S. crispa** Fr.—Whitish, very much branched; *branches 2.5–5 cent. (1–2 in.) broad, intricate, recurved at the apex, zoneless, serrated.*

Very handsome, very large (tufts have occurred as much as 1–2 ft. broad), most frequently wholly sessile, more rarely shortly stipitate.

Beside fir trees and stumps. Rare.  
Aug.–Oct.

Most frequently it has a tinge of pale yellow. Like a large sponge. Edible and delicious; but too rare to be of value as an article of food. Name—*crispa*, a curl. Crisped. *Fr. Hym. Eur. p. 666. Syst. Myc. i. p. 465. Sv. atl. Sv. t. 17. B. & Br. n. 1139\*. C. Hbk. n. 958. S. Mycol. Scot. n. 917. Kl. Bor. t. 463. Hogg & Johnst. t. 24. Bail. t. 27.—Wulf. in Jacqu. Misc. t. 14. f. 1. Krombh. t. 22. f. 2, 3. Schæff. t. 163.*




XC. *Sparassis crispa*. One-fifth natural size.

Clavaria.

GENUS LI.—**Clavaria** (*clava*, a club). Linn.—Fr. Syst. Myc. i. p. 465.

Fleshy, branched or simple, somewhat round, without a distinct stem. Hymenium continuous, dry, homogeneous. *For the most part growing on the ground. Fr. Hym. Eur. p. 666.*



XCI. *Clavaria cinerea*. One-third natural size.

I. **RAMARIA** (*ramus*, a branch). Holmsk. Branched, branches attenuated upwards.

A. *Leucosporæ* (λευκός, white; σπόρος, seed). Spores white or pallid.

\* Colour bright, yellow, red or violet. On the ground.

\*\* Colour white or grey. On the ground.

\*\*\* Colour whitish or dingy. On wood.

B. *Ochrosporæ* (ὀχρός, pale yellow; σπόρος, seed). Spores ochraceous or cinnamon.—There are analogous species in each section which are so similar in form and colour that their figures are with difficulty distinguished. Yet they are most distinct; the *Ochrosporæ* spongy, tough, the majority of them bitter, and not fit to be eaten.

\* Colour yellow, somewhat cinnamon, saffron. On the ground.

\*\* Colour whitish, grey, violet. On the ground.

\*\*\* Growing on trunks.

II. **SYNCORYNE** (σύν, together; κορύνη, a club). Somewhat simple, cæspitose at the base or fasciculate.

\* Reddish.

\*\* Yellowish or white.

\*\*\* Fuliginous or blackish.

III. **HOLOCORYNE** (ὅλος, entire; κορύνη, a club). Somewhat simple, distinct at the base.

\* Colour changeable, becoming dark.

\*\* Colour unchangeable, commonly shining white.

I.—**RAMARIA**. Branched, branches attenuated upwards.

A. *Leucosporæ*. Spores white or pallid.

\* Colour bright, yellow, red or violet.

1. **C. botrytes** Pers.—Varying white, yellow, flesh-colour, 7.5 cent. (3 in.) and more high, but also less, fragile; trunk very thick, 2.5–5 cent. (1–2 in.), unequal, very much branched. Branches swollen, unequal, somewhat wrinkled, red at the apices.

About beech stumps. Inverary, &c. Rare. Aug.—Oct.

Clavaria.

Spores ellipsoid, subhyaline,  $12-15 \times 6$  mk. *K.* Edible. Name—*βότρυς*, a cluster of grapes. From the form. *Pers. Syn.* p. 587. *Fr. Hym. Eur.* p. 667. *Syst. Myc.* i. p. 466. *Sv. atl. Sv. t.* 35. *Berk. Out.* p. 278. *C. Hbk.* n. 959. *S. Mycol. Scot.* n. 918. *Fl. Dan.* t. 1303. *Barla t.* 40. f. 1-3. *Krombh. t.* 53. f. 1-3. *Harz. t.* 67. *Corda Ic. v.* f. 75. *Quel. t.* 21. f. 4.—*Schæff. t.* 176. *Wulf. in Jacqu. Coll.* ii. t. 13. *Holmsk. ii.* p. 117 with fig. *Krombh. t.* 53. f. 4 (white).

**2. *C. amethystina*** Bull.—*Violet, fragile*, very much branched, equal at the base. **Branches** round, *even*, obtuse.

In grassy woods and pastures. Uncommon. Aug.—Oct.

Very variable in size. Sometimes 7.5 cent. (3 in.) or more high and very much branched, sometimes a few lines and nearly simple. *M. J. B.* Edible. Name—*ἀμέθυστος*, amethyst. From the colour. *Bull. t.* 496. f. 2. *Fr. Hym. Eur.* p. 667. *Berk. Out.* p. 279. t. 18. f. 2 small var. *C. Hbk. n.* 960. *S. Mycol. Scot.* n. 919.—*Holmsk. Ot. i.* p. 110 with fig., borrowed in *Nees Syst.* f. 151.

**3. *C. fastigiata*** Linn.—Scarcely exceeding 2.5 cent. (1 in.) long, egg-yellow, slightly tough, cæspitose, very much branched, equal. **Branches** *short, divaricate*, branchlets fastigiate.

It varies with the apices light yellow and fuscous.

In pastures. Very common. Aug.—Oct.

Spores pale buff, irregular, 7 mk. *W.G.S.*; subsphæroid, 6 mk. *K.* Name—*fastigium*, the top. Reaching the top. *Linn. Suec. n.* 1269. *Fr. Hym. Eur.* p. 667. *Berk. Out.* p. 279. *C. Hbk. n.* 961. *S. Mycol. Scot.* n. 920. *Bull. t.* 358. f. D, E. *Holmsk. i.* p. 90 with fig. *Fl. Dan.* t. 836. f. 2. *C. pratensis Syst. Myc.* i. p. 471. *Pers. Comm. t.* 4. f. 5. *Ray Syn. t.* 8. f. 4.

**4. *C. muscoides*** Linn.—*Yellow*, slightly tough, becoming slender, twice or thrice forked, stem thin. **Branchlets** *crenate-shaped*, acute.

In pastures. Frequent. Aug.—Oct.

Taller than *C. fastigiata*, solitary, less branched, dry, very smooth, except the base which is tomentose. *M. J. B.* Name—*muscus*, moss; *εἶδος*, appearance. Like a tuft of moss. *Linn. Suec. n.* 1270. *Fr. Hym. Eur.* p. 667. *Berk. Out.* p. 279. *C. Hbk. n.* 962. *S. Mycol. Scot.* n. 921. *Fl. Dan.* t. 775. f. 2. *Bull. t.* 496. f. O, Z. *Holmsk. i.* p. 89 with fig.—*Schæff. t.* 173. *Krombh. t.* 53. f. 22, 23. *Vaill. Par. t.* 8. f. 4.

**5. *C. curta*** Fr.—*Greenish-yellow*, very small, very much branched, pressed close together. Stem none. **Branches** *short, crowded*, obtuse.

Nearest to *C. fastigiata*, but differing in stature and colour, never egg-yellow as in that species.

On the ground. Coed Coch. Holme Lacy.

Clavaria.

Spores white. *Fr.* Name—*curtus*, curt, short. *Fr. Hym. Eur.* p. 668. *Monogr.* ii. p. 281. *B. & Br.* n. 1448.

**\*\* Colour white or grey.**

**6. C. coralloides** Linn.—*White*, somewhat fragile, *hollow within, trunk* rather thick, repeatedly and *irregularly much branched*. **Branches** unequal, dilated upwards, branchlets crowded, acute.

Commonly cæspitose, growing into each other, unequally fastigiate. It varies with the branchlets not developed, obtuse.

In shady woods. Rare. Aug.–Oct.

Name—κοράλλιον, coral; εἶδος, appearance. Coral-like. *Linn. Suec.* n. 1268. *Fr. Hym. Eur.* p. 668. *Sv. atl. Sv.* t. 92. f. 4, 5. *Berk. Out.* p. 279. *C. Hbk.* n. 963. *S. Mycol. Scot.* n. 922. *Holmsk.* i. p. 113 with fig. *Sow.* t. 278. *Krombh.* t. 53. f. 4.

**7. C. cinerea** Bull.—*Cinereous*, fragile, stuffed, *trunk somewhat thick*, short, very much branched. **Branches** and branchlets *thickened*, irregularly shaped, somewhat wrinkled, rather obtuse.

The trunk varies thicker and thinner. It differs from *C. grisea* in the spores.

In mixed woods. Frequent. Aug.–Oct.

Spores irregular, ellipsoid-sphæroid, 8–10 × 6 mk. *K.* Name—*cinis*, ashes. Ash-coloured. *Bull.* t. 354. *Fr. Hym. Eur.* p. 668. *Berk. Out.* p. 279. *C. Hbk.* n. 965. *S. Mycol. Scot.* n. 923. *Grev.* t. 64, 321. *Letell.* t. 708. f. 1.—*Krombh.* t. 53. f. 9, 10.

**8. C. umbrina** Berk.—*Pale umber*, slightly branched. **Branches** and branchlets *cylindrical*, obtuse, forked.

The habit is that of *C. fastigiata*. It has not, however, the slightest tinge of yellow.

On lawns. Coed Coch. Blair Athole. Aug.–Oct.

Name—from the umber colour. *Berk. Out.* p. 279. t. 18. f. 4. *C. Hbk.* n. 964. *S. Mycol. Scot.* n. 924. *Fr. Hym. Eur.* p. 668.

**9. C. cristata** Pers.—*White*, very variable in stature, 2.5–5 cent. (1–2 in.) high, tough, even, stuffed. **Branches** *dilated above, acutely incised*, crested.

Somewhat cæspitose.

In mixed woods. Frequent. Aug.–Oct.

Spores angulato- or ellipsoid-sphæroid, hyaline, 8 mk. *K.* Name—*crista*, a crest. Crested. *Pers. Syn.* p. 591. *Fr. Hym. Eur.* p. 668. *Syst. Myc.* i. p. 473. *Sv. atl. Sv.* t. 92. f. 1–3. *Berk. Out.* p. 280. *C. Hbk.* n. 966. *S.*



*Mycol. Scot. n. 925. Fl. Dan. t. 1304. f. 2. Grev. t. 190. Krombh. t. 53. Clavaria. f. 12, 13.—Holmsk. i. p. 92 with fig. Schæff. t. 170. Pers. Comm. t. 2. f. 4.*

**10. C. rugosa** Bull.—White, tough, simple or sparingly branched, *thickened upwards, wrinkled. Branches* irregularly shaped, obtuse.

Very variable in form, sometimes wholly simple, club-shaped, sometimes divided into 2-4 variously unequal branches.

In woods and pastures. Very common. Aug.—Nov.

Spores angulato-sphæroid, 8-10 mk. *K.* Name—*ru-ga*, a wrinkle. *Bull. t. 448. f. 2. Fr. Hym. Eur. p. 669. Syst. Myc. i. p. 473. Berk. Out. p. 280. t. 13. f. 3. C. Hbk. n. 967. S. Mycol. Scot. n. 926. Fl. Dan. t. 1301. Grev. t. 328.—Schæff. t. 291. Bolt. t. 115. Sow. t. 278 lower fig. Vaill. Par. t. 8. f. 2. Quel. t. 20. f. 5* slender, even, furnished with short irregular branches from the base, perhaps a distinct species.

**11. C. Krombholzii** Fr.—Shining white, *fragile*, cæspitose, even, sparingly branched. *Branches somewhat compressed*, obtuse.

Like *C. rugosa*, but fragile and even. Varying in stature.

In old pastures. Glamis. Moncreiffe. Sept.

Name—after Krombholz. *Fr. Hym. Eur. p. 669. B. & Br. n. 1586. S. Mycol. Scot. n. 927.—Krombh. t. 53. f. 15, 16, 18-20. Bull. t. 496. f. 3.*

**12. C. Kunzei** Fr.—Shining white, tufts 4-5 cent. (1½-2 in.) high, 2.5 cent. (1 in.) broad, somewhat fragile, *very much branched from the thin, cæspitose base. Branches* elongated, crowded, repeatedly forked, *fastigiate*, even, equal, axils compressed.

In woods. Rare. Sept.

Name—after Kunze. *Fr. Hym. Eur. p. 669. Syst. Myc. i. p. 474. Berk. Out. p. 280. C. Hbk. n. 968. S. Mycol. Scot. n. 928.—Bull. t. 358. f. 1.*

\*\*\* *Colour whitish or dingy.*

**13. C. pyxidata** Pers.—Pallid then tan-colour, somewhat rufescent, tufts as much as 10-12.5 cent. (4-5 in.) high, trunk thin, smooth, branched. *Branches and branchlets all hollowed out in cup-shape at the apex*, cups radiate in a proliferous manner at the margin.

The most distinct of the genus, *white-spored*. Branchlets obconic and even, the cups again and again proliferous in a whorled manner, the terminal ones toothed.

On the ground. Penzance.

Name—*pyxis*, a small box. *Pyxidate. Pers. Comm. t. 1. f. 1. Fr. Hym. Eur. p. 669. B. & Br. n. 1976. Fl. Dan. t. 1304. f. 1.*

Clavaria.

B. *Ochrospora*. Spores ochraceous or cinnamon.

\* Colour yellow, somewhat cinnamon, saffron.

14. **C. aurea** Schæff.—Trunk thick, elastic, pale, *divided into stout*, tense and straight, very dichotomously branched, round, obtuse, somewhat toothed, *yellow branches*.

Large, 7.5–10 cent. (3–4 in.) high, elastic, soft, not pure yellow.

In mixed woods. Rare. Aug.–Oct.

Spores ellipsoid, 10–14 × 5–6 mk. *K.* Name—*aurum*, gold. Yellow. *Schæff. t.* 287 & 285. *Fr. Hym. Eur. p.* 670. *Berk. Out. p.* 280. *C. Hbk. n.* 969. *S. Mycol. Scot. n.* 929.—*Bull. t.* 222. *Krombh. t.* 53. f. 7.

15. **C. formosa** Pers.—Trunk thick, 2.5 cent. (1 in.) and more, whitish, elastic. **Branches** very much branched, *elongated, orange-rose-colour, branchlets obtuse, light yellowish*.

Variable.

In woods. Rare.

Spores elongated oval, covered with papillæ, 16 × 8 mk. *W.G.S.* Name—*formosus*, finely formed, beautiful. *Pers. Ic. & descr. t.* 3. f. 5. *Fr. Hym. Eur. p.* 671. *Syst. Myc. i. p.* 466. *B. & Br. n.* 1031. *C. Hbk. n.* 970. *Krombh. t.* 54. f. 21, 22. *Holmsk. i. n.* 13 with fig. *Corda Ic. iii. f.* 136.—*Harz. t.* 7 lower fig. *Batsch f.* 48.

16. **C. spinulosa** Pers.—Trunk short, rather thick (2.5 cent., 1 in.), pallid. **Branches** elongated, crowded, tense and straight, *attenuated, somewhat cinnamon* (fuliginous-date-brown: Pers.), of the same colour at the apex.

More distinct than *C. aurea* and *formosa*.

In pine woods. Coed Coch.

Name—*spinulosus*, spined. From the appearance of the branches. *Pers. Obs. 2. t.* 3. f. 1. *Fr. Hym. Eur. p.* 671. *Syst. Myc. i. p.* 468. *B. & Br. n.* 1302.—*Krombh. t.* 53. f. 8.

17. **C. abietina** Pers.—Height 5–7.5 cent. (2–3 in.) *Ochraceous, trunk* short, rather thick (8 mm., 4 lin.), *white-tomentose*, very much branched. **Branches** crowded, longitudinally *wrinkled* when dry, branchlets tense and straight, acute.

Becoming green when rubbed. Taste bitter.

Under spruce fir. Common. Aug.–Oct.

Spores oval, greenish-yellow, 5 × 4 mk. *W.G.S.*; subellipsoid, 8–10 × 4–6 mk. *K.* Name—*abies*, fir. *Pers. Comm. p.* 46 (var.) *Fr. Hym. Eur. p.* 671. *Syst. Myc. i. p.* 469. *Berk. Out. p.* 280. *C. Hbk. n.* 971. *S. Mycol. Scot. n.* 930. *Grev. t.* 117. *Fl. Dan. t.* 2030. f. 2.

**18. C. flaccida** Fr.—*Ochraceous*, 2.5–4 cent. (1–1½ in.) long, *Clavaria*. thin, very much branched, *flaccid*. Stem *thin*, 2 mm. (1 lin.) thick, very short, smooth. **Branches** crowded, even, unequal, converging, acute.

Allied to *C. abietina*, but much thinner. It is smooth in itself, but the mycelium is floccose, creeping over leaves.

Among moss in wood. Rare.

Spores ellipsoid-sphaeroid,  $4.5 \times 3$  mk. *K*. Name—*flaccidus*, flaccid. *Fr. Hym. Eur. p. 671. Berk. Out. p. 280. C. Hbk. n. 972.*

**19. C. crocea** Pers.—*Saffron-yellow*, minute, thin, stem naked, pallid. **Branches** and branchlets similar, somewhat forked.

According to the fig. it is wholly distinct. Spores unknown.

On waste ground. Wraxall, Somerset. Glamis. Sept.

Name—*crocus*, saffron. Saffron-yellow. *Pers. Ic. & descr. t. 11. f. 6. Berk. Out. p. 280. C. Hbk. n. 973. S. Mycol. Scot. n. 931.*

**\*\* Colour whitish, grey, violet.**

**20. C. grisea** Pers.—Firm. Trunk 4 cent. (1½ in.) long, 2.5 cent. (1 in.) thick, whitish. **Branches** 7.5 cent. (3 in.) long, attenuated, *somewhat wrinkled*, and as well as the unequal, obtuse branchlets, *fuliginous-cinereous*.

In its exceedingly variable form it is with difficulty distinguished from *C. cinerea*, but is clearly distinct from its rufous-brown spores.

In woods. Rare. Sept.

Name—*griseus*, grey. *Pers. Comm. p. 44. Fr. Hym. Eur. p. 672. Syst. Myc. i. p. 468. Berk. Out. p. 281. C. Hbk. n. 974. S. Mycol. Scot. n. 932.*

**21. C. condensata** Fr.—Tan-rufescent, tufts very dense, 7.5–10 cent. (3–4 in.) high, and without a common trunk, *very much branched from the base, smooth*, unchangeable when bruised. **Branches** *tense and straight, crowded in a parallel manner*, even, fastigate at the apex, twice or thrice toothed, yellow.

Mycelium white, membranaceous, creeping among leaves.

On the ground under trees. West Farleigh, 1874.

Spores ellipsoid, 9 mk. *K*. Name—*condenso*, to press close together. From the manner in which the branches are close together. *Fr. Hym. Eur. p. 672. B. & Br. n. 1587.—Schæff. t. 177* (departing in its reddish colour).

**\*\*\* Growing on trunks.**

**22. C. stricta** Pers.—Yellowish-pallid, *becoming fuscous when*

Clavaria.

*bruised*, trunk about 6 mm. (3 lin.) long and *thick*, very much branched. **Branches** and branchlets tense and straight, even, crowded and adpressed, acute.

Mycelium most frequently fibrillose, naked.

In woods. Uncommon. Aug.—Oct.

Spores dark cinnamon. *Fr.*; creamy-yellow,  $4 \times 6$  mk. *W.G.S.* Name—*stringo*, to draw tight. Tense and straight. *Pers. Comm. p.* 45. *t.* 4. *f.* 1. *Fr. Hym. Eur. p.* 673. *Berk. Out. p.* 281. *t.* 18. *f.* 5. *C. Hbk. n.* 975. *S. Mycol. Scot. n.* 933. *Fl. Dan. t.* 1302. *f.* 1. *Krombh. t.* 54. *f.* 23.

**23. C. crispula** *Fr.*—Entire height 2.5–7.5 cent. (1–3 in.) Tan then ochraceous, very much branched, trunk thin, with villous rootlets. **Branches** *flexuous*, multifid, *branchlets* of the same colour, *divaricate*.

Protean but quite distinct.

At the base of trees, ash. Rare.

Spores creamy-yellow,  $3 \times 5$  mk. *W.G.S.* Name—*crispulus*, slightly curled. *Fr. Hym. Eur. p.* 673. *Syst. Myc. i. p.* 470. *Berk. Out. p.* 281. *C. Hbk. n.* 976.—*Bull. t.* 358. *f.* 1. *a, b.* *Ehrenb. Nov. Act. Nat. Cur. x. t.* 14.

II.—SYNCORYNE. Somewhat simple, cæspitose at the base, &c.

\* *Reddish.*

**24. C. purpurea** *Fr.*—*Purple*, cæspitose. **Clubs** elongated, generally 7.5 cent. (3 in.) long, hollow then compressed, simple, acute.

Very remarkable. When solitary it is more drawn out and decumbent. Fragile.

In grassy places. Rare. Aug.—Oct.

Name—from the colour. *Fr. Hym. Eur. p.* 674. *Syst. Myc. i. p.* 480. *Berk. Out. p.* 281. *C. Hbk. n.* 977. *S. Mycol. Scot. n.* 934. *Fl. Dan. t.* 837. *f.* 2.

**25. C. rufa** *Fl. Dan.*—*Rufous*, 4 cent. ( $1\frac{1}{2}$  in.) long, 4 mm. (2 lin.) thick. **Clubs** stuffed, thickened, *somewhat bifid*, acute.

Cæspitose.

On the ground. Rannoch. The Burn, Forfarshire.

Name—*rufus*, reddish. *Fl. Dan. t.* 775. *f.* 1. *Fr. Hym. Eur. p.* 674. *Syst. Myc. i. p.* 480. *S. Mycol. Scot. n.* 935.

**26. C. rosea** *Fr.*—*Rose-colour*, 4 cent. ( $1\frac{1}{2}$  in.) long, somewhat fasciculate, fragile. **Clubs** stuffed, at length becoming yellow at the apex, attenuated downwards, whitish.

Elegant. No distinct stem, but the club is narrowed at the base.

Clavaria.

In pastures and among moss. Rare.

Name—from the colour. *Fr. Hym. Eur. p. 674. Syst. Myc. i. p. 482. Berk. Out. p. 281. C. Hbk. n. 978. S. Mycol. Scot. n. 936.—Sv. Bot. t. 558. B. attenuata Fr. Obs. 2. t. 5. f. 2. Krombh. t. 53. f. 21.*

**\*\* Yellowish or white.**

**27. C. fusiformis** Sow.—*Yellow*, cæspitoso-connate, slightly firm, *soon hollow*. **Clubs** somewhat fusiform, simple and toothed, even, attenuated to the base which is of the same colour.

Taller than *C. inæqualis*, more connate, at length becoming fuscous at the apex. Both vary unequal, not very different.

In woods, pastures, &c. Common.

Name—*fusus*, a spindle; *forma*, form. Spindle-shaped. *Sow. t. 234. Fr. Hym. Eur. p. 674. Berk. Out. p. 281. C. Hbk. n. 979. S. Mycol. Scot. n. 937. Bolt. t. 110. Var. with the irregularly shaped clubs toothed: Sow. t. 235.*

**28. C. ceranoides** Pers.—Height 7.5 cent. (3 in.) *Yellow*, fasciculate, unequal, slightly divided above, apex brown.

Very cæspitose. Clubs attenuated downwards, curved, somewhat ventricose.

In woods. Rare. Nov.

Name—*κερός*, wax; *εἶδος*, appearance. Waxy-like. *Pers. Syn. p. 594. Berk. Out. p. 282. C. Hbk. n. 980. Fr. Syst. Myc. i. p. 481. Hym. Eur. p. 674* under *C. fusiformis*.

**29. C. inæqualis** Fl. Dan.—*Yellow*, gregarious, somewhat fasciculate, fragile, *stuffed*. **Clubs** variable, simple or forked, continuous *downwards, of the same colour*.

Very changeable in form, clavate and cylindrical. Distinguished from *C. fusiformis* chiefly by its irregularity. Tufts less dense.

In woods and pastures. Very common. Aug.–Oct.

2.5–7.5 cent. (1–3 in.) high. Compressed, angular or channelled, often bifid and variously cut and jagged at the apex, more or less ventricose in the centre. *Grev. Spores white, 11 × 5 mk. W.G.S. Name—inæqualis, unequal. Fl. Dan. t. 873. f. 1. Fr. Hym. Eur. p. 674. Syst. Myc. i. p. 481. Berk. Out. p. 282. C. Hbk. n. 981. S. Mycol. Scot. n. 938. Fl. Bat. t. 814. f. 1. Hussey i. t. 18.—Bull. t. 264. Sow. t. 253 partly. Pers. Comm. t. 1. f. 3. Grev. t. 37. Fl. Dan. t. 1783. f. 2. Sv. Bot. t. 504. f. 4.*

**Var. aurantiaca** Pers. Somewhat wrinkled and compressed. *Krombh. t. 53. f. 19, 20. Vaill. t. 7. f. 5.*

**30. C. argillacea** Fr.—Height 1–2.5 cent. ( $\frac{1}{2}$ –1 in.) *Pallid-clay-colour*, fasciculate, fragile. **Clubs** simple, variable. Stem *yellow-shining*.


## Clavaria.

Constantly thickened upwards.

In woods and moss. Frequent. Aug.-Oct.

There is a variety with a white stem. The plant, moreover, is either dilated or cylindrical. The stem in this species is more distinct from the pileus than is consistent with the generic character. *M. J. B.* Name—*argilla*, clay. Clay-coloured. *A. clavata*, clubs somewhat compressed. *Fr. Hym. Eur. p. 675. Syst. Myc. i. p. 482. Berk. Out. p. 282. C. Hbk. n. 982. S. Mycol. Scot. n. 939. Fr. Obs. t. 5. f. 3. Schmied. Ic. t. 15. Fl. Dan. t. 1852. f. 2. Fl. Bat. t. 814. f. 2. Harz. t. 7. f. 1-7. B. Clubs cylindrical, attenuated. Pers. Comm. t. 1. f. 4. Fl. Dan. t. 1966. f. 2.*

**31. C. vermicularis** Scop.—*Shining white*, cæspitose, fragile. **Clubs** stuffed, simple, *cylindrical, awl-shaped*.



*XCII. Clavaria vermicularis.*  
One-half natural size.

Often flexuous or incurved. Of the thickness of a pigeon's quill.

On lawns. Very common. Sept.

Looks like a little bundle of candles. *M. J. B.* Name—*vermis*, a worm. Flexuous. *Scop.—Fr. Hym. Eur. p. 675. Syst. Myc. i. p. 484. Berk. Out. p. 282. C. Hbk. n. 984. S. Mycol. Scot. n. 940. Swartz in Vet. Ak. Handl. 1811. t. 6. f. 2. Fl. Dan. t. 1966. f. 1, 775. f. 2. Mich. t. 87. f. 2.*

**32. C. fragilis** Holmsk.—Commonly white, sometimes yellowish, fasciculate, *very fragile*, attenuated and *white downwards*. **Clubs** hollow, rather obtuse, variable.

Distinguished by its *white base*, &c., from the three preceding species. 2.5-7.5 cent. (1-3 in.) long, occasionally forked, when young stuffed, round, tense and straight, then hollow, compressed, twisted, often wrinkled.

In pastures, &c. Common. Aug.-Oct.

Name—*fragilis*, fragile. *Holmsk. i. p. 7 with fig. Fr. Hym. Eur. p. 675. Syst. Myc. i. p. 484. Berk. Out. p. 283. C. Hbk. n. 985. S. Mycol. Scot. n. 941. A. club swollen, thickened. Bull. t. 463. f. 1. Sow. t. 90. Mich. t. 87. f. 10. Vaill. t. 7. f. 5. B. more slender, cylindrical, somewhat attenuated. Sow. t. 232. Fl. Dan. t. 775. f. 2. Barla t. 41. f. 14-16. Mich. t. 87. f. 13. C. Cl. gracilis Pers. thin, gregarious, acute. Bolt. t. 111. f. 1. Mich. t. 87. f. 6, perhaps a distinct species.*

\*\*\* *Fuliginous or blackish.*

**33. C. fumosa** Pers.—*Fuliginous*, fasciculate, fistulose, *fragile*. **Clubs** even, tense and straight, somewhat compressed.

It differs from *C. fragilis* only in the colour.

In pastures, &c., uncommon. Aug.-Sept.

Name—*fumus*, smoke. Smoke-coloured. *Pers. Comm. p.* 76. *Fr. Hym. Clavaria. Eur. p.* 676. *B. & Br. n.* 1303. *C. Hbk. n.* 986. *S. Mycol. Scot. n.* 942. *Krombh. t.* 53. *f.* 18.

III.—HOLOCORYNE. Somewhat simple, separate at the base.

\* *Colour changeable, becoming dark.*

**34. C. pistillaris** Linn.—Light yellow then rufescent, simple, large, *fleshy, stuffed, obovato-clavate*, obtuse.

In woods among fern. Uncommon. Sept.

15-30 cent. (6-12 in.) long. Dingy brown in decay. *M. f. B.* Spores white, 10 x 5 mk. *W. G. S.*; ellipsoid, hyaline, 10-11 x 5-6 mk. *K.* Name—*pistillum*, a pestle. Pestle-shaped. *Linn. Suec. n.* 1246. *Fr. Hym. Eur. p.* 676. *Berk. Out. p.* 283. *C. Hbk. n.* 987. *S. Mycol. Scot. n.* 943. *Bull. t.* 244. *Sow. t.* 277. *Hussey i. t.* 62. *Fl. Dan. t.* 1255. *Holmsk. p.* 12 with fig. *Sv. Bot. t.* 504. *f.* 1. *Krombh. t.* 54. *f.* 1-11. *Ventur. t.* 41. *f.* 1-2. *Corda Sturm. ii. t.* 58 and *Icon. v. f.* 76. *Quel. t.* 21. *f.* 2. Forms departing from the type: *Schæff. t.* 290. *Kl. Bor. t.* 395, slender, attenuated upwards. —*Pers. Myc. Eur. t.* 15. *f.* 1 veined at the apex; perhaps distinct.—*Chev. Par. t.* 8. *f.* 2. Var. *alba* *Batt. t.* 3. *A.*

**35. C. ligula** Fr.—Yellowish when young, pallid-rufescent when full grown, simple, gregarious, *spongy-fleshy*, elongato-clavate, obtuse, *villous at the base*.

Smaller and softer than *C. pistillaris*.

In woods. Strachan, Kincardineshire.

Name—*ligula*, a small tongue, strap. From its shape. *Fr. Hym. Eur. p.* 676. *B. & Br. n.* 1820. *S. Mycol. Scot. n.* 944. *A.* young, *Schmied. Ic. t.* 5 upper fig. *Fl. Dan. t.* 837.—*Wulf. in Jacqu. Misc. ii. t.* 2. *f.* 2. *Sv. Bot. t.* 504. *f.* 3. *B.* old,—*Schæff. t.* 171. *Krombh. t.* 54. *f.* 12.

**36. C. contorta** Holmsk.—Watery yellowish, 2.5 cent. (1 in.) long, about 6-8 mm. (3-4 lin.) thick, simple, *erumpent*, stuffed, spongy-fleshy, soft to the touch, *somewhat twisted, wrinkled* obtuse, *pruinose*.

Among the most distinguished, growing in winter. Almost spongy after rain. Sometimes spatulate, compressed, sometimes swollen and somewhat twisted. More or less irregularly curved. Solitary or cæspitose.

On fallen branches. Rare. Sept.

Easily known by its erumpent habit. Name—*contorqueo*, to twist. Twisted. *Holmsk. Ot. i. p.* 29 with fig. *Fr. Hym. Eur. p.* 677. *Syst. Myc. i. p.* 478. *Berk. Out. p.* 283. *C. Hbk. n.* 988. *S. Mycol. Scot. n.* 945.—*Fl. Dan. t.* 1852.

**37. C. fistulosa** Fr.—Light yellowish then rufescent, simple, slender, very long, *tense and straight, fistulose*, rather obtuse; root short, *villous*.

## Clavaria.

Too nearly allied to *C. juncea*. *Fr. Hym. Eur. p. 677. Fl. Dan. t. 1256, 1100. f. 3. Holmsk. i. p. 15 with figure. Krombh. t. 5. f. 19.*

\* **C. Ardenia** Sow.—Ferruginous then date brown, simple, very long, more than 20 cent. (8 in.), thickened upwards, *hollow, obtuse or hollowed out* at the apex, tomentose at the base, not rooting.

The apex is acute when young, then obtuse, &c.

On fallen branches. Rare.

Var. *Ardenia* only is British. Name—after Lady Arden. *Sow. t. 215. Fr. Hym. Eur. p. 677. Syst. Myc. i. p. 478. Berk. Out. p. 283. C. Hbk. n. 989. S. Mycol. Scot. n. 946.*

**38. C. juncea** Fr.—Pallid then rufescent, 5–12.5 cent. (2–5 in.) long, thin, *filiform, flaccid, fistulose*, acute, *the creeping base fibrillose*.

Gregarious. Its affinity with *Typhulæ* is very close. The root is never sclerotoid at the base, stem not distinct. Var. *vivipara*, fibrillose at the sides. *Bull. t. 463. f. H.*

In woods among leaves. Uncommon. Sept.

Name—*juncus*, a rush. *Fr. Hym. Eur. p. 677. Syst. Myc. i. p. 479. Berk. Out. p. 283. C. Hbk. n. 990. S. Mycol. Scot. n. 947.—Fl. Dan. t. 1257. Mich. t. 87. f. 7.*

**39. C. tuberosa** Sow.—*Yellowish*, cæspitose, simple, tough, even, linear, tuberous and rooting at the base.

On birch. Very rare. Dunphail, Morayshire. Nov.

Root a thick, strigose, subglobose tuber, giving out two or three simple linear subacute receptacles from the same base. *M. F. B.* Not recorded since the time of Sowerby till gathered at Dunphail by Dr Keith. Spores ellipsoid, comma-shaped, punctate, 10 mk. *Q.* Name—*tuber*, a tuber. Tuberous-rooted. *Sow. t. 199. B. & Br. n. 1448. S. Mycol. Scot. n. 948. Calocera Fr. Hym. Eur. p. 680. Berk. Out. p. 284. C. Hbk. n. 994.*

\*\* *Colour unchangeable, commonly shining white.*

**40. C. canaliculata** Fr.—*Shining white*, 7.5 cent. (3 in.) long, solitary, fistulose, slightly tough, very smooth, not rooting, at length *compressed, channelled* or splitting longitudinally.

Sometimes in pairs.

On the ground. Coed Coch.

Name—*canaliculus*, a small channel. Marked with channels. *Fr. Hym. Eur. p. 678. B. & Br. n. 1895. Quel. t. 21. f. 1.—Bull. t. 496. f. L. M.*

**41. C. tenuipes** B. & Br.—Pallid *clay-colour*. **Club inflated**, wrinkled. **Stem** slender, flexuous, somewhat distinct from the club.



Not exceeding 2.5 cent. (1 in.) in height. Head swollen, obovate, about 12 Clavaria. mm. ( $\frac{1}{2}$  in.) high, rarely confluent with the stem.

On bare heathy ground. Sherwood Forest. Nov.

Name—*tenuis*, thin; *pes*, a foot. Thin-stemmed. *B. & Br. n.* 369. *t.* 9. *f.* 2. *Out. p.* 282. *C. Hbk. n.* 983. *Fr. Hym. Eur. p.* 678.

**42. C. incarnata** Weinm.—*Flesh-colour, white-pruinose*, internally purple, varying 12 mm.—4 cent. ( $\frac{1}{2}$ –1 $\frac{1}{2}$  in.) long, quite simple, solid, cylindrical.

Gregarious.

On the ground. Terrington, Norfolk.

Name—*incarnatus*, flesh-coloured. *Weinm. p.* 510. *Fr. Hym. Eur. p.* 678. *Grevillea, vol.* viii. *p.* 98.

**43. C. acuta** Sow.—White, 2.5–7.5 cent. (1–3 in.) high, quite simple, tense and straight. **Club** small, distinct, sharp-pointed, *pruinose*. **Stem** cylindrical, equal.

On soil in garden pots. Rare.

Name—*acutus*, acute. Sharp-pointed. *Sow. t.* 333. *Berk. Out. p.* 283. *C. Hbk. n.* 991. *Fr. Hym. Eur. p.* 679. *Syst. Myc. i.* *p.* 485.

**44. C. uncialis** Grev.—*Pale white*, quite simple, stuffed, *tough, tense and straight, obtuse*, smooth. **Stem** continuous, attenuated.

Varying in size, but scarcely exceeding 2.5 cent. (1 in.)

On dead stems of herbaceous plants, &c. Uncommon. May.

Name—*uncia*, an inch. *Grev. t.* 98. *Fr. Hym. Eur. p.* 679. *Elench. p.* 232. *Berk. Out. p.* 284. *C. Hbk. n.* 992. *S. Mycol. Scot. n.* 949.

GENUS LII.—**Calocera** (καλός, beautiful; κέρασ, a horn).

Calocera.

*Fr. Syst. Myc. i.* *p.* 485.

Gelatinoso-cartilaginous, horny when dry, vertical, somewhat cylindrical, simple or branched, viscid, without a distinct stem. Hymenium amphigenous; sporophores bifurcate; spores oblong, curved.

Intermediate between the *Clavariæ* and *Tremellini*. *Fr. Hym. Eur. p.* 679.

\* *Branched.*


\*\* *Cæspitose.*

\*\*\* *Simple, distinct.*

Calocera.

\* *Branched.*

1. **C. viscosa** Fr.—*Golden-egg-yellow*, 2.5 cent. (1 in.) and more high, branched, tough, *rooted*, even, linear. **Branches** round or compressed, tense and straight, repeatedly dichotomous.



The long root paler.

On stumps, chiefly pine. Very common. July-Dec.

Spores white. *Fr.*; hyaline, ellipsoid oblong, mostly slightly curved, 8-10 × 4 mk. *K.* Name—*viscus*, bird-lime. Viscous. *Fr. Hym. Eur. p.* 680. *Syst. Myc. i. p.* 486. *Berk. Out. p.* 284. *C. Hbk. n.* 993. *S. Mycol. Scot. n.* 950. *Bonord. f.* 257.—*Pers. Comm. t. i. f.* 5. *Schæff. t.* 174. *Quel. t. 21. f.* 5.

XCIH. *Calocera viscosa*. One-half natural size.

2. **C. palmata** Fr.—*Orange-yellow*, branched, tremelloso-tough, compressed, *dilated upwards*, divided.

**Branchlets** somewhat round, divaricate, obtuse.

Nearest to *C. cornea*.

On wood. Twycross.

Name—*palma*, the palm. From the branchlets resembling a palmate leaf. *Fr. Hym. Eur. p.* 680. *Grevillea, vol. v. p.* 11.

\*\* *Cæspitose.*

3. **C. cornea** Fr.—*Yellow-orange*, 4-10 mm. (2-5 lin.) and more high, cæspitose, rooted, even, viscus. **Clubs** small, *curt*, awl-shaped, connate at the base.

On dead wood, oak and beech. Frequent.

Name—*cornu*, a horn. Horny. *Fr. Hym. Eur. p.* 680. *Syst. Myc. i. p.* 486. *Berk. Out. p.* 284. *C. Hbk. n.* 995. *S. Mycol. Scot. n.* 951.—*Batsch f.* 161. *Fl. Dan. t.* 1305. *f. 2.* *Sow. t.* 40. *Bull. t.* 463. *f. 4.*

4. **C. corticalis** Fr.—*Pallid-flesh-colour*, cæspitose, erumpent, soft, pellucid. **Clubs** small, awl-shaped, somewhat distinct.

Elegant, delicate, smaller than the rest, scarcely 2 mm. (1 lin.) long.

On dead bark. Penzance.

Name—*cortex*, bark. Growing on bark. *Fr. Hym. Eur. p.* 680. *Elench. p.* 233. *B. & Br. n.* 1977.—*Batsch f.* 162?

\*\*\* *Simple, distinct.*

Calocera.

**5. *C. stricta*** Fr.—Yellow, simple, solitary, elongated, blunt at the base, linear, even when dry.

*A. truncorum*, 12 mm.—2.5 cent. ( $\frac{1}{2}$ –1 in.) long, the base girt with white delicate down. *B. epiphylla*, 5–7.5 cent. (2–3 in.) high, the base naked and bluntly rooted.

On ash. Belvoir Castle. Oct.

Name—*stringo*, to draw tight. Tense and straight. *Fr. Hym. Eur. p. 680.* *A. B. & Br. n. 1032.* *C. Hbk. n. 996.*—*Bonord. f. 255.*

**6. *C. striata*** Fr.—Yellow, simple, solitary, tough, *lanceolate*, acute, *striate* when dry.

Commonly taken for a form of *C. cornea*, but distinct.

On prostrate trunks. Batheaston.

Spores 8×6 mk. *B. & Br.* Name—*stria*, a furrow. Striate. *Fr. Hym. Eur. p. 681.* *B. & Br. n. 1140.* *C. Hbk. n. 997.*—*Hoffm. Germ. t. 7. f. 1.*

**7. *C. glossoides*** Fr.—Light-yellow, 12 mm. ( $\frac{1}{2}$  in.) long, simple, solitary, somewhat tremellose. **Club** small, *thickened*, obtuse, compressed. Stem round.

On decayed oak and stumps. Leighwood, Bristol.

Composed of erect forked flocci. Spores oblong, oblique, somewhat incurved, attached by a short pedicel. *B. & Br.* Name—*γλῶσσα*, the tongue; *εἶδος*, appearance. Tongue-like. *Fr. Hym. Eur. p. 681.* *Syst. Myc. i. p. 487.* *Berk. Out. p. 284.* *C. Hbk. n. 998.*

GENUS LIII.—**Pterula.** Fr. in Linn. 1830. p. 531.

Pterula.

Cartilaginous, filiform, arid, equal, without a distinct stem. Hymenium pubescent then becoming smooth, with 4-spored sporophores (Léveill.) *Fr. Hym. Eur. p. 681.*

Some species are simple, others much branched.

**1. *P. subulata*** Fr.—Height 4 cent. ( $1\frac{1}{2}$  in.), very densely crowded, tense and straight, equal except at the attenuated base, sparingly branched. **Branches** growing into each other, *whitish-cinereous*, *multifid at the apex*, awl-shaped, smooth, becoming yellow.

Tough. Not thicker than a fine thread.

On wood. Burnham Beeches, 1874.

Name—*subula*, an awl. Awl-shaped. *Fr. Hym. Eur. p. 682.* *Syst. Myc. iii. p. 286.* *Linn. 1830. t. 11. f. 4.* *B. & Br. n. 1588.*

Pterula.

**2. P. multifida** Fr.—Height 2.5–5 cent. (1–2 in.), at the first pallid whitish then (especially when dried), dirty pale yellowish, *very much branched*, very delicate, flaccid but slightly tough. **Branches** tense and straight, not much thicker than a hair, heaped as if swept together, somewhat fastigate, *spear-shaped at the apex*, of the same colour.

On dead branches. Sept.

Name—*multus*, many; *findo*, to cleave. From the multitude of branches into which it is cleft. *Fr. Hym. Eur. p. 682. Monogr. ii. p. 282. B. & Br. n. 1304.*

Typhula.

GENUS LIV.—**Typhula** (*Typha*, the Reed-mace, which it somewhat resembles in miniature. *Berk.*) Pers.—Fr. *Epicr. p. 584.*

Minute tender fungi with a filiform stem, which is either heterogeneous and distinct from the linear club, or springs from a sclerotoid hybernaculum. Hymenium waxy, sporophores forked, spicules elongated. *Growing on plants. Fr. Hym. Eur. p. 682.*

I. PHACORRHIZÆ (φακός, a lentil; ῥίζα, a root, from the sclerotoid root). Springing from sclerotoid hybernacula.

II. LEPTORRHIZÆ (λεπτός, slender; ῥίζα, a root). *Radical tubercle (Sclerotium) not discovered or awaiting.*


I.—PHACORRHIZÆ. Springing from sclerotoid hybernacula.

**1. T. erythropus** Fr.—Simple, club 4–6 mm. (2–3 lin.) long, *white*, linear, cylindrical, *smooth*; stem elongated, 12 mm. ( $\frac{1}{2}$  in.), filiform, *reddish-black*, hybernaculum depressed, blackish.

The hybernaculum (sometimes awaiting) is *Sclerotium scutellatum*.

On dead stems of herbaceous plants. Common. Autumn.

Name—ἐρυθρός, red; πούς, a foot. Red-stemmed. *Fr. Hym. Eur. p. 683. Syst. Myc. i. p. 495. Berk. Out. p. 284. C. Hbk. n. 999. S. Mycol. Scot. n. 952. Fl. Dan. t. 2030.—Bolt. t. 112. Grev. t. 43.*



XCIV. *Typhula phacorrhiza*.  
One-half natural size.

**2. T. phacorrhiza** Fr.—*Pallid, be-*

coming somewhat fuscous below, simple, elongated, *filiform*, Typhula. smooth; hybernaculum compressed, pallid then fuscous.

The hybernaculum is *Sclerotium scutellatum*. By most recent authors it has been joined with *Clavaria juncea*, but the best authorities keep them distinct. Cl. juncea Leveill. in *Ann. Sc. Nat.* 1843, t. 7. f. 1 seems a var. of this.

On dead herbaceous stems, leaves, &c. Uncommon. Oct.

Name—*φακός*, a lentil; *ρίζα*, a root. From the sclerotoid hybernaculum. *Fr. Hym. Eur.* p. 683. *Berk. Out.* p. 284. *C. Hbk. n.* 1002. *S. Mycol. Scot. n.* 953. *Schnitzl. Sturm* 31. t. 12. *Reich. in Schr. Naturf. Freund. Berl.* i. t. 9. f. 4. *Sow. t.* 233.

**3. T. incarnata** Lasch.—Whitish, *flesh-colour upwards*, simple, filiform, cylindrical, attenuated into a continuous stem and *hairy downwards*; hybernaculum compressed, fuscous.

Very much allied to *T. phacorrhiza*.

On dead herbaceous stems. Uncommon.

Name—*incarnatus*, flesh-coloured. *Lasch.—Fr. Hym. Eur.* p. 683. *Berk. Out.* p. 285. *C. Hbk. n.* 1001. *S. Mycol. Scot. n.* 954.—*Grev. t.* 93.

**4. T. gyrans** Fr.—*White*, simple, very slender, somewhat cylindrical; stem thin, *pubescent*; hybernaculum pallid, becoming fuscous.

The hybernaculum is *Sclerotium complanatum*.

On straw, &c. Uncommon.

Name—*gyro*, to turn in a circle or make circular. From the somewhat cylindrical head. *Fr. Hym. Eur.* p. 684. *Br. Bath Trans.* 1870, p. 92. *C. Hbk. n.* 1000. *S. Mycol. Scot. n.* 955.—*Batsch f.* 164.

**5. T. muscicola** Fr.—Simple, filiform, club *white*, cylindrical, obtuse, *attenuated into a thin, smooth stem*; hybernaculum even, pallid.

Distinctly stipitate, 5–7.5 cent. (2–3 in.) high.

On the larger mosses. Uncommon.

Name—*muscus*, moss; *colo*, to inhabit. *Fr. Hym. Eur.* p. 684. *Berk. Out.* p. 285. *C. Hbk. n.* 1003.—*Pers. Obs.* 2. t. 3. f. 2.

II.—LEPTORRHIZÆ.—Radical tubercle not discovered, &c.

**6. T. Grevillei** Fr.—*White*, simple, club thickened, obtuse; stem *capillary, pilose*.

Doubtless a form of *T. gyrans*, wanting the tubercle.

Typhula.

On dead leaves, &amp;c. Frequent. Oct.

Name—after Greville. *Fr. Hym. Eur. p. 685. Berk. Out. p. 285. C. Hbk. n. 1004. S. Mycol. Scot. n. 956. Grev. t. 49.*

**7. T. gracillima** White.—White; stem very slender, curved, smooth, club elongated.

On various herbaceous plants. Perth.

Name—*gracilis*, slender. Very slender. *White.—B. & Br. n. 1699. S. Mycol. Scot. n. 957.*

**8. T. filiformis** Fr.—Club whitish, thickened, smooth; *stem somewhat branched, decumbent, date-brown.*

Among dead leaves. Rare.

Name—*filum*, a thread; *forma*, form. Thread-like. *Fr. Hym. Eur. p. 685. Berk. Out. p. 285. C. Hbk. n. 1005.—Bull. t. 448. f. 1. Sow. t. 387. f. 4.*

**9. T. gracilis** Berk. & Desm.—Club pallid, *simple or forked*, acute, rough with spores and little prominent bristles; stem short, distinct, smooth or bristly.

2 mm. (1 lin.) or more high. Rugged with the fruit-bearing cells, which are frosted with the spores, and interspersed sometimes with short bristles, of which some of the upper ones support a small spore; tips often acuminate and then nearly barren. Very near to *Isaria*.

On putrid leaves. Uncommon.

Spores elliptic, having a sparkling appearance under a lens. *B. & Br. ; sphaeroid or sphaeroid-ellipsoid, 8–10 mk. K. Name—gracilis, slender. Berk. Out. p. 285. B. & Br. n. 84. t. 8. f. 1. C. Hbk. n. 1006. Fr. Hym. Eur. p. 686.*

**10. T. translucens** B. & Br.—Minute, white, pellucid; stem short, thickened upwards; head irregular, somewhat obovate.


Resembling somewhat a prematurely dried *Myxogast*, but a true *Hymenomyce*.

On the ground. Glamis, 1875. Oct.

Name—*transluceo*, to be transparent. Translucent. *B. & Br. n. 1589. S. Mycol. Scot. n. 958.*

GENUS LV.—**Pistillaria** (*pistillum*, a pestle. From the form). *Pistillaria*.  
Fr. Syst. Myc. i. p. 496.

Very small, with a turgid club, which is either sessile, or attenuated into a short continuous stem, hardened when dry. Substance somewhat cellular. Sporophores simple or forked. *Fr. Hym. Eur. p. 686.*




1. **P. micans** Fr.—*Glistening-rose-colour*, scarcely exceeding 2 mm. (1 lin.) high, obovate, obtuse; stem short, attenuated, and, as well as the oval spores, white.

Smooth, but somewhat pruinose.

On dead thistles. Rare.

Spores ovoid-ellipsoid,  $9-10 \times 5-6$  mk. *K.* Name—*mico*, to glisten. *Fr. Hym. Eur. p. 686. Berk. Out. p. 285. C. Hbk. n. 1007.—Hoffm. Germ. t. 7. f. 2. Ehrenb. Ber. t. 3. f. 2.*



XCV. *Pistillaria puberula*.  
Natural size. Section four times natural size.

2. **P. culmigena** Fr.—*Hyaline-pellucid*, ovato-clavate, obtuse; stem distinct, very short.

Small, soft when fresh, hardened when dry, sclerotoid.

On stalks of grass. Frequent. Jan.

Name—*culmus*, a haulm; *gigno*, to bear. Growing on stalks. *Fr. Hym. Eur. p. 687. Montagn. Ann. Sc. Nat. 1836, t. 12. f. 2. Berk. Out. p. 285. C. Hbk. n. 1008.*

3. **P. quisquiliaris** Fr.—*Whitish*, 6–8 mm. (3–4 lin.) long, thickened upwards, *somewhat compressed*, soft when fresh, rigid when dry, attenuated *at the base*, *smooth*, somewhat stipitate.

Gregarious, here and there flattened and bifid.

On fern stems. Common. Sept.–Oct.

Often attached to a *Sclerotium*. *M. F. B.* Spores sausage-shaped, 15 mk. *Q.* Name—*quisquiliæ*, sweepings, rubbish. Among refuse of fern. *Fr. Hym. Eur. p. 687. Berk. Out. p. 286. C. Hbk. n. 1009. S. Mycol. Scot. n. 959.—Sow. t. 334 f. 1.*

4. **P. furcata** Smith.—Clubs white or yellowish, 3 cent. ( $1\frac{1}{4}$  in.) high, waxy then tough, compressed, broad at the apex, attenuated downwards, generally furcate and cæspitose.

Pistillaria.

In greenhouses.

Name—*furca*, a fork. Forked. *Worth. Smith in C. Hbk. n. 1012.*

5. **P. puberula** Berk.—White, scarcely 2 mm. (1 lin.) high, obovate, ventricose; stem short, distinct, attenuated upwards, pellucid, *fibrous*, tomentose.

On dead *Pteris aquilina*. Rare. Sept.—Oct.

Name—*pubes*, first downy hair on the chin. From the tomentose stem. *Berk. Out. p. 286. Eng. Fl. v. p. 181. C. Hbk. n. 1010. S. Mycol. Scot. n. 960. Fr. Hym. Eur. p. 688.—Sow. t. 334, f. 2.*

6. **P. pusilla** Fr.—*White*, very small, 2 mm. (1 lin.) long, smooth, even, *linear*; stem not distinct.

On *Equisetum*. Weymouth.

Slightly thickened upwards, nodding when dry. *M. J. B.* Name—*pusio*, a small boy. Very small. *Fr. Hym. Eur. p. 688. Syst. Myc. i. p. 498. Berk. Out. p. 286. Eng. Fl. v. p. 182. C. Hbk. n. 1011.*

Microcera.

GENUS LVI.—**Microcera** (μικρός, small; κέρα, a horn).

Desm. Ann. Sc. Nat. 1848, p. 350.

“External veil persistent, membranaceous-floccose, breaking above into many laciniae. Receptacle fleshy, club-shaped, formed of simple spore-bearing fibres. Spores fusiform, arcuate.” *Fr. Hym. Eur. p. 689.*

No British species.



## ORDER VI.—TREMELLINEI.

Whole fungus homogeneous, gelatinous, shrivelling when dry, reviving when moistened, pervaded internally with branched filaments, terminating towards the surface all round in sporophores; spores somewhat reniform.

The structure of the Tremellini is admirably illustrated by Tulasne in *Ann. Sc. Nat.* 1853. *Fr. Hym. Eur.* p. 689.

GENUS LVII.—**Tremella** (*tremo*, to tremble).

Tremella.

Dill.—*Fr. Syst. Myc.* 2. p. 210.

Distended with jelly when moist, tremulous, immarginate, *not papillate*, sporophores globose, becoming quadripartite, and sending out from each division an elongated free spicule, which terminates in a simple spore. *Fr. Hym. Eur.* p. 689.

I. MESENTERIFORMES (*μεσέντερον*, the mesentery). Gelatinous inclining to cartilaginous, foliaceous, naked.

II. CEREBRINÆ (*cerebrum*, the brain). Firm then pulpy, somewhat pruinose with the spores.

III. CRUSTACEÆ (*crusta*, a crust). Diffused, becoming plane.

IV. TUBERCULIFORMES (*tuberculum*, a little tuber). Small, somewhat erumpent.



XCVI. *Tremella mesenterica*.  
Natural size.

I.—MESENTERIFORMES. Gelatinous, inclining to cartilaginous, &c.

1. **T. fimbriata** Pers.—Olivaceous inclining to black, cæspitose, clusters 5–7.5 cent. (2–3 in.) high and even broader, *erect, corrugated*; lobes flaccid, incised at the margin, *undulato-fimbriate*.

When soaked with water it has a dark tawny tinge.

Tremella.

On dead branches. Rare. Sowerby's Herbarium.

Name—*fimbriæ*, a fringe. From the appearance of the lobes. *Pers. Obs.* 2. p. 97. *Fr. Hym. Eur.* p. 690. *Berk. Out.* p. 286. *C. Hbk. n.* 1013.—*Bull. t.* 272. *Hoffm. Veg. Crypt.* i. t. 7. f. 1.

**2. T. frondosa** Fr.—*Yellow inclining to pale*, cæspitose, very large, *even*, plicate at the base; lobes gyroso-undulated.

The largest of the genus, nearest to *T. foliacea*, but thrice as large, more gyrose. There is no nucleus, and the structure is very different from that of *Nematelia*, wherefore it does not belong to that genus.

On roots of living trees, oak. Rare. Wothorpe, Norths. Careston, Forfarshire.

Colour a peculiar pale pinkish yellow. *M. F. B.* Name—*frondosus*, leafy. From the appearance of the lobes. *Fr. Hym. Eur.* p. 690. *Syst. Myc.* ii. p. 212. *Berk. Out.* p. 286. *C. Hbk. n.* 1014. *S. Mycol. Scot. n.* 961.—*Bull. t.* 499. f. T.

**3. T. foliacea** Pers.—*Cinnamon-flesh-colour*, cæspitose, cluster 2.5–5 cent. (1–2 in.), *even*, undulated, plicate at the base.

When young it is flattened down and nestles under the bark, wrinkled, then protruding from the plane base and forming the cluster. Kickx notes a var. growing on the ground.

On stumps, logs, &c. Frequent. Aug.–Dec.

Surface finely pubescent or granulated, the granulations pale, giving the plant a velvet-like gloss. *Eng. Bot.* Name—*folium*, a leaf. From the leafy appearance. *Pers. Obs.* 2. p. 98. *Berk. Out.* p. 287. *C. Hbk. n.* 1015. *S. Mycol. Scot. n.* 962.—*Bull. t.* 406. f. A. a. *Engl. Bot. t.* 1452.

**4. T. lutescens** Pers.—*Yellowish*, cæspitose, small, cluster 1–2.5 cent. ( $\frac{1}{2}$ –1 in.) broad, very soft, *undulato-gyrose*; lobes entire, naked.

Inclining to be fluid. Whitish when young.

On stumps, &c. Frequent. Aug.–Feb.

Name—*luteus*, yellow. Yellowish. *Pers. Syn.* p. 622.  *Ic. & descr.* p. 33. t. 8. f. 9. *Berk. Out.* p. 287. *C. Hbk. n.* 1016. *S. Mycol. Scot. n.* 963.—*Bull. t.* 406. f. B–D.

## II.—CEREBRINÆ. Firm then pulpy, &c.

**5. T. mesenterica** Retz.—*Orange*, simple, ascending, toughish, changeable in form, plicato-undulate, *gyrose*.

There is no constant form; when full grown pulverulent with the spores.

On sticks, *Ulex*, broom, &c. Very common. July–Dec.

Name—*μεσέντερον*, the mesentery. *Retz. in Vet. Ak. Handl.* 1769. p. 249. Tremella. *Fr. Hym. Eur.* p. 691. *Berk. Out.* p. 287. *C. Hbk. n.* 1017. *S. Mycol. Scot. n.* 964. *Jacqu. Misc. i. t.* 15. *Eng. Bot. t.* 709. *Hussey i. t.* 27.—*Schæff. t.* 168. *Bull. t.* 174. *Hoffm. Veg. Crypt. i. t.* 6. f. 4. *Fl. Dan. t.* 885.

**6. T. intumescens** Eng. Bot.—*Brown*, becoming black when dry, somewhat caespitose, rounded or conglomerate, soft, *obsoletely punctate*, somewhat tortuously lobed.

It is properly developed only in very damp weather, and forms numerous, rounded, soft, pulpy lobes, which are twisted and swollen, resembling the intestines of some animal.

On fallen trunks, beech, &c. Rare.

When cut longitudinally brown vertical streaks are discernible near the surface. *W.G.S.* Name—*intumescere*, to swell up. *Eng. Bot. t.* 1870. *Fr. Hym. Eur.* p. 691. *Syst. Myc. ii. p.* 215. *Berk. Out.* p. 288. *C. Hbk. n.* 1021. *S. Mycol. Scot. n.* 965.

**7. T. vesicaria** Eng. Bot.—*Pallid*, erect, *firm*, gelatinous within, undulated and gyrose.

On the ground. Very rare.

Spores 11 × 6 mk. *B. & Br.* Name—*vesica*, a bladder. *Eng. Bot. t.* 2451. *Fr. Hym. Eur.* p. 691. *Berk. Out.* p. 287. *C. Hbk. n.* 1018.

**8. T. albida** Huds.—*Whitish*, becoming fuscous when dry, 2.5 cent. (1 in.) broad, ascending, tough, expanded, undulated, somewhat gyrose, *pruinose*.

On dead wood. Common. Sept.—May.

Erumpent from bark. Spores oblong obtuse, curved, 2-guttate, subhyaline, 12–14 × 4–5 mk. *K.* Name—*albus*, white. Whitish. *Huds. Angl. ii. p.* 565. *Eng. Bot. t.* 2117. *Fr. Hym. Eur.* p. 691. *Syst. Myc. ii. p.* 215. *Berk. Out.* p. 287. *C. Hbk. n.* 1020. *S. Mycol. Scot. n.* 966.—*Bull. t.* 386. f. A.

### III.—CRUSTACEÆ. Effused, flattened.

**9. T. viscosa** Berk.—At first white then hyaline, effused, resupinate, undulated, somewhat viscous, the similar circumference naked.

On dead branches. Common. Jan.—Dec.

Forming a viscid stratum on the wood. Name—*viscus*, bird-lime. Viscous. *Berk. Out.* p. 288. *Ann. Nat. Hist. Ser. ii. vol. xiii. t.* 15. f. 4. *Fr. Hym. Eur.* p. 691. *C. Hbk. n.* 1027. *S. Mycol. Scot. n.* 967.—*Fl. Dan. t.* 1851. f. 1.

**10. T. epigæa** B. & Br.—*White*, effused, gelatinous, *gyroso-plicate*, sprinkled with the white spores.

Spreading over the naked soil, on which it forms a thin, white, gelatinous stratum.

Tremella

On the ground. Rare. Aug.-Sept.

Name—*ἐπί*, upon; *γῆ*, the earth. *B. & Br. n.* 373. *t.* 9. *f.* 3. *Berk. Out.* p. 289. *C. Hbk. n.* 1028. *S. Mycol. Scot. n.* 968.

## IV.—TUBERCULIFORMES. Small, somewhat erumpent.

11. **T. violacea** Relh.—Small, 6–8 mm. (3–4 lin.) broad, 2 mm. (1 lin.) thick, *violaceous*, becoming black when dry, erumpent, firm, somewhat compressed, *gyrose*.

Gregarious. The habit is that of *Dacrymyces*, but the structure is that of *Tremella*.

On trunks of pear-trees. Rare.

Name—from the colour. *Relh. Cant. p.* 442. *Fr. Hym. Eur. p.* 692. *Dacrymyces Berk. Out. p.* 290. *C. Hbk. n.* 1036. *Tulasne Ann. Sc. Nat.* 1853, *t.* 12. *f.* 3-12.

12. **T. indecorata** Sommerf.—*Dingy*, livid or olivaceous becoming fuscous, black fuscous when dry, 4–8 mm. (2–4 lin.) broad, 4 mm. (2 lin.) thick, erumpent, rounded, convex when swollen, *plicate*, opaque.

At first bursting through bark, firm, then plicate, connate as if with many lobes combined.

On birch, oak, willow. Rare. Oct.-Nov.

Name—*indecorus*, ugly. From its dingy appearance. *Sommerf. Lapp. p.* 306. *Fr. Hym. Eur. p.* 692. *Berk. Out. p.* 288. *C. Hbk. n.* 1022. *S. Mycol. Scot. n.* 969.

13. **T. moriformis** Eng. Bot.—Black, minute, somewhat gelatinous, firm, *spherical*, sinuous, opaque.

Habit exactly that of the mulberry.

On dead branches. Rare. Bath-easton. Dun. Forfarshire. Nov.

Interior mass translucent, violet. Staining paper violet. Name—*morus*, mulberry; *forma*, form. From its resemblance to a mulberry. *Eng. Bot. t.* 2446. *Berk. Out. p.* 287. *C. Hbk. n.* 1019. *S. Mycol. Scot. n.* 970. *Fr. Hym. Eur. p.* 692.

14. **T. tubercularia** Berk.—Erumpent, somewhat stipitate, *head pileate*, *dingy white*, nearly black when dry; stem round, short.

When dry it has a peculiar semi-transparent horny appearance.

On dead branches. Frequent. Oct.-Dec.

Name—*tuberculum*, a tubercle. *Berk. Out. p.* 288. *C. Hbk. n.* 1024. *S. Mycol. Scot. n.* 971. *Fr. Hym. Eur. p.* 692.

**15. T. foliicola** Fckl.—White above, brown below, scattered, Tremella crowded, somewhat sessile, granular, globose, gelatinous above, hard and dry below, when dry cup-shaped.

Conidia on the apices of branched sterigmata, fusiform curved, simple, hyaline,  $6 \times 2$  mk.

On the lower surface of leaves of *Rubus fruticosus* with *Phragmidium*. Castle Rising, 1882. March.

Name—*folium*, a leaf; *colo*, to inhabit. Fckl. *Symb. Myc.* p. 402. Grevillea, vol. xiii. p. 50. Sacc. *Fung. Ital.* n. 1024.

**16. T. torta** Berk.—Yellow or orange, minute, 4–6 mm. (2–3 lin.) broad, round, depressed, gyroso-tuberculate.

On dead wood, oak, &c. Common. Oct.–Dec.

Name—*torqueo*, to twist. Twisted. Berk. *Out.* p. 288. C. Hbk. n. 1025. S. *Mycol. Scot.* n. 972. Fr. *Hym. Eur.* p. 692.

**17. T. versicolor** Berk.—Orange, at length brown, minute, orbicular.

Tear-like, pallid when young.

On *Corticium nudum*. Uncommon.

Name—*versicolor*, changeable in colour. Berk. *Out.* p. 288. C. Hbk. n. 1026. S. *Mycol. Scot.* n. 973. Fr. *Hym. Eur.* p. 693.



GENUS LVIII.—**Exidia** (ἐξιδίω, to exude. Exuding from the receptacle). Fr. Syst. Myc. ii. p. 220. Exidia.

Distended with jelly when moist, tremulous, somewhat marginate, papillose, with a gelatinous coloured stratum covering the sporophores, the spicules of which are erumpent only at their apices.

An easily recognised genus from its habit and from the papillæ. Fr. *Hym. Eur.* p. 693.

I. AURICULINÆ (*auricula*, the external ear). Pezizoid.

II. SPICULARIÆ (*spiculum*, a little point, spicule). Somewhat plane, unpolished beneath, very evidently glandular, here and there gyrose.



XCVII. *Exidia glandulosa*.  
Natural size.

Exidia.

## I.—AURICULINÆ. Pezizoid.

1. **E. truncata** Fr.—Scarcely reaching 2.5 cent. (1 in.) in height and breadth, *very black*, shining, soft, distended with jelly, *disc truncato-plane*, glandular, at length cavernous, *rough with dots beneath*; stem very short.

It does not become swollen when moistened.

On lime branches. Yester Gardens. Dec.

Spores oblong, curved, chlorino-hyaline,  $14-16 \times 5-6$  mk. *K.* Name—*trunco*, to maim, shorten. Cut short, truncate. *Fr. Hym. Eur. p. 693. Syst. Myc. ii. p. 224. Scot. Nat. iii. p. 20. S. Mycol. Scot. n. 974.*

2. **E. recisa** Fr.—Fuscous-amber-colour, 1–2.5 cent. ( $\frac{1}{2}$ –1 in.) broad, very soft, distended with jelly, *disc truncato-plane*, *ribbed*, papillose, somewhat repand, rough with dots beneath; stem very short, *excentric*, oblique.

On dead wood, willow, &c. Common. Sept.–Oct.

Name—*recido*, to cut short. Shortened. *Fr. Hym. Eur. p. 693. Syst. Myc. ii. p. 223. Berk. Out. p. 289. C. Hbk. n. 1029. S. Mycol. Scot. n. 975. —Dittm. Sturm i. t. 13. Eng. Bot. t. 1819.*

## II.—SPICULARIÆ. Somewhat plane, &amp;c.

3. **E. glandulosa** Fr.—Blackish, 5–7.5 cent. (2–3 in.) broad, effused, *somewhat plane*, thick, *undulated*, studded with conical papillæ, *cinereous and somewhat tomentose beneath*.

Extremely variable in form and size.

On dead oak. Common. Sept.–March.

The under side feels like black crape. Sometimes truncate, sometimes pendulous. *M. F. B.* Known as Witch's butter. Spores oblong, curved, hyaline,  $12-14 \times 5$  mk. *K.* Name—*glandulæ*, glands. Full of glands. From the papillæ. *Fr. Hym. Eur. p. 694. Syst. Myc. ii. p. 224. Berk. Out. p. 289. C. Hbk. n. 1030. S. Mycol. Scot. n. 976. Hussey i. t. 42.—Bull. t. 420. f. 1. Fl. Dan. t. 884. Eng. Bot. t. 2448, 2452.*

4. **E. saccharina** Fr.—*Tawny-cinnamon*, effused, tubercular, *gyroso-undulated*, thick, sprinkled with few papillæ.

The colour is that of crystallised sugar. Papillæ minute, obtuse, vanishing.

On larch and Scotch fir. Rare. Nov.


Name—*saccharon*, sugar. From the colour. *Fr. Hym. Eur. p. 694. Berk. Out. p. 289. C. Hbk. n. 1031. S. Mycol. Scot. n. 977.*

GENUS LIX.—**Hirneola** (*hirnea*, a jug). Fr. Fung. Nat. p. 24. Hirneola.

Gelatinous inclining to cartilaginous, soft when moist, and tremulous but not distended with jelly; excipulum (cavity) cup-shaped, when dry coriaceous-horny, reviving when moistened, but scarcely swelling. The hard skin which forms the hymenium superior, discoid, of a different colour from the excipulum, and when soaked for a considerable time admitting of being separated from it entire.

Sporophores not involved in jelly; spores oblong, curved.

A remarkable genus, clearly distinct alike from *Auricularia* and *Exidia*. Fr. Hym Eur. p. 695.



XCVIII. *Hirneola auricula-Judæ*.  
One-third natural size.

1. **H. auricula-Judæ** Berk.—Thin, concave, flexuous, at length black, *venoso-plicate without and within*, tomentose and olive-cinereous beneath.

On elder and elm. Frequent. March–Aug.

Plant 2.5–7.5 cent. (1–3 in.) broad. Upper substance corrugated, the plaits branching from the middle part where they are strongest and somewhat convoluted, so as to give an idea of a human ear. When the plant grows on a perpendicular stump or tree it turns upwards. *Eng. Bot.* Name—Jew's-ear. *Berk. Out.* p. 289. t. 18. f. 7. *C. Hbk.* n. 1032. *S. Mycol. Scot.* n. 978. *Fr. Hym. Eur.* p. 695. *Hussey i. t.* 53.—*Bolt.* t. 107. *Mich. Gen.* t. 66. f. 1. B. more even. *Berk. Out.* t. 18. f. 7. C. reddish-fuscos. *Bull.* t. 427. f. 2. *Eng. Bot.* t. 2447.

GENUS LX.—**Femsjonía** (from Femsjö). Fr. Summ.  
Veg. Scand. p. 341.


Femsjonía.

Cup-shaped, distended with firm jelly, different on the two sides, disc thick, heterogeneous, even and not glandular. Sporophores immersed, globose, the oblong spores curved. *Growing on wood, erumpent, brightly coloured.* Fr. Hym. Eur. p. 695.

No British species.

Næmatelia. GENUS LXI.—**Næmatelia** (*ναιμαδ*, a word coined by Fries from *ναίω*, and applied to the gelatinous substance which surrounds the nucleus; *εἰλέω*, to roll or wrap round). Fr. Syst. Myc. ii. p. 227.

Solid, convex, immarginate, nucleus fleshy firm, entirely enveloped with a thick gelatinous stratum, which is internally fibroso-floccose, and everywhere covered with sporophores. *Growing on wood, rather firm, not shrivelling when dry.* Fr. Hym. Eur. p. 696.



XCIX. *Næmatelia encephala*.  
Natural size.

1. **N. encephala** Fr.—*Pallid-flesh-colour*, 8–12 mm. (4–6 lin.) broad and thick, somewhat sessile, pulvinate, plicato-rugose.

Solitary or clustered. Nucleus large, hard, white.

On pine branches, chiefly Scotch fir. Frequent. Sept.–Feb.

Looks like the brain of some animal. *M. J. B.* Name—*εν*, within; *κεφαλή*, head. Fr. Hym. Eur. p. 696. Syst. Myc. ii. p. 227. Berk. Out. p. 290. C. Hbk. n. 1033. S. Mycol. Scot. n. 979.—Willd. Bot. Mag. i. t. 4. f. 14.

2. **N. nucleata** Fr.—Whitish then *light yellow-fuscous*, 2–8 mm. (1–4 lin.) broad, sessile, becoming plane, somewhat gyrose.

On rotten wood. Rare.

Sometimes confounded with *Tremella albida* from which it differs in the presence of a small white nucleus. *M. J. B.* Name—*nucleus*, a kernel. Fr. Hym. Eur. p. 696. Syst. Myc. ii. p. 228. Berk. Out. p. 290. C. Hbk. n. 1034.

3. **N. virescens** Corda.—*Greenish*, sessile, somewhat orbicular, depressed, gyroso-tubercular.

On branches of *Ulex*, &c. Common. Jan.–Dec.

Name—*vireo*, to be green. Corda Ic. iii. f. 90. Fr. Hym. Eur. p. 696. Berk. Out. p. 290. C. Hbk. n. 1035. S. Mycol. Scot. n. 980.—Fl. Dan. t. 1857. f. 1.



GENUS LXII.—**Guepinia** (after Guepin). Fr. Elench. ii. p. 30. Guepinia.

Gelatinous inclining to cartilaginous, free, different on the two sides, variable in form, substipitate; hymenium confined to one side; sporophores linear, long-forked, 2-spored; spores curved. Fr. Hym. Eur. p. 697.

1. **G. peziza** Tul.—Yellow, cup-shaped, somewhat sessile, smooth on both sides, adnate behind; stem slender.

Very remarkable in having the hymenium on the upper side.

On wood. Shere.

Name—From its being like *Peziza*. Tul. Ann. Sc. Nat. 1853, p. 224. Fr. Hym. Eur. p. 697. Grevillea, vol. vi. p. 71. Pl. 97. f. 19, 20.



C. *Guepinia helvelloides*. One-half natural size.

GENUS LXIII.—**Dacrymyces** (δάκρυ, a tear; μύκης, a fungus). Dacrymyces. Nees Syst. p. 89.

Gelatinous, homogeneous, pervaded internally with septate fibres; conidia linked together like a necklace; sporophores clavate at the end of the filaments, bifurcate when mature; spores septate. Tul. Ann. Sc. Nat. 1853, t. II. f. 19. t. 12. Fr. Hym. Eur. p. 697.

\* Red.


\*\* Yellowish.

\*\*\* Pallid or fuscous.

\* Red.

1. **D. macrosporus** B. & Br.—Rose-colour, gelatinous, tuberculated.

Forming irregular gyrate and tuberculated masses of a rosy colour, about 6 mm. ( $\frac{1}{4}$  in.) long, parasitic on old remains of *Sphaeria stigma*. It preserves its rosy tint when dry.



CI. *Dacrymyces sebacus*. Natural size. Section five hundred times natural size.

Dacrymyces. On dead branches. Batheaston. Dec.—March.

Primary spores oblong, 3-5 septate, 40-50 × 8-11 mk. Secondary spores elliptic, 14 mk. *B. & Br.* Name—μακρός, long; σπόρος, seed. *B. & Br. n.* 1374. *t. 7. f. 1.*

\*\* *Yellowish.*

**2. *D. deliquescens*** Dub.—*Yellowish*, somewhat round, rooted, convex, immarginate, *at length twisted, hyaline*; spores triseptate.

At first pezizæform, even, dingy yellowish, becoming pale, yellow-fuscescent when dry.

On dead wood, pine rails, &c. Common. Jan.—Dec.

Easily distinguished from *D. stillatus* by its larger size, more caespitose habit, and paler yellowish colour. 6-12 mm. ( $\frac{1}{4}$ – $\frac{1}{2}$  in.) across. *M. F. B.* Spores triseptate, curved, oblong, hyaline, 12-22 × 6 mk. *K.* Name—*deliquesco*, to dissolve. Deliquescent. *Dub. Bot. Gall. p. 729. Fr. Hym. Eur. p. 698. Berk. Out. p. 290. C. Hbk. n. 1038. S. Mycol. Scot. n. 981. —Bull. t. 455. f. 3. Corda Ic. ii. f. 115.*

**3. *D. stillatus*** Nees.—*Yellow then orange*, colour persistent, 2-8 mm. (1-4 lin.) broad, somewhat round, convex, at length plicate, spores multiseptate.

Gregarious, somewhat confluent, variable in form, spherical when young, then turbinate or irregular, pezizoid, plicate, sessile or stipitate. It is readily distinguished from *D. deliquescens* by its colour being persistently bright.

On pine stumps, rails, &c. Common. Jan.—Dec.

Spores multiseptate, subhyaline, slightly curved, 10-22 × 8 mk. *K.* Name—*stillo*, to drop. *Nees Syst. p. 89. f. 90. Fr. Hym. Eur. p. 699. Syst. Myc. ii. p. 230. Berk. Out. p. 291. t. 18. f. 8. C. Hbk. n. 1039. S. Mycol. Scot. n. 982. Grev. t. 159. Corda Ic. ii. f. 114?*

**4. *D. chrysocomus*** Tul.—Golden, orbicular, *when young spherical*, immarginate, *soon collapsing, pezizoid*, and at length flattened, persistently *even*; spores multiseptate.

It agrees in its structure with *D. stillatus*. When dry it is very much contracted, somewhat horny.

On fir sticks. Rare.

Spores multiseptate, ellipsoid, pale yellow, 20-28 × 9-11 mk. *K.* Name—χρυσός, gold; κόμη, hair. Golden lock. *Tul.—Fr. Hym. Eur. p. 699. Berk. Out. p. 291. C. Hbk. n. 1040. S. Mycol. Scot. n. 983.—Peziza Fr. Syst. Myc. ii. p. 140. Bull. t. 376. f. 2. Sow. t. 152.*

\*\*\* *Pallid or fuscous.*

**5. *D. sebaceus*** B. & Br.—*Whitish*, somewhat round, composed

of filaments which are variously branched and often clavate Dacrymyces. above.

Conspicuous only in rainy weather. Forming circular patches. Individual plants 4-8 mm. (2-4 lin.) broad.

On twigs of ash and maple. Bath, 1868. Winter.

Spores ovato-triangular,  $14 \times 6-8$  mk. *B. & Br.* Name—*sebum*, tallow. From its resemblance to little spots of tallow. *B. & Br. n. 1305. t. 18. f. 2. C. Hbk. n. 1037. Fr. Hym. Eur. p. 699.*

**6. *D. succineus*** Fr.—*Amber-colour*, punctiform, somewhat gelatinous, smooth, becoming pale externally when moistened, disc darker and immarginate.

Gregarious, very small, rather thick, collapsing when dry as *D. chrysocomus* does.

On pine leaves. Grantown, Morayshire.


Minute, cup-shaped, stipitate, pale-yellow; stem short, stout, black; spores cylindrical,  $11 \times 1\frac{1}{2}$  mk. The British specimens exactly accord with those from Sprée in Rabenhorst's '*Fungi Europæi*,' No. 680. *Phillips.* Name—*succinum*, amber. From the colour. *Fr. Hym. Eur. p. 699. S. Mycol. Scot. n. 984.*

**7. *D. vermiformis*** B. & Br.—Grey, minute, worm-shaped. Sporophores globose; spores globose, pallid fuscous.

Sporophores 14 mk.; spores 6 mk.

On rotten wood. Batheaston. April.

Name—*vermis*, a worm; *forma*, form. *B. & Br. n. 1700. t. 3. f. 1.*



## A D D E N D A.

---

THE following were unfortunately omitted from Vol. I.:—

**Acetabularia**, a subgenus of *Agaricus*, belongs to the third series *Dermini*, and takes its place before Subgenus XIX., *Pholiota*. Acetabularia.

**Acetabularia** (*acetabulum*, a vinegar cup. From the cup-like volva). Berk. Linn. Journ. xviii. p. 389.

Universal veil distinct from the pileus; hymenophore distinct; gills free; spores pallid tawny or brown.

Analogous to *Volvaria* and *Chitonina*.

**1. A. acetabulosus** Sow.—**Pileus** about 2.5 cent. (1 in.) broad, light-brown, convex, striate especially at the margin. **Stem** 5 cent. (2 in.) long, white, equal, hollow; volva nearly white. **Gills** free, narrow, hispid, light-brown.



CII. *Acetabularia*. One-half natural size.

Bank of Thames near high-water mark at Millbank, 1795. May.

This is very like a poor specimen of *A. congregatus*, but the pileus is more plaited. The lamellæ are remarkably glandular on their sides; and instead of a bare base or foot, it stands in a little socket-like volva. Sow. The accompanying illustration is from the original specimen in the British Museum. Sowerby gives also an enlarged section of the gills, and a side of a gill showing very large cystidia. The pileus appears to be slightly split as in *A. rimosus*. Name—from the cup-like volva. Sow. t. 303. Berk. Linn. Journ. xviii. p. 389. C. *Illust. Pl.* 345.

## Hiatula

**Hiatula** holds as yet an uncertain position. From the colour of the spores, and the habit being that of *Mycena*, it might be referred to the *Leucospori*, but the mode of growth is entirely that of the *Coprini*. At present it is placed next to *Coprinus*.



CIII. *Hiatula Wynnii*. One-half natural size.

**Hiatula** (*hio*, to gape. From the manner in which the pileus splits open). Mont. Ann. Sc. Nat. 1854. Vol. i. p. 107.

Pileus very thin, without a pellicle, formed by the junction of the back of the gills, when expanded veliform and splitting open as in the most delicate *Coprini*, but not melting away, and the sporidia are white. *Fr. Nov. Symb. Myc. p. 11.*

1. **H. Wynnii** B. & Br.—Shining white. **Pileus** 4 cent. ( $1\frac{1}{2}$  in.) broad, tender, striate, pulverulent, centre darker. **Stem** 2.5 cent. (1 in.) long, 2 mm. (1 lin.) thick, slender, striate. **Gills** scarious, rather broad.

One specimen became darker in drying, and had a longer and thicker stem.

In a stove. Kew.

Name—after Mrs Lloyd Wynne. *B. & Br. n. 1772.*

## GLOSSARY.

---

ADNATE (*adnascor*, to grow to), growing into; of gills *e.g.* closely attached to the stem.

ADNEXED (*ad*, to; *necto*, to join), of gills attached to the stem but not adnate to it.

ADRESSED (*ad*, to; *premo*, to press), pressed in close contact.

ÆRUGINOUS (*ærugeo*, rust of copper), verdigris-green.

AGGLUTINATED (*agglutino*, to cement), glued to a surface.

ALVEOLATE (*alveus*, a hollow), in the form of small depressions like shallow honeycomb.

AMPHIGENOUS (*ἀμφί*, about; *γεννᾶω*, to beget), not confined to one surface, all round; *e.g.* hymenium of *Clavaria*.

ANASTOMOSING (*ἀναστόμωσις*, a bringing to a point, a contraction), uniting by running together irregularly.

ANNULAR (*annulus*, a ring), in the form of a ring.

APICAL (*apex*, the top), close to the apex.

APICULATE (*apex*), terminating in a small point.

APPENDICULATE (*appendix*, an appendage), hanging in small fragments.

APPROXIMATE (*ad*, to; *proximo*, to approach), of gills which approach but do not reach the stem.

ARACHNOID (*ἀράχνη*, a spider's web; *εἶδος*, resemblance), like a cobweb.

ARCUATE (*arcus*, a bow), bow-shaped.

AREOLATE (*area*, a plot), divided into little areas or patches.

ASCENDING (*ascendo*, to climb), inclining upwards.

ATOMATE (*ἄτομος*, an atom), sprinkled with atoms.

BULLATE (*bullā*, a bubble), furnished with a boss or stud.

CÆSPITOSE (*cæspes*, a clump), growing in tufts.

CALYPTRA (*calyptra*, a hood), applied *e.g.* to the portion of the volva covering the pileus.

CAMPANULATE (*campana*, a bell), bell-shaped.

CINEREOUS (*cinis*, ashes), ash-coloured.

CIRCINATE (*circino*, to make round), rounded.

CLAVATE (*clava*, a club), club-shaped.

- CONFEROID (*Conferva*, a name applied to certain of the *Algæ*, sea-weeds; *εἶδος*, form), like a *Conferva*; from the finely branched threads.
- CONNATE (*con*, together; *nascor*, to be born), united by growing together.
- CONTINUOUS (*continuo*, to unite), without a break; *e.g.* of a surface which is not cracked, or of one part which runs into another without interruption.
- CORRUGATED (*corrugo*, to wrinkle), drawn into wrinkles or folds.
- CORTICATED (*cortex*, bark), furnished with bark-like covering.
- CORTINA (*cortina*, a veil), a veil of a spider-web structure; applied to the peculiar veil of the genus *Cortinarius*.
- CORTINATE, furnished with a cortina.
- CRENATE (*crena*, a notch), notched at the edge. The notches are blunt or rounded, not sharp as in a serrated edge.
- CRIBROSE (*cribra*, a sieve), pierced with holes.
- CUSPIDATE (*cuspis*, a point, spear), with a sharp spear-like point.
- CYATHIFORM (*cyathus*, a cup; *forma*, form), cup-shaped.
- DECURRENT (*decurro*, to run down), running down; *e.g.* of gills on the apex of the stem.
- DETERMINATE (*determino*, to limit), ending definitely; having a distinctly defined outline.
- DIAPHANOUS (*διδ*, through; *φαίνω*, to appear), of a transparent nature.
- DICHOTOMOUS (*δίχα*, *τέμνω*, to cut), divided into two; regularly forked.
- DIMIDIATE (*dimidio*, to halve), halved; *e.g.* of gills which reach half-way to the stem.
- DISCIFORM (*discus*, a quoit; *forma*, form), disk-shaped; of a circular and flat form.
- DISSEPIMENTS (*dissepimentum*, a partition), dividing walls.
- DISTANT (*disto*, to stand apart), far apart; of gills which have a wide distance between them.
- DIVARICATE (*divarico*, to spread asunder), separating at an obtuse angle.
- ECHINULATE (*echinus*, a hedgehog), beset with short bristles.
- EFFUSED (*effundo*, to pour out), spread over without regular form.
- EMARGINATE (*e*, out of; *margo*, the margin), of gills with a sudden curve as if scooped out at the point of attachment to the stem.
- EPIPHYTAL (*ἐπί*, upon; *φύτον*, a plant), growing upon another plant.
- EVEN, of a surface which is quite plane as contrasted *e.g.* with one which is striate, pitted, &c. Distinguished from smooth. A surface may not be smooth, and yet even.
- FARINACEOUS (*farina*, meal), mealy.
- FASCIATE (*fascia*, a band or girth), zoned with coloured bands.
- FASCICULATE (*fascis*, a bundle), growing in small bundles.
- FASTIGIATE (*fastigium*, the top), reaching the top.
- FIBRILLOSE (*fibra*, a fibre), clothed with small fibres.
- FIBROUS, composed of fibres.
- FILIFORM (*filum*, a thread; *forma*, form), thread-like.
- FIMBRIATE (*fimbriæ*, a fringe), fringed.
- FISSILE (*findo*, to split), capable of being split.
- FISTULOSE (*fistula*, a pipe), hollow in the centre like a pipe.
- FLOCCI (*flocus*, a lock of wool), woolly locks.



FLOCCULOSE, covered with small flocci.

FREE, of gills which reach the stem but are not attached to it.

FURFURACEOUS (*furfur*, bran), with branny scales or scurf.

FUSCOUS (*fuscus*, dusky), brownish in colour, but dingy, not pure.

GIBBOUS (*gibbus*, hump-backed), in the form of a swelling; of a pileus *e.g.* which is more convex or tumid on one side than the other.

GILLS, the plates of an Agaric on which the hymenium is situated.

GREAVED (*grève*, the shin-bone), of a stem clothed like a leg in armour.

GRUMOUS (*grumus*, a little heap), clotted; of flesh *e.g.* composed of little clustered grains.

GUTTATE (*gutta*, a tear), marked with tear-like spots.

GYROSE (*γῦρος*, a circle), circling in wavy folds.

HETEROGENEOUS (*ἑτερος*, one of two; *γεννάω*, to beget), of a structure which is different from adjacent ones.

HOMOGENEOUS (*ὁμός*, one and the same; *γεννάω*, to beget), similar in structure.

HYALINE (*ὑἄλος*, clear), transparent.

HYBERNACULUM (*hiberno*, to winter), applied to bodies which are the forms in which certain fungi, *e.g.* *Typhulæ*, pass the winter.

HYGROPHANOUS (*ὑγρος*, moist; *φαίνω*, to appear), of a watery appearance when moist and opaque when dry.

HYMENIUM (*ὑμήν*, a membrane), the fruit-bearing surface; *e.g.* on each side of the gills of an Agaric.

HYMENOPHORE (*ὑμήν*, a membrane; *φέρω*, to bear), the structure which bears the hymenium; in Agarics *e.g.* the under surface of the pileus to which the gills are attached.

IMMARGINATE (*in*, negative; *margo*, a margin), without a distinct border.

IMMERSED (*immergo*, to immerse), sunk into the matrix.

INFERIOR, of the ring of an Agaric which is far down on the stem.

INFUNDIBULIFORM (*infundibulum*, a funnel; *forma*, form), funnel-shaped.

INNATE (*in*, into; *nascor*, to be born), adhering by growing into.

INSERTED (*insero*, to insert), growing like a graft from its stock.

INVOLUTE (*in*; *volvo*, to roll), rolled inwards.

ISABELLINE, of the colour of soiled linen; brownish-yellow-clay.

LACINIATE (*lacinia*, a lappet, flap), divided into flaps.

LACUNOSE (*lacuna*, a pit), marked with small hollows.

LANCEOLATE (*lancea*, a spear), lance-shaped; tapering to both ends.

LINEAR (*linea*, a line), narrow and straight.

MAMMIFORM (*mamma*, a breast; *forma*, form), breast-shaped.

MARGINATE (*margo*, a margin), having a distinct border.

MATRIX (*matrix*, the womb), the substance on which a fungus grows.

MEDIAL (*medium*, the middle), at the middle; of the ring of an Agaric *e.g.* which is between *superior* or near the apex of the stem, and *distant* or far removed from the apex.

MERISMOID (Merisma, from *μερίζω*, to divide, a generic name applied by Persoon to certain fungi with branched or lacinate pilei, now included under

*Thelephora*; *εἶδος*, form), resembling a *Merisma*—*i.e.* having a branched or lacinate pileus.

MYCELIUM (*μύκης*, a fungus), spawn of fungi resulting from the germination of spores; in *Agarics* *e.g.* forming root-like threads.

OBCONIC (*ob*; *conus*, a cone), inversely conical.

OBOVATE (*ob*; *ovum*, an egg), inversely egg-shaped.

PAPILLATE (*papilla*, a nipple), furnished with one or more nipple-like elevations.

PARTIAL (*pars*, a part), of a veil clothing the stem and reaching to the edge of the pileus, but not extending beyond it.

PECTINATE (*pecten*, a comb), like the teeth of a comb.

PELLICULOSE (*pellis*, skin), furnished with a pellicle or distinct skin.

PENCILLED (*penicillum*, a painter's brush), with pencil-like hairs either on the tip or border.

PERONATE (*pero*, a kind of high boot), sheathed; of a stem *e.g.* which has a covering like a legged boot.

PERVIOUS (*per*, through; *via*, a way), forming an open tube-like passage.

PILEUS (*pileus*, a hat), a part of the receptacle of a fungus; *e.g.* the cap-like head of *Agarics*.

PILEOLI, secondary pilei; arising from the division of a primary pileus.

PILOSE (*pilus*, a hair), covered with hairs.

PORIFORM (*porus*, a pore; *forma*, form), in the form of pores.

POROUS, furnished with pores.

PROLIFEROUS (*proles*, offspring; *fero*, to bear), applied to an organ which gives rise to secondary ones of the same kind.

PRUINOSE (*pruina*, hoar-frost), covered with frost-like bloom.

PRUNIFORM (*prunus*, a plum; *forma*, form), plum-shaped.

PULVINATE (*pulvinus*, a cushion), cushion-shaped.

PUNCTATE (*punctus*, a point), dotted with points.

QUATERNATE (*quatuor*, four; *nascor*, to be born), arranged in groups of four.

REMOTE (*removeo*, to remove), of gills which do not reach the stem, but leave a free space between them and it.

RENIFORM (*renes*, the kidneys; *forma*, form), kidney-shaped.

REPAND (*repando*, to throw open), bent backwards.

RESUPINATE (*resupino*, to throw on the back), lying on the back; of an *Agaric* *e.g.* lying on the back without a stem; or of fungi spread over the matrix without any stem and with the hymenium upwards.

REVOLUTE (*revolvo*, to roll back), rolled backwards; of the margin of a pileus *e.g.*, the opposite of *involute*.

RIMOSE (*rima*, a crack), cracked.

RING, a part of the veil adhering in the form of a ring to the stem of an *Agaric*.

RIVULOSE (*rivula*, a small stream), marked with lines like rivulets.

RUBIGINOUS (*rubigo*, rust), rust-coloured.

RUFESCENT (*rufesco*, to become reddish), reddish in colour.

RUGOSE (*ruga*, a wrinkle), wrinkled.

- SCABROUS (*scaber*, rough), rough on the surface.
- SCARIOUS (*σκάδρος*, a kind of sea-fish), applied to a shrivelled membrane.
- SCISSILE (*scindo*, to cleave), capable of being cleft asunder; of two plates *e.g.* lying together, but capable of being separated.
- SCLEROTIROID (*σκληρότος*, hard; *εἶδος*, resemblance), in the form of a *Sclerotium*, a form assumed by the mycelium of certain fungi.
- SCROBICULATE (*scrobis*, a trench), marked with small pits.
- SEPARATING (*separo*, to separate), becoming detached as gills from the stem, or resupinate fungi from the matrix.
- SERRATE (*serra*, a saw), with teeth like a saw.
- SESSILE (*sedeo*, to sit), seated without a stem.
- SINUATE (*sinus*, a curve), waved; of a pileus with a wavy margin; of gills which have a sudden wave or sinus where they reach the stem.
- SMOOTH, applied to a surface which is destitute of hairs, &c. See Even.
- SPATHULATE (*spatha*, a broad flat instrument for stirring liquids), shaped like a spatula or spoon.
- SQUAMOSE (*squama*, a scale), scaly.
- SQUAMULOSE, covered with small scales.
- SQUARROSE (*squarrosus*, rough), rough with scales.
- STIPITATE (*stipes*, a stalk), stemmed.
- STOLONIFEROUS (*stolo*, a sucker; *fero*, to bear), stolon-bearing; applied to a trailing and rooting branch.
- STRIATE (*stria*, a furrow, flute of a column), marked with lines in the form of channels.
- STRIGOSE (*striga*, a swath), rough with fascicles of hairs.
- STUFFED, of a stem filled with substance of a different texture from its walls.
- SUBICULUM (*subiculum*, an under layer), the thready mycelium forming the under layer of the plant.
- SULCATE (*sulcus*, a furrow), marked with furrows.
- SUPERIOR (*super*, above), the upper surface; or applied to a ring when it is near the apex of the stem.
- TOMENTOSE (*tomentum*, wool, &c.), downy.
- TRAMA (*trama*, the web or filling of a web), the substance proceeding from the hymenophore, between the plates of (central in) the gills in Agarics, and between the double membranes of which the dissepiments of the pores are composed in *Polyporei*.
- TREMELLOID (*tremo*, to tremble), of the consistency of jelly.
- TRUNCATE (*trunco*, to maim), ending abruptly as if cut short.
- TUBÆFORM (*tuba*, a trumpet; *forma*, form), trumpet-shaped.
- TUBERCLE (*tuber*, a swelling), a small wart-like excrescence.
- UMBILICATE (*umbilicus*, the navel), with a central depression.
- UMBONATE (*umbo*, the boss of a shield), with a central boss-like elevation.
- UNCINATE (*uncinus*, a hook), hooked.
- UNIVERSAL (*universus*, whole), of the veil or volva which entirely envelops the fungus when young.
- VEIL (*velum*, a covering), a covering of various texture more or less completely enwrapping a fungus; occurring chiefly among the *Agaricini*.
- VELIFORM (*velum*; *forma*, form), of a thin veil-like covering.

VENTRICOSE (*venter*, the belly), swollen in the middle.

VERNICOSE (*vernix*, varnish), shining as if varnished.

VILLOUS (*villus*, a tuft of hair), downy with soft hairs.

VIRGATE (*virga*, a twig, stripe), streaked.

VOLVA (*volva*, a wrapper), a substance (universal veil) covering the plant ;  
*e.g.* in young *Amanitæ*.

ZONED (*zona*, a girdle), furnished with one or more concentric circles of colour,  
depression, &c.

# INDEX TO VOL. II.

	PAGE		PAGE
ACETABULARIA Berk. . . . .	321	BOLETUS Dill.—continued.	
acetabulosus Sow. . . . .	321	piperatus Bull. . . . .	171
AGARICUS Linn. . . . .		pruinatus Fr. . . . .	174
adhesivus With. . . . .	156	purpureus Fr. . . . .	178
filamentosus Fr. . . . .	68	radicans Pers. . . . .	173
glaucoptus Sow. . . . .	5	rubinus Smith . . . . .	173
helvelloides Bull. . . . .	136	sanguineus With. . . . .	170
obscuratus Lasch. . . . .	112	satanas Lenz. . . . .	177
paradoxus Kalchbr. . . . .	67	scaber Fr. . . . .	180
sublanatus Huss. . . . .	3	spadiceus Schæff. . . . .	173
tristis Krombh. . . . .	112	striipes Secr. . . . .	172
turbinatus Sow. . . . .	8	strobilaceus Scop. . . . .	182
turfosus Sow. . . . .	161	subtomentosus Linn. . . . .	172
validus Berk. . . . .	26	sulphureus Fr. . . . .	171
ARRHENIA Fr. . . . .	137	varicolor B. & Br. . . . .	174
AURICULARIA Bull. . . . .	271	variegatus Sw. . . . .	171
lobata Sommerf. . . . .	272	versipellis Fr. . . . .	179
mesenterica Fr. . . . .	272	viscidus Linn. . . . .	179
BOLETUS Dill. . . . .	166	CALOCERA Fr. . . . .	301
æreus Bull. . . . .	176	cornea Fr. . . . .	302
æstivalis Fr. . . . .	177	corticalis Fr. . . . .	302
alutarius Fr. . . . .	180	glossoides Fr. . . . .	303
arboresus Sow. . . . .	229	palmata Fr. . . . .	302
badius Fr. . . . .	170	striata Fr. . . . .	303
bovinus Linn. . . . .	170	stricta Fr. . . . .	303
calopus Fr. . . . .	175	tuberosa Fr. . . . .	300
carnosus Rostk. . . . .	182	viscosa Fr. . . . .	302
castaneus Bull. . . . .	181	CANTHARELLUS Adans. . . . .	131
chrysenteron Fr. . . . .	172	albidus Fr. . . . .	133
collinitus Fr. . . . .	169	anthracophilus Lev. . . . .	133
cyanescens Bull. . . . .	181	aurantiacus Fr. . . . .	132
edulis Bull. . . . .	176	Brownii B. & Br. . . . .	133
elegans Schum. . . . .	168	carbonarius A. & S. . . . .	133
erythropus Pers. . . . .	178	cibarius Fr. . . . .	132
felleus Bull. . . . .	180	cinereus Fr. . . . .	135
flavidus Fr. . . . .	169	cupulatus Fr. . . . .	135
flavus With. . . . .	168	Friesii Quel. . . . .	132
fragrans Vitt. . . . .	176	Houghtoni Phill. . . . .	134
granulatus Linn. . . . .	169	infundibuliformis Fr. . . . .	134
impolitus Fr. . . . .	177	lobatus Fr. . . . .	137
laricinus Berk. . . . .	179	lutescens Bull. . . . .	134
luridus Schæff. . . . .	178	muscigenus Fr. . . . .	136
luteus Linn. . . . .	168	radicosus B. & Br. . . . .	133
olivaceus Schæff. . . . .	175	retirugus Fr. . . . .	136
pachypus Fr. . . . .	175	Stevensoni B. & Br. . . . .	136
pachypus Krombh. . . . .	176	tubæformis Fr. . . . .	134
parasiticus Bull. . . . .	174	umbonatus Fr. . . . .	133
		CLADODERRIS Pers. . . . .	266
		minima B. & Br. . . . .	266

	PAGE		PAGE
<b>CLAVARIA</b> Linn.	290	<b>CORTICIUM</b> Fr.—continued.	
<i>abietina</i> Pers.	294	<i>ferrugineum</i> Fr.	288
<i>acuta</i> Sow.	301	<i>foetidum</i> B. & Br.	275
<i>amethystina</i> Bull.	291	<i>giganteum</i> Fr.	274
<i>Ardenia</i> Sow.	300	<i>incarnatum</i> Fr.	279
<i>argillacea</i> Fr.	297	<i>lactescens</i> Berk.	276
<i>aurantiaca</i> Pers.	297	<i>lacteum</i> Fr.	274
<i>aurea</i> Schæff.	294	<i>lacunosum</i> B. & Br.	284
<i>botrytes</i> Pers.	290	<i>læve</i> Pers.	275
<i>canaliculata</i> Fr.	300	<i>laxum</i> Fr.	282
<i>ceranoides</i> Pers.	297	<i>limitatum</i> Fr.	280
<i>cinerea</i> Bull.	292	<i>lividum</i> Pers.	278
<i>condensata</i> Fr.	295	<i>maculæforme</i> Fr.	280
<i>contorta</i> Holmsk.	299	<i>nudum</i> Fr.	279
<i>coralloides</i> Linn.	292	<i>ochraceum</i> Fr.	278
<i>crispula</i> Fr.	296	<i>olivaceum</i> Fr.	283
<i>cristata</i> Pers.	292	<i>polygonium</i> Pers.	280
<i>crocea</i> Pers.	295	<i>porosum</i> B. & Curt.	275
<i>curta</i> Fr.	291	<i>puberum</i> Fr.	277
<i>fastigiata</i> Linn.	291	<i>puteum</i> Fr.	281
<i>fistulosa</i> Fr.	299	<i>quercinum</i> Fr.	278
<i>flaccida</i> Fr.	295	<i>roseum</i> Pers.	275
<i>formosa</i> Pers.	294	<i>salicinum</i> Fr.	273
<i>fragilis</i> Holmsk.	298	<i>sambuci</i> Fr.	283
<i>fumosa</i> Pers.	298	<i>sanguineum</i> Fr.	276
<i>fusiformis</i> Sow.	297	<i>scutellare</i> B. & Curt.	278
<i>gracilis</i> Pers.	298	<i>serum</i> Fr.	283
<i>grisea</i> Pers.	295	<i>stabile</i> Fr.	282
<i>inæqualis</i> Fl. Dan.	297	<i>subdealbatum</i> B. & Br.	276
<i>incarnata</i> Weinm.	301	<i>sulphureum</i> Fr.	276
<i>juncea</i> Fr.	300	<i>typhæ</i> Fckl.	281
<i>juncea</i> Lev.	305	<i>umbrinum</i> Fr.	282
<i>Kromholzii</i> Fr.	293	<i>velutinum</i> Fr.	275
<i>Kunzei</i> Fr.	293	<i>violaceo-lividum</i> Fr.	280
<i>ligula</i> Fr.	299		
<i>muscoïdes</i> Linn.	291	<b>CORTINARIUS</b> Fr.	1
<i>pistillaris</i> Linn.	299	<i>acutus</i> Fr.	61
<i>pratensis</i> Pers.	291	<i>albo-violaceus</i> Fr.	24
<i>purpurea</i> Fr.	296	<i>anfractus</i> Fr.	7
<i>pyxidata</i> Pers.	293	<i>anomalous</i> Fr.	32
<i>rosea</i> Fr.	296	<i>anthracinus</i> Fr.	34
<i>rufa</i> Fl. Dan.	296	<i>arenatus</i> Fr.	28
<i>rugosa</i> Bull.	293	<i>argentatus</i> Fr.	22
<i>spinulosa</i> Pers.	294	<i>armeniacus</i> Fr.	52
<i>stricta</i> Pers.	295	<i>armillatus</i> Fr.	43
<i>tenuipes</i> B. & Br.	300	<i>arvinaceus</i> Fr.	17
<i>tuberosa</i> Sow.	300	<i>bivelus</i> Fr.	38
<i>umbrina</i> Berk.	292	<i>bolaris</i> Fr.	26
<i>uncialis</i> Grev.	301	<i>bulbosus</i> Fr.	39
<i>vermicularis</i> Scop.	298	<i>Bulliardii</i> Fr.	26
		<i>cærulescens</i> Fr.	9
<b>CORTICIUM</b> Fr.	272	<i>cærulescens</i> Saund. & Sm.	14
<i>amorphum</i> Fr.	274	<i>callisteus</i> Fr.	26
<i>anthochroum</i> Fr.	284	<i>calochrous</i> Fr.	9
<i>arachnoideum</i> Berk.	275	<i>camphoratus</i> Fr.	24
<i>aridum</i> Fr.	282	<i>camurus</i> Fr.	30
<i>atrovirens</i> Fr.	277	<i>caninus</i> Fr.	31
<i>aurora</i> Berk.	281	<i>castaneus</i> Fr.	55
<i>byssoides</i> Fr.	282	<i>cinnabarinus</i> Fr.	33
<i>cæruleum</i> Fr.	277	<i>cinnamomeus</i> Fr.	35
<i>calceum</i> Fr.	277	<i>claricolor</i> Fr.	3
<i>cinereum</i> Fr.	279	<i>collinitus</i> Fr.	17
<i>cinnamomeum</i> Fr.	276	<i>colus</i> Fr.	56
<i>citrinum</i> Pers.	279	<i>cotoneus</i> Fr.	36
<i>comedens</i> Fr.	281	<i>croceo-cæruleus</i> Fr.	15
<i>confluens</i> Fr.	279	<i>cumatilis</i> Fr.	13
<i>corrugatum</i> Fr.	280	<i>cyanites</i> Fr.	23
<i>evolvens</i> Fr.	273	<i>cyanopus</i> Fr.	5
		<i>decipiens</i> Fr.	59

CORTINARIUS *Fr.*—continued.

	PAGE
decolorans <i>Fr.</i>	15
decoloratus <i>Fr.</i>	14
decumbens <i>Fr.</i>	30
delibutus <i>Fr.</i>	20
detonsus <i>Fr.</i>	60
diabolicus <i>Fr.</i>	31
dibaphus <i>Fr.</i>	11
dilutus <i>Fr.</i>	54
duracinus <i>Fr.</i>	53
elatio <i>Fr.</i>	18
erythrinus <i>Fr.</i>	59
evernius <i>Fr.</i>	42
fasciatus <i>Fr.</i>	62
finitimus <i>Weinm.</i>	25
flabellus <i>Fr.</i>	48
flexipes <i>Fr.</i>	47
fulgens <i>Fr.</i>	11
fulmineus <i>Fr.</i>	12
gentilis <i>Fr.</i>	45
germanus <i>Fr.</i>	60
glaucopus <i>Fr.</i>	8
grallipes <i>Fr.</i>	19
hæmatochelis <i>Fr.</i>	44
helvelloides <i>Fr.</i>	46
helvolus <i>Fr.</i>	44
hemitrichus <i>Fr.</i>	50
hinnuleus <i>Fr.</i>	45
iliopodius <i>Fr.</i>	49
illibatus <i>Fr.</i>	20
imbutus <i>Fr.</i>	55
impennis <i>Fr.</i>	41
incisus <i>Fr.</i>	49
infucatus <i>Fr.</i>	36
jubarinus <i>Fr.</i>	57
Jungkuhnii <i>Fr.</i>	62
largus <i>Fr.</i>	5
leucopus <i>Fr.</i>	58
licinipes <i>Fr.</i>	40
limonius <i>Fr.</i>	44
livido-ochraceus <i>Berk.</i>	19
macropus <i>Fr.</i>	38
miltinus <i>Fr.</i>	33
milvinus <i>Fr.</i>	62
mucifluus <i>Fr.</i>	18
multiformis <i>Fr.</i>	7
myrtillinus <i>Fr.</i>	32
napus <i>Fr.</i>	8
obtusus <i>Fr.</i>	61
ochroleucus <i>Fr.</i>	29
orellanus <i>Fr.</i>	35
orichalceus <i>Batsch</i>	12
paleaceus <i>Fr.</i>	51
papulosus <i>Fr.</i>	16
penicillatus <i>Fr.</i>	28
periscelis <i>Fr.</i>	47
pholideus <i>Fr.</i>	27
plumiger <i>Fr.</i>	41
pluvius <i>Fr.</i>	21
porphyropus <i>Fr.</i>	15
prasinus <i>Fr.</i>	13
privignus <i>Fr.</i>	53
psammoccephalus <i>Fr.</i>	48
punctatus <i>Fr.</i>	46
purpurascens <i>Fr.</i>	10
quadricolor <i>Fr.</i>	43
raphanoides <i>Fr.</i>	36
Reedii <i>Berk.</i>	58
redimitus <i>Fr.</i>	25

CORTINARIUS *Fr.*—continued.

	PAGE
renidens <i>Fr.</i>	56
Riederi <i>Fr.</i>	6
rigens <i>Fr.</i>	58
russus <i>Fr.</i>	6
saginus <i>Fr.</i>	6
salor <i>Fr.</i>	19
sanguineus <i>Fr.</i>	34
saturninus <i>Fr.</i>	54
scandens <i>Fr.</i>	59
scaurus <i>Fr.</i>	13
scutulatus <i>Fr.</i>	42
sebaceus <i>Fr.</i>	4
spilomeus <i>Fr.</i>	33
stemmaus <i>Fr.</i>	50
stillatitius <i>Fr.</i>	21
subferrugineus <i>Fr.</i>	52
sublanatus <i>Fr.</i>	27
subpurpurascens <i>Fr.</i>	10
tabularis <i>Fr.</i>	30
talus <i>Fr.</i>	8
tophaceus <i>Fr.</i>	25
torvus <i>Fr.</i>	40
traganus <i>Fr.</i>	24
triformis <i>Fr.</i>	47
triumphans <i>Fr.</i>	2
turbinatus <i>Fr.</i>	11
tormalis <i>Fr.</i>	3
uliginosus <i>Berk.</i>	35
uraceus <i>Fr.</i>	57
urbicus <i>Fr.</i>	39
varius <i>Fr.</i>	4
venetus <i>Fr.</i>	37
violaceus <i>Fr.</i>	23
CRATERELLUS <i>Fr.</i>	259
clavatus <i>Fr.</i>	261
cornucopioides <i>Pers.</i>	260
crispus <i>Fr.</i>	260
lutescens <i>Fr.</i>	259
sinuosus <i>Fr.</i>	260
CYPHELLA <i>Fr.</i>	284
Bloxami <i>B. &amp; Phill.</i>	284
brunnea <i>Phill.</i>	285
capula <i>Fr.</i>	287
catilla <i>Smith</i>	286
cuticulosa <i>Berk.</i>	288
dochmiospora <i>B. &amp; Br.</i>	288
fraxinicola <i>B. &amp; Br.</i>	286
fulva <i>B. &amp; Rav.</i>	285
galeata <i>Fr.</i>	286
Goldbachii <i>Weinm.</i>	288
griseo-pallida <i>Weinm.</i>	285
lacera <i>Fr.</i>	287
musciicola <i>Fr.</i>	286
muscigena <i>Fr.</i>	286
ochroleuca <i>B. &amp; Br.</i>	285
pallida <i>B. &amp; Br.</i>	287
Pini <i>Phill.</i>	287
stuppea <i>B. &amp; Br.</i>	285
DACRYMYCES <i>Nees</i>	317
chrysocomus <i>Tul.</i>	318
deliquescent <i>Dub.</i>	318
macrosporus <i>B. &amp; Br.</i>	317
sebaceus <i>B. &amp; Br.</i>	318
stillatus <i>Nees</i>	318
succineus <i>Fr.</i>	319
verniformis <i>B. &amp; Br.</i>	319

	PAGE	HYDNUM Linn.—continued.	PAGE
<i>DÆDALEA Pers.</i> . . . . .	224	<i>ferrugineum Fr.</i> . . . . .	237
<i>aurea Fr.</i> . . . . .	224	<i>ferruginosum Fr.</i> . . . . .	242
<i>cinerea Fr.</i> . . . . .	225	<i>fragile Fr.</i> . . . . .	235
<i>confragosa Pers.</i> . . . . .	225	<i>fuscum Berk.</i> . . . . .	242
<i>ferruginea Schum.</i> . . . . .	226	<i>gelatinosum Pers.</i> . . . . .	247
<i>latissima Fr.</i> . . . . .	225	<i>graveolens Delast.</i> . . . . .	238
<i>quercina Pers.</i> . . . . .	224	<i>imbricatum Linn.</i> . . . . .	234
<i>unicolor Fr.</i> . . . . .	225	<i>lævigatum Sw.</i> . . . . .	235
<i>vermicularis Pers.</i> . . . . .	226	<i>limonicolor B. &amp; Br.</i> . . . . .	243
<i>EXIDIA Fr.</i> . . . . .	313	<i>macrodon Fr.</i> . . . . .	245
<i>glandulosa Fr.</i> . . . . .	314	<i>melaleucum Fr.</i> . . . . .	238
<i>recisa Fr.</i> . . . . .	314	<i>mellem B. &amp; Br.</i> . . . . .	244
<i>saccharina Fr.</i> . . . . .	314	<i>membranaceum Bull.</i> . . . . .	242
<i>truncata Fr.</i> . . . . .	34	<i>multiforme B. &amp; Br.</i> . . . . .	244
<i>FAVULUS Fr.</i> . . . . .	227	<i>nigrum Fr.</i> . . . . .	238
<i>FEMSJONIA Fr.</i> . . . . .	315	<i>niveum Pers.</i> . . . . .	246
<i>FISTULINA Bull.</i> . . . . .	183	<i>nodulosum Fr.</i> . . . . .	245
<i>hepatica Fr.</i> . . . . .	183	<i>ochraceum Pers.</i> . . . . .	241
<i>GOMPHIDIUS Fr.</i> . . . . .	63	<i>repandum Linn.</i> . . . . .	236
<i>glutinosus Fr.</i> . . . . .	63	<i>rufescens Pers.</i> . . . . .	236
<i>gracilis B. &amp; Br.</i> . . . . .	64	<i>scabrosum Fr.</i> . . . . .	234
<i>roseus Fr.</i> . . . . .	64	<i>scrobiculatum Fr.</i> . . . . .	237
<i>stillatus Strauss</i> . . . . .	65	<i>sepultum B. &amp; Br.</i> . . . . .	244
<i>viscidus Fr.</i> . . . . .	64	<i>sordidum Weinm.</i> . . . . .	243
<i>GRANDINIA Fr.</i> . . . . .	255	<i>spatulatum Fr.</i> . . . . .	244
<i>crustosa Fr.</i> . . . . .	256	<i>squalinum Fr.</i> . . . . .	241
<i>granulosa Fr.</i> . . . . .	255	<i>squamosum Schæff.</i> . . . . .	234
<i>mucida Fr.</i> . . . . .	255	<i>Stevensoni B. &amp; Br.</i> . . . . .	245
<i>ocellata Fr.</i> . . . . .	255	<i>stipatum Fr.</i> . . . . .	246
<i>papillosa Fr.</i> . . . . .	256	<i>tomentosum B. &amp; Br.</i> . . . . .	239
<i>GUEPINIA Fr.</i> . . . . .	317	<i>udum Fr.</i> . . . . .	245
<i>peziza Tul.</i> . . . . .	317	<i>variecolor Fr.</i> . . . . .	242
<i>HERICIUM Pers.</i> . . . . .	247	<i>Weinmanni Fr.</i> . . . . .	242
<i>HEXAGONA Poll.</i> . . . . .	226	<i>zonatum Batsch</i> . . . . .	237
<i>HIATULA Fr.</i> . . . . .	322	<i>HYGROPHORUS Fr.</i> . . . . .	70
<i>Wynnæ B. &amp; Br.</i> . . . . .	322	<i>agathosmus Fr.</i> . . . . .	76
<i>HIRNEOLA Fr.</i> . . . . .	315	<i>arbutivus Fr.</i> . . . . .	73
<i>auricula-Judæ Berk.</i> . . . . .	315	<i>calyptræformis B. &amp; Br.</i> . . . . .	89
<i>HYDNUM Linn.</i> . . . . .	233	<i>caprinus Fr.</i> . . . . .	77
<i>alutaceum Fr.</i> . . . . .	243	<i>ceraceus Fr.</i> . . . . .	86
<i>anomalum B. &amp; Br.</i> . . . . .	244	<i>cerasinus Berk.</i> . . . . .	76
<i>argutum Fr.</i> . . . . .	246	<i>chlorophanus Fr.</i> . . . . .	90
<i>aurantiacum Alb. &amp; Schw.</i> . . . . .	236	<i>chrysodon Fr.</i> . . . . .	71
<i>aureum Fr.</i> . . . . .	243	<i>cinereus Fr.</i> . . . . .	79
<i>auriscalpium Linn.</i> . . . . .	239	<i>Clarkii B. &amp; Br.</i> . . . . .	82
<i>bicolor Alb. &amp; Schw.</i> . . . . .	245	<i>coccineus Fr.</i> . . . . .	86
<i>caput-Medusæ Bull.</i> . . . . .	240	<i>Colemannianus Blox.</i> . . . . .	84
<i>cirrhatum Pers.</i> . . . . .	240	<i>conicus Fr.</i> . . . . .	89
<i>compactum Pers.</i> . . . . .	236	<i>cossus Fr.</i> . . . . .	72
<i>coralloides Scop.</i> . . . . .	239	<i>discoides Fr.</i> . . . . .	74
<i>crinale Fr.</i> . . . . .	242	<i>distans Berk.</i> . . . . .	81
<i>cyathiforme Schæff.</i> . . . . .	239	<i>eburneus Fr.</i> . . . . .	71
<i>denticulatum Pers.</i> . . . . .	243	<i>erubescens Fr.</i> . . . . .	73
<i>diversidens Fr.</i> . . . . .	240	<i>foetens Phill.</i> . . . . .	79
<i>erinaceum Bull.</i> . . . . .	240	<i>fornicatus Fr.</i> . . . . .	81
<i>farinaceum Pers.</i> . . . . .	246	<i>fusco-albus Fr.</i> . . . . .	76
		<i>glauco-nitens Fr.</i> . . . . .	91
		<i>glutifer Fr.</i> . . . . .	73
		<i>Houghtoni B. &amp; Br.</i> . . . . .	84
		<i>hypothecus Fr.</i> . . . . .	75
		<i>irrigatus Fr.</i> . . . . .	83
		<i>lacmus Fr.</i> . . . . .	83
		<i>lætus Fr.</i> . . . . .	85
		<i>leporinus Fr.</i> . . . . .	78
		<i>limacinus Fr.</i> . . . . .	74
		<i>livido-albus Fr.</i> . . . . .	77
		<i>mesotephrus B. &amp; Br.</i> . . . . .	77



HYGROPHORUS Fr.—continued.		PAGE	LACTARIUS Fr.—continued.		PAGE
metapodius Fr.		82	mammosus Fr.		107
micaceus B. & Br.		81	marginato-tomentoso B. & Br.		96
miniatus Fr.		87	minimus Smith		112
mollis B. & Br.		87	mitissimus Fr.		110
mucronellus Fr.		87	obnubilus Lasch		112
murinaceus Fr.		91	pallidus Fr.		104
nemoreus Fr.		78	pallidus Saund. & Sm.		105
nitratus Fr.		91	pargamenus Fr.		101
niveus Fr.		80	picinus Fr.		109
obrusseus Fr.		88	piperatus Fr.		102
olivaceo-albus Fr.		75	pubescens Fr.		95
ovinus Fr.		82	pyrogalus Fr.		100
pallidus B. & Br.		79	quietus Fr.		104
penarius Fr.		72	roseo-zonatus Post.		99
pratensis Fr.		78	rufus Fr.		106
psittacinus Fr.		90	scoticus B. & Br.		103
pulverulentus B. & Br.		72	scrobiculatus Fr.		93
puniceus Fr.		88	serifluus Fr.		110
russo-coriaceus B. & Br.		81	squalidus Fr.		100
sciophanus Fr.		84	subdulcis Fr.		111
subradiatus Fr.		83	Terrei B. & Br.		111
turundus Fr.		87	theiogalus Fr.		105
unguinosis Fr.		91	torminosus Fr.		93
ventricosus B. & Br.		80	trivialis Fr.		98
virgineus Fr.		79	turpis Fr.		94
vitellinus Fr.		85	uvidus Fr.		98
Wynnii B. & Br.		80	vellereus Fr.		102
			vietus Fr.		105
			volemus Fr.		109
			zonarius Fr.		96
HYMENOCHÆTE Lev.			LENTINUS Fr.		
corrugata Berk.		281	adhærens Fr.		156
rubiginosa Lev.		269	cochleatus Fr.		156
Stevensoni B. & Br.		269	Dunalii Fr.		154
tabacina Lev.		269	fimbriatus Curr.		157
IRPEX Fr.			flabelliformis Fr.		158
carneus Fr.		248	leontopodius Schulz.		155
deformis Fr.		250	lepideus Fr.		155
fusco-violaceus Fr.		249	pulverulentus Fr.		155
Johnstoni Berk.		249	resinaceus Fr.		156
lacteus Eng. Fl.		250	scoticus B. & Br.		157
obliquus Fr.		250	tigrinus Fr.		164
pendulus Fr.		249	vulpinus Fr.		157
spathulatus Fr.		250			
KNEIFFIA Fr.			LENZITES Fr.		
setigera Fr.		257	abietina Fr.		163
subgelatinosa B. & Br.		258	betulina Fr.		163
LACTARIUS Fr.			flaccida Fr.		164
acris Fr.		101	sæpiaria Fr.		164
blennius Fr.		97			
camphoratus Fr.		111	MARASMIUS Fr.		
capsicum Schulz.		100	alliaceus Fr.		148
chrysorheus Fr.		101	amadelphus Fr.		148
cilicoides Fr.		94	androsaceus Fr.		150
circellatus Fr.		98	angulatus B. & Br.		147
controversus Fr.		95	archyropus Fr.		145
cyathula Fr.		106	Broomei Berk.		153
deliciosus Fr.		103	calopus Fr.		146
exsuccus Smith		103	caulicinalis Fr.		149
flexuosus Fr.		99	Curreyi B. & Br.		151
fuliginosus Fr.		108	epichloe Fr.		152
glycosmus Fr.		107	epiphyllus Fr.		152
helvus Fr.		107	erythropus Fr.		144
hyssinus Fr.		97	foetidus Fr.		147
ichoratus Fr.		110	fusco-purpureus Fr.		143
insulsius Fr.		96	gramineus Lev.		152
lilacinus Lasch		109	graminum Berk.		150

MARASMIUS Fr.—continued.	PAGE	PAXILLUS Fr.—continued.	PAGE
Hudsoni Fr. . . . .	151	involutus Fr. . . . .	68
impudicus Fr. . . . .	145	lepista Fr. . . . .	66
insititius Fr. . . . .	151	leptopus Fr. . . . .	68
languidus Fr. . . . .	147	panæolus Fr. . . . .	67
oreades Fr. . . . .	142	panuoides Fr. . . . .	69
perforans Fr. . . . .	151	paradoxus Berk. . . . .	67
peronatus Fr. . . . .	141	spilomæolus Fr. . . . .	67
porreus Fr. . . . .	141		
ramealis Fr. . . . .	148	PHLEBIA Fr. . . . .	253
rotula Fr. . . . .	149	contorta Fr. . . . .	254
saccharinus Fr. . . . .	152	lirellosa Pers. . . . .	254
scorodonius Fr. . . . .	146	merismoides Fr. . . . .	253
scorteus Fr. . . . .	142	radiata Fr. . . . .	253
splachnoides Fr. . . . .	150	vaga Fr. . . . .	254
spodolencus Berk. . . . .	153		
Stephensii Berk. . . . .	144	PISTILLARIA Fr. . . . .	307
terginus Fr. . . . .	143	culmigena Fr. . . . .	307
torquescens Quel. . . . .	145	furcata Smith. . . . .	307
urens Fr. . . . .	140	nicans Fr. . . . .	307
Vaillantii Fr. . . . .	146	quisquiliaris Fr. . . . .	307
varicosus Fr. . . . .	143	puberula Berk. . . . .	303
Wynnei B. & Br. . . . .	144	pusilla Fr. . . . .	308
MERULIUS Fr. . . . .	227	POLYPORUS Fr. . . . .	183
aurantiacus Klotzsch . . . . .	228	abietinus Fr. . . . .	211
Carmichaelianus Berk. . . . .	230	acanthoides Fr. . . . .	195
corium Fr. . . . .	228	adiposus B. & Br. . . . .	201
himantoides Fr. . . . .	228	adustus Fr. . . . .	200
lachrymans Fr. . . . .	230	alligatus Fr. . . . .	195
laticolor B. & Br. . . . .	228	alutaceus Fr. . . . .	197
molluscus Fr. . . . .	229	amorphus Fr. . . . .	201
pallens Berk. . . . .	229	aneirinus Sommerf. . . . .	214
porinoides Fr. . . . .	229	annosus Fr. . . . .	208
pulverulentus Fr. . . . .	230	applanatus Fr. . . . .	204
rufus Pers. . . . .	229	armeniaceus Berk. . . . .	215
serpens Tode. . . . .	229	bathyporus Rostk. . . . .	220
tremellosus Schrad. . . . .	227	betulinus Fr. . . . .	203
		blepharistoma B. & Br. . . . .	220
MICROCERA Desm. . . . .	308	bombycinus Fr. . . . .	214
		borealis Fr. . . . .	202
MUCRONELLA Fr. . . . .	258	brumalis Fr. . . . .	188
		cæsius Fr. . . . .	198
NÆMATELIA Fr. . . . .	316	callosus Fr. . . . .	217
encephala Fr. . . . .	316	carneus Nees . . . . .	208
nucleata Fr. . . . .	316	cerebrinus B. & Br. . . . .	197
virescens Corda . . . . .	316	cervinus Pers. . . . .	223
		chioneus Fr. . . . .	197
NYCTALIS Fr. . . . .	137	cinctus Berk. . . . .	215
asterophora Fr. . . . .	138	collabefactus B. & Br. . . . .	218
caliginosa Smith . . . . .	138	conchatus Fr. . . . .	206
parasitica Fr. . . . .	138	connatus Fr. . . . .	207
		contiguus Fr. . . . .	212
ODONTIA Fr. . . . .	256	crispus Fr. . . . .	200
barba-Jovis Fr. . . . .	256	cristatus Fr. . . . .	194
fimbriata Pers. . . . .	257	cryptarum Fr. . . . .	209
		cuticularis Fr. . . . .	202
PANUS Fr. . . . .	158	cytisinus Berk. . . . .	207
conchatus Fr. . . . .	159	destructor Fr. . . . .	199
farinaceus Schum. . . . .	160	dryadeus Fr. . . . .	202
patellaris Fr. . . . .	160	elegans Fr. . . . .	192
Stevensoni B. & Br. . . . .	160	epileucus Fr. . . . .	196
stipticus Fr. . . . .	160	farinellus Fr. . . . .	220
torulosus Fr. . . . .	159	ferruginosus Fr. . . . .	212
		fibula Fr. . . . .	210
PAXILLUS Fr. . . . .	65	floccipes Rostk. . . . .	188
atro-tomentosus Fr. . . . .	68	fomentarius Fr. . . . .	204
fagi B. & Br. . . . .	69	fragilis Fr. . . . .	198
giganteus Fr. . . . .	66	fraxineus Fr. . . . .	207

POLYPORUS <i>Fr.</i> —continued.	PAGE	POLYPORUS <i>Fr.</i> —continued.	PAGE
<i>frondosus Fr.</i> . . . . .	193	<i>sulphureus Fr.</i> . . . . .	195
<i>fulvus Fr.</i> . . . . .	205	<i>terrestris Fr.</i> . . . . .	216
<i>fumosus Fr.</i> . . . . .	200	<i>trabeus Rostk.</i> . . . . .	198
<i>fuscidulus Fr.</i> . . . . .	188	<i>ulnarius Fr.</i> . . . . .	207
<i>giganteus Fr.</i> . . . . .	194	<i>umbellatus Fr.</i> . . . . .	193
<i>gilvus Schw.</i> . . . . .	199	<i>umbrinus Fr.</i> . . . . .	211
<i>Gordoniensis B. &amp; Br.</i> . . . . .	219	<i>Vallantii Fr.</i> . . . . .	219
<i>gossypinus Lev.</i> . . . . .	209	<i>vaporarius Fr.</i> . . . . .	219
<i>Herbergii Rostk.</i> . . . . .	195	<i>variegatus Secr.</i> . . . . .	208
<i>heteroclitus Fr.</i> . . . . .	196	<i>varius Fr.</i> . . . . .	191
<i>hirsutus Fr.</i> . . . . .	210	<i>vegetus Fr.</i> . . . . .	203
<i>hispidus Fr.</i> . . . . .	201	<i>velutinus Fr.</i> . . . . .	210
<i>hybridus B. &amp; Br.</i> . . . . .	221	<i>versicolor Fr.</i> . . . . .	210
<i>hymenocystis B. &amp; Br.</i> . . . . .	220	<i>violaceus Fr.</i> . . . . .	212
<i>igniarius Fr.</i> . . . . .	205	<i>viridans Berk.</i> . . . . .	216
<i>incarnatus Fr.</i> . . . . .	213	<i>vitreus Fr.</i> . . . . .	217
<i>intybaceus Fr.</i> . . . . .	193	<i>vulgaris Fr.</i> . . . . .	218
<i>Keithii B. &amp; Br.</i> . . . . .	201	<i>Wynnæ B. &amp; Br.</i> . . . . .	211
<i>lacteus Fr.</i> . . . . .	197		
<i>Læstadii Fr. &amp; Berk.</i> . . . . .	214	<b>POROTHELIUM <i>Fr.</i></b> . . . . .	230
<i>lentus Berk.</i> . . . . .	187	<i>confusum B. &amp; Br.</i> . . . . .	231
<i>leptocephalus Fr.</i> . . . . .	188	<i>Friesii Mont.</i> . . . . .	231
<i>leucomelas Fr.</i> . . . . .	187	<i>Keithii B. &amp; Br.</i> . . . . .	231
<i>lucidus Fr.</i> . . . . .	193	<i>Stevensoni B. &amp; Br.</i> . . . . .	231
<i>medulla-panis Fr.</i> . . . . .	216		
<i>melanopus Fr.</i> . . . . .	190	<b>PTERULA <i>Fr.</i></b> . . . . .	303
<i>micans Fr.</i> . . . . .	213	<i>multifida Fr.</i> . . . . .	304
<i>Michellii Fr.</i> . . . . .	190	<i>subulata Fr.</i> . . . . .	303
<i>mollis Fr.</i> . . . . .	198		
<i>molluscus Fr.</i> . . . . .	218	<b>RADULUM <i>Fr.</i></b> . . . . .	251
<i>mucidus Fr.</i> . . . . .	217	<i>aterrimum Fr.</i> . . . . .	253
<i>nidulans Fr.</i> . . . . .	199	<i>corallinum B. &amp; Br.</i> . . . . .	252
<i>nigricans Fr.</i> . . . . .	204	<i>deglubens B. &amp; Br.</i> . . . . .	252
<i>nitidus Fr.</i> . . . . .	214	<i>epileucum B. &amp; Br.</i> . . . . .	252
<i>nummularius Fr.</i> . . . . .	192	<i>erectum Sow.</i> . . . . .	253
<i>obducens Pers.</i> . . . . .	217	<i>fagineum Fr.</i> . . . . .	252
<i>pallescens Fr.</i> . . . . .	197	<i>orbiculare Fr.</i> . . . . .	251
<i>pectinatus Klotzsch.</i> . . . . .	205	<i>quercinum Fr.</i> . . . . .	251
<i>perennis Fr.</i> . . . . .	189	<i>tomentosum Fr.</i> . . . . .	252
<i>petaloides Fr.</i> . . . . .	192		
<i>picipes Fr.</i> . . . . .	191	<b>RUSSULA <i>Pers.</i></b> . . . . .	113
<i>polymorphus Rostk.</i> . . . . .	209	<i>adusta Fr.</i> . . . . .	114
<i>populinus Fr.</i> . . . . .	208	<i>alutacea Fr.</i> . . . . .	129
<i>purpureus Fr.</i> . . . . .	212	<i>aurata Fr.</i> . . . . .	128
<i>quercinus Fr.</i> . . . . .	203	<i>chamæleontina Fr.</i> . . . . .	130
<i>radiatus Fr.</i> . . . . .	209	<i>consobrina Fr.</i> . . . . .	123
<i>radula Fr.</i> . . . . .	219	<i>cutefracta Cke.</i> . . . . .	121
<i>ramentaceus B. &amp; Br.</i> . . . . .	214	<i>cyanoxantha Fr.</i> . . . . .	122
<i>Rennyi B. &amp; Br.</i> . . . . .	215	<i>decolorans Fr.</i> . . . . .	128
<i>reticulatus Fr.</i> . . . . .	220	<i>delica Fr.</i> . . . . .	115
<i>rhodellus Fr.</i> . . . . .	213	<i>depallens Fr.</i> . . . . .	117
<i>ribis Fr.</i> . . . . .	206	<i>drimeia Cke.</i> . . . . .	118
<i>roseus Fr.</i> . . . . .	206	<i>Du Portii Phill.</i> . . . . .	119
<i>Rostkovii Fr.</i> . . . . .	191	<i>elephantina Fr.</i> . . . . .	115
<i>rufescens Fr.</i> . . . . .	189	<i>emetica Fr.</i> . . . . .	125
<i>rufus Fr.</i> . . . . .	213	<i>fellea Fr.</i> . . . . .	124
<i>rutilans Fr.</i> . . . . .	199	<i>foetens Fr.</i> . . . . .	124
<i>rutilans Saund. &amp; Sm.</i> . . . . .	199	<i>fragilis Fr.</i> . . . . .	126
<i>salicinus Fr.</i> . . . . .	206	<i>furcata Fr.</i> . . . . .	116
<i>salignus Fr.</i> . . . . .	196	<i>galochroa Fr.</i> . . . . .	123
<i>sanguinolentus Fr.</i> . . . . .	218	<i>heterophylla Fr.</i> . . . . .	123
<i>Schweinizii Fr.</i> . . . . .	189	<i>integra Fr.</i> . . . . .	127
<i>secernibilis B. &amp; Br.</i> . . . . .	219	<i>lactea Fr.</i> . . . . .	118
<i>spumeus Fr.</i> . . . . .	202	<i>lepida Fr.</i> . . . . .	119
<i>squamosus Fr.</i> . . . . .	190	<i>Linnæi Fr.</i> . . . . .	120
<i>Stephensti B. &amp; Br.</i> . . . . .	224	<i>lutea Fr.</i> . . . . .	129
<i>subfusco-flavidus Rostk.</i> . . . . .	215	<i>nauseosa Fr.</i> . . . . .	130
<i>subgelatinosus B. &amp; Br.</i> . . . . .	216	<i>nigricans Fr.</i> . . . . .	114

## RUSSULA Pers.—continued.

	PAGE
nitida Fr. . . . .	129
ochroleuca Fr. . . . .	126
olivacea Fr. . . . .	121
pectinata Fr. . . . .	126
Queletii Fr. . . . .	125
rosea Fr. . . . .	117
rubra Fr. . . . .	120
sanguinea Fr. . . . .	116
sardonia Fr. . . . .	117
semicrema Fr. . . . .	115
sororia Larbr. . . . .	124
subfoetens Smith . . . . .	124
vesca Fr. . . . .	122
veternosa Fr. . . . .	127
virescens Fr. . . . .	119
vitellina Fr. . . . .	130
xerampelina Fr. . . . .	121

SCHIZOPHYLLUM Fr. . . . .	162
commune Fr. . . . .	162

SISTOTREMA Pers. . . . .	248
confluens Pers. . . . .	248

SOLENIA Hoffm. . . . .	232
ochracea Hoffm. . . . .	232

SPARASSIS Fr. . . . .	289
crispa Fr. . . . .	289

STEREUM Pers. . . . .	267
acerinum Pers. . . . .	271
avellanum Fr. . . . .	270
frustulosum Fr. . . . .	270
hirsutum Fr. . . . .	268
lauro-cerasi Berk. . . . .	270
ochroleucum Fr. . . . .	267
pinii Fr. . . . .	270
purpureum Pers. . . . .	268
rubiginosum Fr. . . . .	269
rufo-hispidum Stev. . . . .	269
rufum Fr. . . . .	271
rugosum Fr. . . . .	270
sanguinolentum Fr. . . . .	269
spadiceum Fr. . . . .	268
stratosum B. & Br. . . . .	271
tabacinum Fr. . . . .	269
vorticolum Fr. . . . .	268

STROBILOMYCES Berk. . . . .	182
strobilaceus Berk. . . . .	182

THELEPHORA Ehrb. . . . .	261
anthocephala Fr. . . . .	263
biennis Fr. . . . .	264
caesia Pers. . . . .	266
caryophyllea Pers. . . . .	262
clavularis Fr. . . . .	263
cristata Fr. . . . .	265
crustacea Schum. . . . .	266
digitata Fr. . . . .	263
fastidiosa Fr. . . . .	265

## THELEPHORA Ehrb.—continued.

	PAGE
intybacea Pers. . . . .	264
laciniata Pers. . . . .	264
mollissima Pers. . . . .	265
multizonata B. & Br. . . . .	262
palmata Fr. . . . .	263
sebacea Pers. . . . .	265
Sowerbeii B. & Br. . . . .	261
tuberosa Fr. . . . .	262
terrestris Ehrb. . . . .	264
undulata Fr. . . . .	262

TRAMETES Fr. . . . .	221
Bulliardii Fr. . . . .	222
gibbosa Fr. . . . .	222
inodora Fr. . . . .	223
mollis Fr. . . . .	223
odora Fr. . . . .	222
pini Fr. . . . .	221
serpens Fr. . . . .	223
suaveolens Fr. . . . .	222
Terrei B. & Br. . . . .	223

TREMELLA Dill. . . . .	309
albida Huds. . . . .	311
epigæa B. & Br. . . . .	311
fimbriata Pers. . . . .	309
foliacea Pers. . . . .	310
foliicola Fekl. . . . .	313
frondosa Fr. . . . .	310
indecorata Sommerf. . . . .	312
intumescens Eng. Bot. . . . .	311
lutescens Pers. . . . .	310
mesenterica Retz. . . . .	310
moriformis Eng. Bot. . . . .	312
torta Berk. . . . .	313
tubercularia Berk. . . . .	312
versicolor Berk. . . . .	313
vesicaria Eng. Bot. . . . .	311
violacea Relh. . . . .	312
viscosa Berk. . . . .	311

TREMELLODON Pers. . . . .	247
gelatinosum Pers. . . . .	247

TROGIA Fr. . . . .	162
crispa Fr. . . . .	162

TYPHULA Pers. . . . .	304
erythropus Fr. . . . .	304
filiformis Fr. . . . .	306
Grevillei Fr. . . . .	305
gracilis Berk. & Desm. . . . .	306
gracillima White . . . . .	306
gyrans Fr. . . . .	305
incarnata Lasch . . . . .	305
musciola Fr. . . . .	305
phacorrhiza Fr. . . . .	304
translucens B. & Br. . . . .	306

XEROTUS Fr. . . . .	161
degener Fr. . . . .	161



**THE LIBRARY  
UNIVERSITY OF CALIFORNIA  
San Francisco Medical Center**

**THIS BOOK IS DUE ON THE LAST DATE STAMPED BELOW**

Books not returned on time are subject to fines according to the Library Lending Code.

Books not in demand may be renewed if application is made before expiration of loan period.

172552

